

Mark Scheme (Results)

January 2011

GCE

GCE Core Mathematics C3 (6665) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M marks:** method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A marks:** Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B marks** are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod - benefit of doubt
- ft - follow through
- the symbol \checkmark will be used for correct ft
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw - ignore subsequent working
- awrt - answers which round to
- SC: special case
- oe - or equivalent (and appropriate)
- dep - dependent
- indep - independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- \square The second mark is dependent on gaining the first mark

January 2011
Core Mathematics C3 6665
Mark Scheme

Question Number	Scheme	Marks
1.		
(a)	$7\cos x - 24\sin x = R\cos(x + \alpha)$ $7\cos x - 24\sin x = R\cos x \cos \alpha - R\sin x \sin \alpha$ <p>Equate $\cos x$: $7 = R\cos \alpha$ Equate $\sin x$: $24 = R\sin \alpha$</p> $R = \sqrt{7^2 + 24^2} = 25 \qquad R = 25$ $\tan \alpha = \frac{24}{7} \Rightarrow \alpha = 1.287002218...^{\circ}$ <p style="text-align: right;">$\tan \alpha = \frac{24}{7}$ or $\tan \alpha = \frac{7}{24}$ awrt 1.287</p> <p>Hence, $7\cos x - 24\sin x = 25\cos(x + 1.287)$</p>	<p>B1</p> <p>M1 A1</p> <p>(3)</p>
(b)	<p>Minimum value = <u>-25</u></p> <p style="text-align: right;">-25 or -R</p>	<p>B1ft</p> <p>(1)</p>
(c)	$7\cos x - 24\sin x = 10$ $25\cos(x + 1.287) = 10$ $\cos(x + 1.287) = \frac{10}{25}$ <p>PV = 1.159279481...^c or 66.42182152...^o</p> <p>So, $x + 1.287 = \{1.159279...^{\circ}, 5.123906...^{\circ}, 7.442465...^{\circ}\}$</p> <p>gives, $x = \{3.836906..., 6.155465...\}$</p>	<p>$\cos(x \pm \text{their } \alpha) = \frac{10}{(\text{their } R)}$ M1</p> <p>For applying $\cos^{-1}\left(\frac{10}{\text{their } R}\right)$ M1</p> <p>either $2\pi +$ or $-$ their PV^c or $360^{\circ} +$ or $-$ their PV^o M1</p> <p>awrt 3.84 OR 6.16 A1 awrt 3.84 AND 6.16 A1</p> <p>(5) [9]</p>


Question Number	Scheme	Marks
2.		
(a)	$\frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)}$ $= \frac{(4x-1)(2x-1) - 3}{2(x-1)(2x-1)}$ $= \frac{8x^2 - 6x - 2}{\{2(x-1)(2x-1)\}}$ $= \frac{2(x-1)(4x+1)}{\{2(x-1)(2x-1)\}}$ $= \frac{4x+1}{2x-1}$	<p>An attempt to form a single fraction</p> <p>Simplifies to give a correct quadratic numerator over a correct quadratic denominator</p> <p>An attempt to factorise a 3 term quadratic numerator</p> <p>M1</p> <p>A1 aef</p> <p>M1</p> <p>A1</p> <p>(4)</p>
(b)	$f(x) = \frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)} - 2, \quad x > 1$ $f(x) = \frac{(4x+1)}{(2x-1)} - 2$ $= \frac{(4x+1) - 2(2x-1)}{(2x-1)}$ $= \frac{4x+1-4x+2}{(2x-1)}$ $= \frac{3}{(2x-1)}$	<p>An attempt to form a single fraction</p> <p>Correct result</p> <p>M1</p> <p>A1 *</p> <p>(2)</p>
(c)	$f(x) = \frac{3}{(2x-1)} = 3(2x-1)^{-1}$ $f'(x) = 3(-1)(2x-1)^{-2}(2)$ $f'(2) = \frac{-6}{9} = -\frac{2}{3}$	<p>$\pm k(2x-1)^{-2}$</p> <p>Either $\frac{-6}{9}$ or $-\frac{2}{3}$</p> <p>M1</p> <p>A1 aef</p> <p>A1</p> <p>(3)</p> <p>[9]</p>


Question Number	Scheme	Marks
3.	$2 \cos 2\theta = 1 - 2 \sin \theta$ $2(1 - 2 \sin^2 \theta) = 1 - 2 \sin \theta$ $2 - 4 \sin^2 \theta = 1 - 2 \sin \theta$ $4 \sin^2 \theta - 2 \sin \theta - 1 = 0$ $\sin \theta = \frac{2 \pm \sqrt{4 - 4(4)(-1)}}{8}$ PVs: $\alpha_1 = 54^\circ$ or $\alpha_2 = -18^\circ$ $\theta = \{54, 126, 198, 342\}$	<p>Substitutes either $1 - 2 \sin^2 \theta$ or $2 \cos^2 \theta - 1$ or $\cos^2 \theta - \sin^2 \theta$ for $\cos 2\theta$. M1</p> <p>Forms a “quadratic in sine” = 0 M1(*)</p> <p>Applies the quadratic formula See notes for alternative methods. M1</p> <p>Any one correct answer 180-their pv All four solutions correct. A1 dM1(*) A1</p> <p>[6]</p>

Question Number	Scheme	Marks
4.		
(a)	$\theta = 20 + Ae^{-kt}$ (eqn *) $\{t = 0, \theta = 90 \Rightarrow\} \quad 90 = 20 + Ae^{-k(0)} \quad \text{Substitutes } t = 0 \text{ and } \theta = 90 \text{ into eqn } *$ $90 = 20 + A \Rightarrow \underline{A = 70} \quad \underline{A = 70}$	M1 A1 (2)
(b)	$\theta = 20 + 70e^{-kt}$ $\{t = 5, \theta = 55 \Rightarrow\} \quad 55 = 20 + 70e^{-k(5)} \quad \text{Substitutes } t = 5 \text{ and } \theta = 55 \text{ into eqn } *$ $\frac{35}{70} = e^{-5k} \quad \text{and rearranges eqn } * \text{ to make } e^{\pm 5k} \text{ the subject.}$ $\ln\left(\frac{35}{70}\right) = -5k \quad \text{Takes 'lns' and proceeds to make '}\pm 5k\text{' the subject.}$ $-5k = \ln\left(\frac{1}{2}\right)$ $-5k = \ln 1 - \ln 2 \Rightarrow -5k = -\ln 2 \Rightarrow \underline{k = \frac{1}{5}\ln 2} \quad \text{Convincing proof that } k = \frac{1}{5}\ln 2$	M1 dM1 A1 * (3)
(c)	$\theta = 20 + 70e^{-\frac{1}{5}t\ln 2}$ $\frac{d\theta}{dt} = -\frac{1}{5}\ln 2 \cdot (70)e^{-\frac{1}{5}t\ln 2} \quad \pm \alpha e^{-kt} \text{ where } k = \frac{1}{5}\ln 2$ $-14\ln 2 e^{-\frac{1}{5}t\ln 2}$ When $t = 10$, $\frac{d\theta}{dt} = -14\ln 2 e^{-2\ln 2}$ $\frac{d\theta}{dt} = -\frac{7}{2}\ln 2 = -2.426015132...$ Rate of decrease of $\theta = 2.426^\circ \text{C/min}$ (3 dp.) awrt ± 2.426	M1 A1 oe A1 (3) [8]

Question Number	Scheme	Marks
5.		
(a)	<p>Crosses x-axis $\Rightarrow f(x) = 0 \Rightarrow (8 - x)\ln x = 0$</p> <p>Either $(8 - x) = 0$ or $\ln x = 0 \Rightarrow x = 8, 1$</p> <p>Coordinates are $A(1, 0)$ and $B(8, 0)$.</p>	<p>Either one of $\{x\}=1$ OR $x=\{8\}$ B1</p> <p>Both $A(1, \{0\})$ and $B(8, \{0\})$ B1</p> <p>(2)</p>
(b)	<p>Apply product rule: $\left\{ \begin{array}{l} u = (8 - x) \quad v = \ln x \\ \frac{du}{dx} = -1 \quad \frac{dv}{dx} = \frac{1}{x} \end{array} \right\}$</p> <p>$f'(x) = -\ln x + \frac{8-x}{x}$</p>	<p>$vu' + uv'$ M1</p> <p>Any one term correct A1</p> <p>Both terms correct A1</p> <p>(3)</p>
(c)	<p>$f'(3.5) = 0.032951317...$</p> <p>$f'(3.6) = -0.058711623...$</p> <p>Sign change (and as $f'(x)$ is continuous) therefore the x-coordinate of Q lies between 3.5 and 3.6.</p>	<p>Attempts to evaluate both $f'(3.5)$ and $f'(3.6)$ M1</p> <p>both values correct to at least 1 sf, sign change and conclusion A1</p> <p>(2)</p>
(d)	<p>At Q, $f'(x) = 0 \Rightarrow -\ln x + \frac{8-x}{x} = 0$</p> <p>$\Rightarrow -\ln x + \frac{8}{x} - 1 = 0$</p> <p>$\Rightarrow \frac{8}{x} = \ln x + 1 \Rightarrow 8 = x(\ln x + 1)$</p> <p>$\Rightarrow x = \frac{8}{\ln x + 1}$ (as required)</p>	<p>Setting $f'(x) = 0$. M1</p> <p>Splitting up the numerator and proceeding to $x=$ M1</p> <p>For correct proof. No errors seen in working. A1</p> <p>(3)</p>

Question Number	Scheme	Marks
(e)	<p>Iterative formula: $x_{n+1} = \frac{8}{\ln x_n + 1}$</p> <p>$x_1 = \frac{8}{\ln(3.55) + 1}$</p> <p>$x_1 = 3.528974374...$</p> <p>$x_2 = 3.538246011...$</p> <p>$x_3 = 3.534144722...$</p> <p>$x_1 = 3.529, x_2 = 3.538, x_3 = 3.534, \text{ to 3 dp.}$</p>	<p>An attempt to substitute $x_0 = 3.55$ into the iterative formula. Can be implied by $x_1 = 3.528(97)...$ Both $x_1 = \text{awrt } 3.529$ and $x_2 = \text{awrt } 3.538$</p> <p>M1</p> <p>A1</p> <p>x_1, x_2, x_3 all stated correctly to 3 dp</p> <p>A1</p> <p>(3) [13]</p>

Question Number	Scheme	Marks
6.		
(a)	$y = \frac{3-2x}{x-5} \Rightarrow y(x-5) = 3-2x$ <p>Attempt to make x (or swapped y) the subject</p> $xy - 5y = 3 - 2x$ $\Rightarrow xy + 2x = 3 + 5y \Rightarrow x(y+2) = 3 + 5y$ <p>Collect x terms together and factorise.</p> $\Rightarrow x = \frac{3+5y}{y+2} \therefore f^{-1}(x) = \frac{3+5x}{x+2}$	<p>M1</p> <p>M1</p> <p>A1 oe (3)</p>
(b)	<p>Range of g is $-9 \leq g(x) \leq 4$ or $-9 \leq y \leq 4$</p> <p>Correct Range</p>	<p>B1 (1)</p>
(c)	<p>$g g(2) = g(0) = -6$, from sketch.</p>	<p>Deduces that $g(2)$ is 0. Seen or implied.</p> <p>-6</p> <p>M1</p> <p>A1 (2)</p>
(d)	<p>$fg(8) = f(4)$</p> $= \frac{3-4(2)}{4-5} = \frac{-5}{-1} = 5$	<p>Correct order g followed by f</p> <p>5</p> <p>M1</p> <p>A1 (2)</p>
(e)(i)	 <p>Correct shape</p> <p>$(2, \{0\}), (\{0\}, 6)$</p>	<p>B1</p> <p>B1</p>

Question Number	Scheme	Marks
(e)(ii)	 <p>Correct shape</p> <p>Graph goes through $(\{0\}, 2)$ and $(-6, \{0\})$ which are marked.</p>	<p>B1</p> <p>B1</p> <p>(4)</p>
(f)	<p>Domain of g^{-1} is $-9 \leq x \leq 4$</p> <p>Either correct answer or a follow through from part (b) answer</p>	<p>B1 $\sqrt{\quad}$</p> <p>(1)</p> <p>[13]</p>

Question Number	Scheme	Marks
7	<p>(a) $y = \frac{3 + \sin 2x}{2 + \cos 2x}$</p> <p>Apply quotient rule:</p> $\left\{ \begin{array}{l} u = 3 + \sin 2x \quad v = 2 + \cos 2x \\ \frac{du}{dx} = 2 \cos 2x \quad \frac{dv}{dx} = -2 \sin 2x \end{array} \right\}$ $\frac{dy}{dx} = \frac{2 \cos 2x(2 + \cos 2x) - (-2 \sin 2x)(3 + \sin 2x)}{(2 + \cos 2x)^2}$ $= \frac{4 \cos 2x + 2 \cos^2 2x + 6 \sin 2x + 2 \sin^2 2x}{(2 + \cos 2x)^2}$ $= \frac{4 \cos 2x + 6 \sin 2x + 2(\cos^2 2x + \sin^2 2x)}{(2 + \cos 2x)^2}$ $= \frac{4 \cos 2x + 6 \sin 2x + 2}{(2 + \cos 2x)^2} \quad (\text{as required})$	<p>Applying $\frac{u'v - uv'}{v^2}$ M1</p> <p>Any one term correct on the numerator A1</p> <p>Fully correct (unsimplified). A1</p> <p>For correct proof with an understanding that $\cos^2 2x + \sin^2 2x = 1$. A1*</p> <p>No errors seen in working. A1*</p> <p>(4)</p>
(b)	<p>When $x = \frac{\pi}{2}$, $y = \frac{3 + \sin \pi}{2 + \cos \pi} = \frac{3}{1} = 3$</p> <p>At $(\frac{\pi}{2}, 3)$, $m(\mathbf{T}) = \frac{6 \sin \pi + 4 \cos \pi + 2}{(2 + \cos \pi)^2} = \frac{-4 + 2}{1^2} = -2$</p> <p>Either \mathbf{T}: $y - 3 = -2(x - \frac{\pi}{2})$ or $y = -2x + c$ and $3 = -2(\frac{\pi}{2}) + c \Rightarrow c = 3 + \pi$; \mathbf{T}: $y = -2x + (\pi + 3)$</p>	<p>$y = 3$ B1</p> <p>$m(\mathbf{T}) = -2$ B1</p> <p>$y - y_1 = m(x - \frac{\pi}{2})$ with 'their TANGENT gradient' and their y_1; M1 or uses $y = mx + c$ with 'their TANGENT gradient';</p> <p>$y = -2x + \pi + 3$ A1</p> <p>(4) [8]</p>

Question Number	Scheme	Marks
8.		
(a)	$y = \sec x = \frac{1}{\cos x} = (\cos x)^{-1}$ $\frac{dy}{dx} = -1(\cos x)^{-2}(-\sin x)$ $\frac{dy}{dx} = \left\{ \frac{\sin x}{\cos^2 x} \right\} = \left(\frac{1}{\cos x} \right) \left(\frac{\sin x}{\cos x} \right) = \underline{\underline{\sec x \tan x}}$	<p>Writes $\sec x$ as $(\cos x)^{-1}$ and gives</p> $\frac{dy}{dx} = \pm((\cos x)^{-2}(\sin x))$ <p>$-1(\cos x)^{-2}(-\sin x)$ or $(\cos x)^{-2}(\sin x)$</p> <p>Convincing proof. Must see both <u>underlined steps</u>.</p> <p>M1 A1 A1 AG (3)</p>
(b)	$x = \sec 2y, \quad y \neq (2n+1)\frac{\pi}{4}, n \in \mathbb{Z}.$ $\frac{dx}{dy} = 2 \sec 2y \tan 2y$	<p>$K \sec 2y \tan 2y$ $2 \sec 2y \tan 2y$</p> <p>M1 A1 (2)</p>
(c)	$\frac{dy}{dx} = \frac{1}{2 \sec 2y \tan 2y}$ $\frac{dy}{dx} = \frac{1}{2x \tan 2y}$ $1 + \tan^2 A = \sec^2 A \Rightarrow \tan^2 2y = \sec^2 2y - 1$ <p>So $\tan^2 2y = x^2 - 1$</p> $\frac{dy}{dx} = \frac{1}{2x\sqrt{(x^2-1)}}$	<p>Applies $\frac{dy}{dx} = \frac{1}{\left(\frac{dx}{dy}\right)}$</p> <p>Substitutes x for $\sec 2y$.</p> <p>Attempts to use the identity $1 + \tan^2 A = \sec^2 A$</p> <p>$\frac{dy}{dx} = \frac{1}{2x\sqrt{(x^2-1)}}$</p> <p>M1 M1 M1 A1 (4) [9]</p>

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481

Email publications@linneydirect.com

Order Code US026238 January 2011

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH