



1.




Figure 1

Figure 1 shows part of the curve with equation  $y = \sqrt{0.75 + \cos^2 x}$ . The finite region  $R$ , shown shaded in Figure 1, is bounded by the curve, the  $y$ -axis, the  $x$ -axis and the line with equation  $x = \frac{\pi}{3}$ .

(a) Complete the table with values of  $y$  corresponding to  $x = \frac{\pi}{6}$  and  $x = \frac{\pi}{4}$ .

$x$	0	$\frac{\pi}{12}$	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$
$y$	1.3229	1.2973			1

(2)

(b) Use the trapezium rule

(i) with the values of  $y$  at  $x = 0$ ,  $x = \frac{\pi}{6}$  and  $x = \frac{\pi}{3}$  to find an estimate of the area of  $R$ .

Give your answer to 3 decimal places.

(ii) with the values of  $y$  at  $x = 0$ ,  $x = \frac{\pi}{12}$ ,  $x = \frac{\pi}{6}$ ,  $x = \frac{\pi}{4}$  and  $x = \frac{\pi}{3}$  to find a

further estimate of the area of  $R$ . Give your answer to 3 decimal places.

(6)























4. A curve  $C$  has parametric equations

$$x = \sin^2 t, \quad y = 2 \tan t, \quad 0 \leq t < \frac{\pi}{2}$$

(a) Find  $\frac{dy}{dx}$  in terms of  $t$ .

**(4)**

The tangent to  $C$  at the point where  $t = \frac{\pi}{3}$  cuts the  $x$ -axis at the point  $P$ .

(b) Find the  $x$ -coordinate of  $P$ .

**(6)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave blank

Question 4 continued

Lined area for writing the answer to Question 4 continued.











Leave blank

**Question 5 continued**

Ruled area for writing the answer to Question 5.







6.  $f(\theta) = 4 \cos^2 \theta - 3 \sin^2 \theta$

(a) Show that  $f(\theta) = \frac{1}{2} + \frac{7}{2} \cos 2\theta$ . (3)

(b) Hence, using calculus, find the exact value of  $\int_0^{\frac{\pi}{2}} \theta f(\theta) d\theta$ . (7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





Leave blank

Question 6 continued

A large rectangular area containing numerous horizontal lines for writing, intended for the student's answer to Question 6.





7. The line  $l_1$  has equation  $\mathbf{r} = \begin{pmatrix} 2 \\ 3 \\ -4 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ , where  $\lambda$  is a scalar parameter.

The line  $l_2$  has equation  $\mathbf{r} = \begin{pmatrix} 0 \\ 9 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix}$ , where  $\mu$  is a scalar parameter.

Given that  $l_1$  and  $l_2$  meet at the point  $C$ , find

(a) the coordinates of  $C$ . (3)

The point  $A$  is the point on  $l_1$  where  $\lambda = 0$  and the point  $B$  is the point on  $l_2$  where  $\mu = -1$ .

(b) Find the size of the angle  $ACB$ . Give your answer in degrees to 2 decimal places. (4)

(c) Hence, or otherwise, find the area of the triangle  $ABC$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









Leave  
blank

**Question 7 continued**

Lined area for writing the answer to Question 7 continued.

Q7


--	--

(Total 12 marks)



Leave blank

8.



**Figure 2**

Figure 2 shows a cylindrical water tank. The diameter of a circular cross-section of the tank is 6 m. Water is flowing into the tank at a constant rate of  $0.48\pi \text{ m}^3 \text{ min}^{-1}$ . At time  $t$  minutes, the depth of the water in the tank is  $h$  metres. There is a tap at a point  $T$  at the bottom of the tank. When the tap is open, water leaves the tank at a rate of  $0.6\pi h \text{ m}^3 \text{ min}^{-1}$ .

(a) Show that  $t$  minutes after the tap has been opened

$$75 \frac{dh}{dt} = (4 - 5h) \tag{5}$$

When  $t = 0$ ,  $h = 0.2$

(b) Find the value of  $t$  when  $h = 0.5$  (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









**BLANK PAGE**

