

2.




Figure 1

Figure 1 shows a metal cube which is expanding uniformly as it is heated. At time t seconds, the length of each edge of the cube is x cm, and the volume of the cube is V cm³.

(a) Show that $\frac{dV}{dx} = 3x^2$ (1)

Given that the volume, V cm³, increases at a constant rate of 0.048 cm³s⁻¹,

(b) find $\frac{dx}{dt}$, when $x = 8$ (2)

(c) find the rate of increase of the total surface area of the cube, in cm²s⁻¹, when $x = 8$ (3)



Leave
blank

Question 2 continued

Ruled area for writing answers, consisting of multiple horizontal lines.

(Total 6 marks)

Q2



Leave blank

3.

$$f(x) = \frac{6}{\sqrt{9 - 4x}}, \quad |x| < \frac{9}{4}$$

(a) Find the binomial expansion of $f(x)$ in ascending powers of x , up to and including the term in x^3 . Give each coefficient in its simplest form.

(6)

Use your answer to part (a) to find the binomial expansion in ascending powers of x , up to and including the term in x^3 , of

$$(b) \quad g(x) = \frac{6}{\sqrt{9 + 4x}}, \quad |x| < \frac{9}{4}$$

(1)

$$(c) \quad h(x) = \frac{6}{\sqrt{9 - 8x}}, \quad |x| < \frac{9}{8}$$

(2)



Leave blank

Question 4 continued

Handwritten student response area consisting of 27 horizontal lines.

Q4

(Total 5 marks)



P 4 1 4 8 4 A 0 1 5 3 2

Leave
blank

Question 5 continued

Lined area for writing the answer to Question 5.

(Total 12 marks)

Q5



Leave blank

6.




Figure 2

Figure 2 shows a sketch of the curve C with parametric equations

$$x = (\sqrt{3})\sin 2t, \quad y = 4 \cos^2 t, \quad 0 \leq t \leq \pi$$

- (a) Show that $\frac{dy}{dx} = k(\sqrt{3})\tan 2t$, where k is a constant to be determined. (5)

- (b) Find an equation of the tangent to C at the point where $t = \frac{\pi}{3}$.
Give your answer in the form $y = ax + b$, where a and b are constants. (4)

- (c) Find a cartesian equation of C . (3)

Leave blank

Question 6 continued

Area with horizontal lines for writing the answer to Question 6.

Q6

--	--

(Total 12 marks)



P 4 1 4 8 4 A 0 2 3 3 2

Leave blank

7.




Figure 3

Figure 3 shows a sketch of part of the curve with equation $y = x^{\frac{1}{2}} \ln 2x$.

The finite region R , shown shaded in Figure 3, is bounded by the curve, the x -axis and the lines $x = 1$ and $x = 4$

- (a) Use the trapezium rule, with 3 strips of equal width, to find an estimate for the area of R , giving your answer to 2 decimal places. (4)
- (b) Find $\int x^{\frac{1}{2}} \ln 2x \, dx$. (4)
- (c) Hence find the exact area of R , giving your answer in the form $a \ln 2 + b$, where a and b are exact constants. (3)



Leave
blank

Question 7 continued

Lined area for writing answers to Question 7.



Leave blank

Question 7 continued

Lined area for writing the answer to Question 7.



Leave
blank

Question 8 continued

Ruled area for writing the answer to Question 8.



Leave
blank

Question 8 continued

Blank lined area for student response.

(Total 10 marks)

Q8

TOTAL FOR PAPER: 75 MARKS

END

