

Mark Scheme (Results)

June 2011

GCE Core Mathematics C4 (6666) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025 or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:
<http://www.edexcel.com/Aboutus/contact-us/>

June 2011

Publications Code UA027663

All the material in this publication is copyright

© Edexcel Ltd 2011

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
- ft – follow through
- the symbol \checkmark will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- \square The second mark is dependent on gaining the first mark

June 2011
Core Mathematics C4 6666
Mark Scheme

Question Number	Scheme	Marks
1.	$9x^2 = A(x-1)(2x+1) + B(2x+1) + C(x-1)^2$	B1
	$x \rightarrow 1 \quad 9 = 3B \Rightarrow B = 3$	M1
	$x \rightarrow -\frac{1}{2} \quad \frac{9}{4} = \left(-\frac{3}{2}\right)^2 C \Rightarrow C = 1$	Any two of A, B, C A1
	$x^2 \text{ terms} \quad 9 = 2A + C \Rightarrow A = 4$	All three correct A1
	<p><i>Alternatives for finding A.</i></p> $x \text{ terms} \quad 0 = -A + 2B - 2C \Rightarrow A = 4$ Constant terms $0 = -A + B + C \Rightarrow A = 4$	(4) [4]

Question Number	Scheme	Marks
2.	$f(x) = (\dots + \dots)^{-\frac{1}{2}}$ $= 9^{-\frac{1}{2}} (\dots + \dots)^{\dots}$ $(1+kx^2)^n = 1+nkx^2 + \dots$ $(1+kx^2)^{-\frac{1}{2}} = \dots + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2}(kx^2)^2$ $\left(1 + \frac{4}{9}x^2\right)^{-\frac{1}{2}} = 1 - \frac{2}{9}x^2 + \frac{2}{27}x^4$ $f(x) = \frac{1}{3} - \frac{2}{27}x^2 + \frac{2}{81}x^4$	<p>M1</p> <p>B1 $3^{-1}, \frac{1}{3}$ or $\frac{1}{9^{\frac{1}{2}}}$</p> <p>M1 n not a natural number, $k \neq 1$</p> <p>A1 ft ft their $k \neq 1$</p> <p>A1</p> <p>A1 (6) [6]</p>

Question Number	Scheme	Marks
3.	(a) $\frac{dV}{dh} = \frac{1}{2}\pi h - \pi h^2$	or equivalent M1 A1
	At $h = 0.1$, $\frac{dV}{dh} = \frac{1}{2}\pi(0.1) - \pi(0.1)^2 = 0.04\pi$	$\frac{\pi}{25}$ M1 A1 (4)
	(b) $\frac{dh}{dt} = \frac{dV}{dt} \div \frac{dV}{dh} = \frac{\pi}{800} \times \frac{1}{\frac{1}{2}\pi h - \pi h^2}$	or $\frac{\pi}{800} \div$ their (a) M1
	At $h = 0.1$, $\frac{dh}{dt} = \frac{\pi}{800} \times \frac{25}{\pi} = \frac{1}{32}$	awrt 0.031 A1 (2) [6]

Question Number	Scheme	Marks
<p>4.</p>	<p>(a) 0.0333, 1.3596 awrt 0.0333, 1.3596</p>	<p>B1 B1 (2)</p>
	<p>(b) $\text{Area}(R) \approx \frac{1}{2} \times \frac{\sqrt{2}}{4} [\dots]$ $\approx \dots [0 + 2(0.0333 + 0.3240 + 1.3596) + 3.9210]$ ≈ 1.30 Accept</p>	<p>B1 M1 A1 (3)</p>
	<p>(c) $u = x^2 + 2 \Rightarrow \frac{du}{dx} = 2x$ $\text{Area}(R) = \int_0^{\sqrt{2}} x^3 \ln(x^2 + 2) dx$ $\int x^3 \ln(x^2 + 2) dx = \int x^2 \ln(x^2 + 2) x dx = \int (u - 2)(\ln u) \frac{1}{2} du$</p>	<p>B1 B1 M1</p>
	<p>Hence $\text{Area}(R) = \frac{1}{2} \int_2^4 (u - 2) \ln u du$ * cso</p>	<p>A1 (4)</p>
	<p>(d) $\int (u - 2) \ln u du = \left(\frac{u^2}{2} - 2u \right) \ln u - \int \left(\frac{u^2}{2} - 2u \right) \frac{1}{u} du$ $= \left(\frac{u^2}{2} - 2u \right) \ln u - \int \left(\frac{u}{2} - 2 \right) du$ $= \left(\frac{u^2}{2} - 2u \right) \ln u - \left(\frac{u^2}{4} - 2u \right) (+C)$</p>	<p>M1 A1 M1 A1</p>
	<p>$\text{Area}(R) = \frac{1}{2} \left[\left(\frac{u^2}{2} - 2u \right) \ln u - \left(\frac{u^2}{4} - 2u \right) \right]_2^4$ $= \frac{1}{2} [(8 - 8) \ln 4 - 4 + 8 - ((2 - 4) \ln 2 - 1 + 4)]$ $= \frac{1}{2} (2 \ln 2 + 1)$ ln 2 + 1/2</p>	<p>M1 A1 (6) [15]</p>

Question Number	Scheme	Marks
5.	$\frac{1}{y} \frac{dy}{dx} = \dots$ $\dots = 2 \ln x + 2x \left(\frac{1}{x} \right)$	B1 M1 A1
	<p>At $x = 2$, leading to</p> $\ln y = 2(2) \ln 2$ $y = 16$	M1 A1
	<p>At $(2, 16)$</p> $\frac{1}{16} \frac{dy}{dx} = 2 \ln 2 + 2$	M1
	$\frac{dy}{dx} = 16(2 + 2 \ln 2)$	A1
	<p><i>Alternative</i></p>	(7) [7]
	$y = e^{2x \ln x}$	B1
	$\frac{d}{dx}(2x \ln x) = 2 \ln x + 2x \left(\frac{1}{x} \right)$	M1 A1
	$\frac{dy}{dx} = \left(2 \ln x + 2x \left(\frac{1}{x} \right) \right) e^{2x \ln x}$	M1 A1
	<p>At $x = 2$,</p> $\frac{dy}{dx} = (2 \ln 2 + 2) e^{4 \ln 2}$	M1
	$= 16(2 + 2 \ln 2)$	A1 (7)

Question Number	Scheme	Marks
6.	<p>(a) i: $6 - \lambda = -5 + 2\mu$ j: $-3 + 2\lambda = 15 - 3\mu$ leading to $\lambda = 3, \mu = 4$ $\mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + 3 \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 7 \end{pmatrix}$ or $\mathbf{r} = \begin{pmatrix} -5 \\ 15 \\ 3 \end{pmatrix} + 4 \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 7 \end{pmatrix}$ k: LHS = $-2 + 3(3) = 7$, RHS = $3 + 4(1) = 7$ (As LHS = RHS, lines intersect)</p> <p>Alternatively for B1, showing that $\lambda = 3$ and $\mu = 4$ both give $\begin{pmatrix} 3 \\ 3 \\ 7 \end{pmatrix}$</p> <p>(b) $\begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} = -2 - 6 + 3 = \sqrt{14}\sqrt{14}\cos\theta$ ($\theta \approx 110.92^\circ$) Acute angle is 69.1°</p> <p>(c) $\mathbf{r} = \begin{pmatrix} 6 \\ -3 \\ -2 \end{pmatrix} + 1 \begin{pmatrix} -1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix}$ ($\Rightarrow B$ lies on l_1)</p> <p>(d) Let d be shortest distance from B to l_2</p> <div style="display: flex; align-items: center;"> <div style="flex: 1;"> <p>$\overline{AB} = \begin{pmatrix} 5 \\ -1 \\ 1 \end{pmatrix} - \begin{pmatrix} 3 \\ 3 \\ 7 \end{pmatrix} = \begin{pmatrix} 2 \\ -4 \\ -6 \end{pmatrix}$</p> <p>$\overline{AB} = \sqrt{(2)^2 + (-4)^2 + (-6)^2} = \sqrt{56}$</p> <p>$\frac{d}{\sqrt{56}} = \sin\theta$ $d = \sqrt{56}\sin 69.1^\circ \approx 6.99$</p> </div> <div style="flex: 1; text-align: center;"> </div> </div>	<p>Any two equations</p> <p>M1 M1 A1 M1 A1</p> <p>B1 (6)</p> <p>M1 A1</p> <p>A1 (3)</p> <p>B1 (1)</p> <p>M1 A1 M1 A1 (4) [14]</p>

Question Number	Scheme	Marks
7.	(a) $\tan \theta = \sqrt{3}$ or $\sin \theta = \frac{\sqrt{3}}{2}$ $\theta = \frac{\pi}{3}$	M1 A1 (2)
	(b) $\frac{dx}{d\theta} = \sec^2 \theta, \frac{dy}{d\theta} = \cos \theta$ $\frac{dy}{dx} = \frac{\cos \theta}{\sec^2 \theta} (= \cos^3 \theta)$	awrt 1.05 M1 A1
	At P, $m = \cos^3 \left(\frac{\pi}{3} \right) = \frac{1}{8}$	Can be implied A1
	Using $mm' = -1, m' = -8$ For normal $y - \frac{1}{2}\sqrt{3} = -8(x - \sqrt{3})$	M1 M1
	At Q, $y = 0$ $-\frac{1}{2}\sqrt{3} = -8(x - \sqrt{3})$	
	leading to $x = \frac{17}{16}\sqrt{3} \quad (k = \frac{17}{16})$	1.0625 A1 (6)
	(c) $\int y^2 dx = \int y^2 \frac{dx}{d\theta} d\theta = \int \sin^2 \theta \sec^2 \theta d\theta$ $= \int \tan^2 \theta d\theta$ $= \int (\sec^2 \theta - 1) d\theta$ $= \tan \theta - \theta (+C)$	M1 A1 A1 M1 A1
	$V = \pi \int_0^{\frac{\pi}{3}} y^2 dx = [\tan \theta - \theta]_0^{\frac{\pi}{3}} = \pi \left[\left(\sqrt{3} - \frac{\pi}{3} \right) - (0 - 0) \right]$	M1
	$= \sqrt{3}\pi - \frac{1}{3}\pi^2 \quad (p = 1, q = -\frac{1}{3})$	A1 (7) [15]

Question Number	Scheme	Marks
8.	<p>(a) $\int (4y+3)^{-\frac{1}{2}} dx = \frac{(4y+3)^{\frac{1}{2}}}{(4)(\frac{1}{2})} (+C)$ $(= \frac{1}{2}(4y+3)^{\frac{1}{2}} + C)$</p> <p>(b) $\int \frac{1}{\sqrt{4y+3}} dy = \int \frac{1}{x^2} dx$ $\int (4y+3)^{-\frac{1}{2}} dy = \int x^{-2} dx$ $\frac{1}{2}(4y+3)^{\frac{1}{2}} = -\frac{1}{x} (+C)$</p> <p>Using $(-2, 1.5)$ $\frac{1}{2}(4 \times 1.5 + 3)^{\frac{1}{2}} = -\frac{1}{-2} + C$ leading to $C = 1$ $\frac{1}{2}(4y+3)^{\frac{1}{2}} = -\frac{1}{x} + 1$ $(4y+3)^{\frac{1}{2}} = 2 - \frac{2}{x}$ $y = \frac{1}{4} \left(2 - \frac{2}{x} \right)^2 - \frac{3}{4}$</p>	<p>M1 A1 (2)</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 (6)</p> <p>[8]</p>

or equivalent

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publication.orders@edexcel.com

Order Code UA027663 June 2011

For more information on Edexcel qualifications, please visit
www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual
.....



Llywodraeth Cynulliad Cymru
Welsh Assembly Government

