Mathematics C2

Examiner's use only

Team Leader's use only

Question

2

3

4

5

6

7

8

9

10

Leave

Past Paper

This resource was created and owned by Pearson Edexcel

Centre No.			Paper Reference			Surname	Initial(s)				
Candidate No.			6	6	6	4	/	0	1	Signature	

6664/01

Edexcel GCE

Core Mathematics C2 **Advanced Subsidiary**

Wednesday 10 January 2007 – Afternoon

Time: 1 hour 30 minutes

Materials required for examination

Items included with question papers

Mathematical Formulae (Green)

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature.

Check that you have the correct question paper.

You must write your answer for each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 10 questions in this question paper. The total mark for this paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2007 Edexcel Limited



Turn over

Total



W850/R6664/57570 4/4/3/3/3/3/21,400

■ Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

6664

Find (a) $f''(x)$,		$f(x) = x^3 + 3x^2 + 5.$
(a) $f''(x)$, (b) $\int_{1}^{2} f(x) dx$.		$1(\lambda) - \lambda + 3\lambda + 3.$
(b) $\int_{1}^{2} f(x) dx$.	Find	
(b) $\int_{1}^{2} f(x) dx$.	(a) $f''(x)$,	
(4) (b) $\int_{1}^{1} f(x) dx$.		(3)
	(b) $\int_1^1 f(x) dx$.	(4)

/inter 2007 ast Paper	www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics Ca
		Leave blank
Question 1 conti	inued	Ulank
		<u>Q1</u>

(Total 7 marks)

■ Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

6664

(4)	$(1-2x)^5$. Give each term in its simplest form.
show that	b) If x is small, so that x^2 and higher powers can be ignored
	$(1+x)(1-2x)^5 \approx 1-9x$.
(2)	

_			
\A/:	nter	\sim	$^{-7}$
vv	ntor	711	

aper	This resource was created and owned by Pearson Edexcel	
Question 2 continu	ıed	
C		

Mathematics C2

Dact Daner	This recour

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

6664 Leave blank

Find an equation for C .				
		(6)		
_				
_				

Mathematics C2

		_
D (_	
Past	Paper	

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

6664

Solve the equation	
$5^x = 17,$	
3 – 17,	
giving your answer to 3 significant figures.	
	(3)

(Total 3 marks)

(2)

■ Past Paper

This resource was created and owned by Pearson Edexcel

6664

		Leave blank
5.	$f(x) = x^3 + 4x^2 + x - 6.$	
	(a) Use the factor theorem to show that $(r + 2)$ is a factor of $f(r)$	

(b) Factorise f(x) completely. **(4)**

(c) Write down all the solutions to the equation

$$x^3 + 4x^2 + x - 6 = 0. ag{1}$$

Question 5 continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Continued Contin	st Paper	This resource was created and owned by Pearson Edexcel	666
Question 5 continued			Leave
Q5	Ouestion 5 co	ntinued	biank
	Question 5 col	nunueu	
			05
(Total 7 marks)			
		(Total 7 m	arks)

■ Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

6664

	_
	7
Leave	
blank	
Olullic	

giving each solution in terms of π. (6)	$2\cos^2 x + 1 = 5\sin x,$	
	giving each solution in terms of π .	
		(6)

Wint	or 2	007
vvint	er z	W/ /

	This resource was created and owned by Fearson Edexcer	
uestion 6 continued		
uestion o continuet		

This resource was created and owned by Pearson Edexcel

6664 Leave

blank

7.






Figure 1 shows a sketch of part of the curve C with equation

$$y = x(x-1)(x-5).$$

Use calculus to find the total area of the finite region, shown shaded in Figure 1, that is between x = 0 and x = 2 and is bounded by C, the x-axis and the line x = 2.

(9)

Winter	2007
vvinter	/UU/

Winter 2007	www.mystudybro.com	Mathematics	C2
Past Paper	This resource was created and owned by Pearson Edexcel	(6664
		Le	ave

estion 7 continued		

	y = y	
Past Paper	This resource was created and owner	ed by Pearson Edexcel

•		J	•	_
		0	366	: 1

stion 7 continued		

₩.	ntar	201	17

Paper	This resource was created and owned by Pearson Edexcel	66
		Leav blank
Question 7 continued	I	
		Q7
		al 9 marks)

Past Paper

This resource was created and owned by Pearson Edexcel

Leave blank

A diesel lorry is driven from Birmingham to Bury at a steady speed of v kilometres per hour. The total cost of the journey, £C, is given by

 $C = \frac{1400}{v} + \frac{2v}{7}$.

(a) Find the value of v for which C is a minimum.

(5)

(b) Find $\frac{d^2C}{dv^2}$ and hence verify that C is a minimum for this value of v.

(2)

(c) Calculate the minimum total cost of the journey.

(2)

₩.	ntar	201	17

per	This resource was created and owned by Pearson Edexcel	
)	a a	
Question 8 continue	a	

6664

Leave blank

9.




Figure 2 shows a plan of a patio. The patio PQRS is in the shape of a sector of a circle with centre Q and radius $6 \, \text{m}$.

Given that the length of the straight line PR is $6\sqrt{3}$ m,

(a) find the exact size of angle PQR in radians.

(3)

(b) Show that the area of the patio PQRS is 12π m².

(2)

(c) Find the exact area of the triangle *PQR*.

(2)

(d) Find, in m^2 to 1 decimal place, the area of the segment *PRS*.

(2)

(e) Find, in m to 1 decimal place, the perimeter of the patio PQRS.

(2)

Winter	2007
vviiitei	ZUUI

Willer ZUU1	www.iiiyStudybiO.com	Maniemancs CZ
Past Paper	This resource was created and owned by Pearson Edexcel	6664

iestion 9 continu	-	

www.mvstudvbro.com

M

	,,
Past Paper	This resource was created and owned by Pearson Edexcel

lathematics	C2
	6664

	Leave blank
Question 9 continued	

aper	This resource was created and owned by Pearson Edexcel	66
		Lea
0	1	blar
Question 9 continue	ed .	
		Q
	(Total 11 ma	arks)

(3)

(3)

■ Past Paper

This resource was created and owned by Pearson Edexcel

Leave blank

- **10.** A geometric series is $a + ar + ar^2 + ...$
 - (a) Prove that the sum of the first n terms of this series is given by

$$S_n = \frac{a(1-r^n)}{1-r}.$$

(4)

(b) Find

$$\sum_{k=1}^{10} 100(2^k).$$

(c) Find the sum to infinity of the geometric series

$$\frac{5}{6} + \frac{5}{18} + \frac{5}{54} + \dots$$

(d) State the condition for an infinite geometric series with common ratio r to be convergent.

(1)



\A/:	nter	20	^7
vvi	nter	70	.,,

William 2001	www.mystudybro.com	Mathematics CZ
Past Paper	This resource was created and owned by Pearson Edexcel	6664

estion 10 continued	

Mathematics C2

www.mystudybro.comThis resource was created and owned by Pearson Edexcel ■ Past Paper

6664