Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	7	/	0	1	Signature	

Paper Reference(s)

6667/01

Examiner's use only

Team Leader's use only

Edexcel GCE

Further Pure Mathematics FP1

Advanced/Advanced Subsidiary

Tuesday 22 June 2010 – Afternoon

Time: 1 hour 30 minutes

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
9	
Total	

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer to each question in the space following the question.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 9 questions in this question paper. The total mark for this paper is 75.
There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

Printer's Log. No.
N35387A
W850/R6667/57570 4/3

N 3 5 3 8 7 A 0 1 2 8

Turn over

edexcel
advancing learning, changing lives

Leave blank

1. $z = 2 - 3i$

(a) Show that $z^2 = -5 - 12i$.

(2)

Find, showing your working,

(b) the value of $|z^2|$,

(2)

(c) the value of $\arg(z^2)$, giving your answer in radians to 2 decimal places.

(2)

(d) Show z and z^2 on a single Argand diagram.

(1)

Leave blank

Question 1 continued

Q1

(Total 7 marks)

Leave blank

2. $\mathbf{M} = \begin{pmatrix} 2a & 3 \\ 6 & a \end{pmatrix}$, where a is a real constant.

(a) Given that $a = 2$, find $\mathbf{M}^{-1}$.

(3)

(b) Find the values of a for which $\mathbf{M}$ is singular.

(2)

Leave blank

Question 2 continued

Q2

(Total 5 marks)

Leave blank

3. $$f(x) = x^3 - \frac{7}{x} + 2, \quad x > 0$$

(a) Show that $f(x) = 0$ has a root α between 1.4 and 1.5

(2)

(b) Starting with the interval $[1.4, 1.5]$, use interval bisection twice to find an interval of width 0.025 that contains α.

(3)

(c) Taking 1.45 as a first approximation to α, apply the Newton-Raphson procedure once to $f(x) = x^3 - \frac{7}{x} + 2$ to obtain a second approximation to α, giving your answer to 3 decimal places.

(5)

Leave blank

Question 3 continued

Leave blank

Question 3 continued

Leave blank

Question 3 continued

Q3

(Total 10 marks)

Leave blank

4. $f(x) = x^3 + x^2 + 44x + 150$

Given that $f(x) = (x+3)(x^2 + ax + b)$, where a and b are real constants,

(a) find the value of a and the value of b.

(2)

(b) Find the three roots of $f(x) = 0$.

(4)

(c) Find the sum of the three roots of $f(x) = 0$.

(1)

Leave blank

Question 4 continued

Q4

(Total 7 marks)

Leave blank

5. The parabola C has equation $y^2 = 20x$.

(a) Verify that the point $P(5t^2, 10t)$ is a general point on C.

(1)

The point A on C has parameter $t = 4$.
The line l passes through A and also passes through the focus of C.

(b) Find the gradient of l.

(4)

Leave blank

Question 5 continued

Q5

(Total 5 marks)

Leave blank

6. Write down the 2×2 matrix that represents

(a) an enlargement with centre $(0, 0)$ and scale factor 8,

(1)

(b) a reflection in the x-axis.

(1)

Hence, or otherwise,

(c) find the matrix **T** that represents an enlargement with centre $(0, 0)$ and scale factor 8, followed by a reflection in the x-axis.

(2)

$$\mathbf{A} = \begin{pmatrix} 6 & 1 \\ 4 & 2 \end{pmatrix} \text{ and } \mathbf{B} = \begin{pmatrix} k & 1 \\ c & -6 \end{pmatrix}, \text{ where } k \text{ and } c \text{ are constants.}$$

(d) Find **AB**.

(3)

Given that **AB** represents the same transformation as **T**,

(e) find the value of k and the value of c.

(2)

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Leave blank

Question 6 continued

Q6

(Total 9 marks)

Leave blank

7. $f(n) = 2^n + 6^n$

(a) Show that $f(k+1) = 6f(k) - 4(2^k)$.

(3)

(b) Hence, or otherwise, prove by induction that, for $n \in \mathbb{Z}^+$, $f(n)$ is divisible by 8.

(4)

Leave blank

Question 7 continued

Q7

(Total 7 marks)

Leave blank

8. The rectangular hyperbola H has equation $xy = c^2$, where c is a positive constant.

The point A on H has x-coordinate $3c$.

(a) Write down the y-coordinate of A.

(1)

(b) Show that an equation of the normal to H at A is

$$3y = 27x - 80c$$

(5)

The normal to H at A meets H again at the point B.

(c) Find, in terms of c, the coordinates of B.

(5)

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Q8

(Total 11 marks)

Leave blank

9. (a) Prove by induction that

$$\sum_{r=1}^{n} r^2 = \frac{1}{6}n(n+1)(2n+1)$$

(6)

Using the standard results for $\sum_{r=1}^{n} r$ and $\sum_{r=1}^{n} r^2$,

(b) show that

$$\sum_{r=1}^{n} (r+2)(r+3) = \frac{1}{3}n(n^2 + an + b),$$

where a and b are integers to be found.

(5)

(c) Hence show that

$$\sum_{r=n+1}^{2n} (r+2)(r+3) = \frac{1}{3}n(7n^2 + 27n + 26)$$

(3)

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Q9

(Total 14 marks)

TOTAL FOR PAPER: 75 MARKS

END