Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	7	/	0	1	Signature	

Paper Reference(s)

6667/01

Edexcel GCE

Further Pure Mathematics FP1

Advanced/Advanced Subsidiary

Wednesday 22 June 2011 – Morning

Time: 1 hour 30 minutes

Examiner's use only

Team Leader's use only

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
9	
Total	

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer to each question in the space following the question.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 9 questions in this question paper. The total mark for this paper is 75.
There are 32 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2011 Edexcel Limited.

Printer's Log. No.
P38168A
W850/R6667/57570 5/5/5/3

P 3 8 1 6 8 A 0 1 3 2

Turn over

edexcel
advancing learning, changing lives

Leave blank

1. $$f(x) = 3^x + 3x - 7$$

(a) Show that the equation $f(x) = 0$ has a root α between $x = 1$ and $x = 2$.

(2)

(b) Starting with the interval $[1, 2]$, use interval bisection twice to find an interval of width 0.25 which contains α.

(3)

Leave blank

Question 1 continued

Q1

(Total 5 marks)

Leave blank

2.

$$z_1 = -2 + \mathrm{i}$$

(a) Find the modulus of z_1.

(1)

(b) Find, in radians, the argument of z_1, giving your answer to 2 decimal places.

(2)

The solutions to the quadratic equation

$$z^2 - 10z + 28 = 0$$

are z_2 and z_3.

(c) Find z_2 and z_3, giving your answers in the form $p \pm \mathrm{i}\sqrt{q}$, where p and q are integers.

(3)

(d) Show, on an Argand diagram, the points representing your complex numbers z_1, z_2 and z_3.

(2)

P 3 8 1 6 8 A 0 4 3 2

Leave blank

Question 2 continued

Leave blank

Question 2 continued

Leave blank

Question 2 continued

Q2

(Total 8 marks)

Leave blank

3. (a) Given that

$$\mathbf{A} = \begin{pmatrix} 1 & \sqrt{2} \\ \sqrt{2} & -1 \end{pmatrix}$$

(i) find $\mathbf{A}^2$,

(ii) describe fully the geometrical transformation represented by $\mathbf{A}^2$.

(4)

(b) Given that

$$\mathbf{B} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

describe fully the geometrical transformation represented by **B**.

(2)

(c) Given that

$$\mathbf{C} = \begin{pmatrix} k+1 & 12 \\ k & 9 \end{pmatrix}$$

where k is a constant, find the value of k for which the matrix **C** is singular.

(3)

P 3 8 1 6 8 A 0 8 3 2

Leave blank

Question 3 continued

Leave blank

Question 3 continued

P 3 8 1 6 8 A 0 1 0 3 2

Leave blank

Question 3 continued

Q3

(Total 9 marks)

Leave blank

4. $$\mathrm{f}(x) = x^2 + \frac{5}{2x} - 3x - 1\,, \quad x \neq 0$$

(a) Use differentiation to find $\mathrm{f}'(x)$.

(2)

The root α of the equation $\mathrm{f}(x) = 0$ lies in the interval $[0.7,\ 0.9]$.

(b) Taking 0.8 as a first approximation to α, apply the Newton-Raphson process once to $\mathrm{f}(x)$ to obtain a second approximation to α. Give your answer to 3 decimal places.

(4)

P 3 8 1 6 8 A 0 1 2 3 2

Leave blank

Question 4 continued

Q4

(Total 6 marks)

Leave blank

5. $$\mathbf{A} = \begin{pmatrix} -4 & a \\ b & -2 \end{pmatrix}, \text{ where } a \text{ and } b \text{ are constants.}$$

Given that the matrix **A** maps the point with coordinates $(4, 6)$ onto the point with coordinates $(2, -8)$,

(a) find the value of a and the value of b.

(4)

A quadrilateral R has area 30 square units.
It is transformed into another quadrilateral S by the matrix **A**.
Using your values of a and b,

(b) find the area of quadrilateral S.

(4)

P 3 8 1 6 8 A 0 1 4 3 2

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Leave blank

Question 5 continued

Q5

(Total 8 marks)

Leave blank

6. Given that $z = x + \mathrm{i}y$, find the value of x and the value of y such that

$$z + 3\mathrm{i}z^* = -1 + 13\mathrm{i}$$

where z^* is the complex conjugate of z.

(7)

Leave blank

Question 6 continued

Q6

(Total 7 marks)

Leave blank

7. (a) Use the results for $\sum_{r=1}^{n} r$ and $\sum_{r=1}^{n} r^2$ to show that

$$\sum_{r=1}^{n}(2r-1)^2 = \frac{1}{3}n(2n+1)(2n-1)$$

for all positive integers n.

(6)

(b) Hence show that

$$\sum_{r=n+1}^{3n}(2r-1)^2 = \frac{2}{3}n\left(an^2+b\right)$$

where a and b are integers to be found.

(4)

P 3 8 1 6 8 A 0 2 0 3 2

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Leave blank

Question 7 continued

Q7

(Total 10 marks)

Leave blank

8. The parabola C has equation $y^2 = 48x$.

The point $P(12t^2, 24t)$ is a general point on C.

(a) Find the equation of the directrix of C.

(2)

(b) Show that the equation of the tangent to C at $P(12t^2, 24t)$ is

$$x - ty + 12t^2 = 0$$

(4)

The tangent to C at the point $(3, 12)$ meets the directrix of C at the point X.

(c) Find the coordinates of X.

(4)

P 3 8 1 6 8 A 0 2 4 3 2

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Leave blank

Question 8 continued

Q8

(Total 10 marks)

Leave blank

9. Prove by induction, that for $n \in \mathbb{Z}^+$,

(a) $\begin{pmatrix} 3 & 0 \\ 6 & 1 \end{pmatrix}^n = \begin{pmatrix} 3^n & 0 \\ 3(3^n - 1) & 1 \end{pmatrix}$,

(6)

(b) $\mathrm{f}(n) = 7^{2n-1} + 5$ is divisible by 12.

(6)

P 3 8 1 6 8 A 0 2 8 3 2

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Leave blank

Question 9 continued

Q9

(Total 12 marks)

TOTAL FOR PAPER: 75 MARKS

END

P 3 8 1 6 8 A 0 3 2 3 2