

Mark Scheme (Results) Summer 2010

GCE

Further Pure Mathematics FP1 (6667)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.


For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

June 2010
Further Pure Mathematics FP1 6667
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) $(2 - 3i)(2 - 3i) = \dots$ Expand and use $i^2 = -1$, getting completely correct expansion of 3 or 4 terms</p> <p>Reaches $-5 - 12i$ after completely correct work (must see $4 - 9$) (*)</p>	<p>M1 A1cso (2)</p>
	<p>(b) $z^2 = \sqrt{(-5)^2 + (-12)^2} = 13$ or $z^2 = \sqrt{5^2 + 12^2} = 13$</p> <p>Alternative methods for part (b)</p> <p>$z^2 = z ^2 = 2^2 + (-3)^2 = 13$ Or: $z^2 = zz^* = 13$</p>	<p>M1 A1 (2) M1 A1 (2)</p>
	<p>(c) $\tan \alpha = \frac{12}{5}$ (allow $-\frac{12}{5}$) or $\sin \alpha = \frac{12}{13}$ or $\cos \alpha = \frac{5}{13}$</p> <p>$\arg(z^2) = -(\pi - 1.176\dots) = -1.97$ (or 4.32) allow awrt</p> <p>Alternative method for part (c) $\alpha = 2 \times \arctan\left(-\frac{3}{2}\right)$ (allow $\frac{3}{2}$) or use $\frac{\pi}{2} + \arctan \frac{5}{12}$</p> <p>so $\arg(z^2) = -(\pi - 1.176\dots) = -1.97$ (or 4.32) allow awrt</p>	<p>M1 A1 (2) M1 A1</p>
	<p>(d)</p> <div style="display: flex; align-items: center; justify-content: center;">  <div style="margin-left: 20px;"> <p>Both in correct quadrants. Approximate relative scale No labels needed Allow two diagrams if some indication of scale Allow points or arrows</p> </div> </div>	<p>B1 (1) 7 marks</p>
	<p>Notes: (a) M1: for $4 - 9 - 12i$ or $4 - 9 - 6i - 6i$ or $4 - 3^2 - 12i$ but must have correct statement seen and see i^2 replaced by -1 maybe later A1: Printed answer. Must see $4 - 9$ in working. Jump from $4 - 6i - 6i + 9i^2$ to $-5 - 12i$ is M0A0</p> <p>(b) Method may be implied by correct answer. NB $z^2 = 169$ is M0 A0</p> <p>(c) Allow $\arctan \frac{12}{5}$ for M1 or $\pm \frac{\pi}{2} \pm \arctan \frac{5}{12}$</p>	

Question Number	Scheme	Marks
2.	<p>(a) $\mathbf{M} = \begin{pmatrix} 4 & 3 \\ 6 & 2 \end{pmatrix}$ Determinant: $(8 - 18) = -10$</p> $\mathbf{M}^{-1} = \frac{1}{-10} \begin{pmatrix} 2 & -3 \\ -6 & 4 \end{pmatrix} \quad \left[= \begin{pmatrix} -0.2 & 0.3 \\ 0.6 & -0.4 \end{pmatrix} \right]$	<p>B1</p> <p>M1 A1</p> <p>(3)</p>
	<p>(b) Setting $\Delta = 0$ and using $2a^2 \pm 18 = 0$ to obtain $a = .$</p> $a = \pm 3$	<p>M1</p> <p>A1 cao</p> <p>(2)</p> <p>5 marks</p>
	<p>Notes:</p> <p>(a) B1: must be -10</p> <p>M1: for correct attempt at changing elements in major diagonal and changing signs in minor diagonal. Three or four of the numbers in the matrix should be correct – eg allow one slip</p> <p>A1: for any form of the correct answer, with correct determinant then isw.</p> <p>Special case: a not replaced is B0M1A0</p> <p>(b) Two correct answers, $a = \pm 3$, with no working is M1A1</p> <p>Just $a = 3$ is M1A0, and also one of these answers rejected is A0.</p> <p>Need 3 to be simplified (not $\sqrt{9}$).</p>	

Question Number	Scheme	Marks
3.	<p>(a) $f(1.4) = \dots$ and $f(1.5) = \dots$ Evaluate both</p> <p>$f(1.4) = -0.256$ (or $-\frac{32}{125}$), $f(1.5) = 0.708\dots$ (or $\frac{17}{24}$) Change of sign, \therefore root</p> <p>Alternative method: Graphical method could earn M1 if 1.4 and 1.5 are both indicated A1 then needs correct graph and conclusion, i.e. change of sign \therefore root</p>	<p>M1 A1 (2)</p>
	<p>(b) $f(1.45) = 0.221\dots$ or 0.2 [\therefore root is in [1.4, 1.45]]</p> <p>$f(1.425) = -0.018\dots$ or -0.019 or -0.02</p> <p>\therefore root is in [1.425, 1.45]</p>	<p>M1 M1 A1cso (3)</p>
	<p>(c) $f'(x) = 3x^2 + 7x^{-2}$</p> <p>$f'(1.45) = 9.636\dots$ (Special case: $f'(x) = 3x^2 + 7x^{-2} + 2$ then $f'(1.45) = 11.636\dots$)</p> <p>$x_1 = 1.45 - \frac{f(1.45)}{f'(1.45)} = 1.45 - \frac{0.221\dots}{9.636\dots} = 1.427$</p>	<p>M1 A1 A1ft M1 A1cao (5) 10 marks</p>
<p>Notes</p> <p>(a) M1: Some attempt at two evaluations A1: needs accuracy to 1 figure truncated or rounded and conclusion including sign change indicated (One figure accuracy sufficient)</p> <p>(b) M1: See $f(1.45)$ attempted and positive M1: See $f(1.425)$ attempted and negative A1: is cso – any slips in numerical work are penalised here even if correct region found. Answer may be written as $1.425 \leq \alpha \leq 1.45$ or $1.425 < \alpha < 1.45$ or (1.425, 1.45) must be correct way round. Between is sufficient. There is no credit for linear interpolation. This is M0 M0 A0 Answer with no working is also M0M0A0</p> <p>(c) M1: for attempt at differentiation (decrease in power) A1 is cao Second A1 may be implied by correct answer (do not need to see it) ft is limited to special case given. 2nd M1: for attempt at Newton Raphson with their values for $f(1.45)$ and $f'(1.45)$. A1: is cao and needs to be correct to 3dp Newton Raphson used more than once – isw. Special case: $f'(x) = 3x^2 + 7x^{-2} + 2$ then $f'(1.45) = 11.636\dots$) is M1 A0 A1ft M1 A0 This mark can also be given by implication from final answer of 1.43</p>		

Question Number	Scheme	Marks
4.	(a) $a = -2, b = 50$	B1, B1 (2)
	(b) -3 is a root Solving 3-term quadratic $x = \frac{2 \pm \sqrt{4 - 200}}{2}$ or $(x-1)^2 - 1 + 50 = 0$ $= 1 + 7i, 1 - 7i$	B1 M1 A1, A1ft (4)
	(c) $(-3) + (1 + 7i) + (1 - 7i) = -1$	B1ft (1) 7 marks
	Notes (a) Accept $x^2 - 2x + 50$ as evidence of values of a and b . (b) B1: -3 must be seen in part (b) M1: for solving quadratic following usual conventions A1: for a correct root (simplified as here) and A1ft: for conjugate of first answer. Accept correct answers with no working here. If answers are written down as factors then isw. Must see roots for marks. (c) ft requires the sum of two non-real conjugate roots and a real root resulting in a real number. Answers including x are B0	

Question Number	Scheme	Marks
5.	<p>(a) $y^2 = (10t)^2 = 100t^2$ and $20x = 20 \times 5t^2 = 100t^2$</p> <p>Alternative method: Compare with $y^2 = 4ax$ and identify $a = 5$ to give answer.</p>	<p>B1 (1)</p> <p>B1 (1)</p>
	<p>(b) Point A is $(80, 40)$ (stated or seen on diagram). May be given in part (a) Focus is $(5, 0)$ (stated or seen on diagram) or $(a, 0)$ with $a = 5$ May be given in part (a). Gradient: $\frac{40-0}{80-5} = \frac{40}{75} \left(= \frac{8}{15} \right)$</p>	<p>B1</p> <p>B1</p> <p>M1 A1 (4)</p> <p>5 marks</p>
	<p>Notes:</p> <p>(a) Allow substitution of x to obtain $y = \pm 10t$ (or just $10t$) or of y to obtain x</p> <p>(b) M1: requires use of gradient formula correctly, for their values of x and y. This mark may be implied by correct answer. Differentiation is M0 A0 A1: Accept 0.533 or awrt</p>	

Question Number	Scheme	Marks
6.	(a) $\begin{pmatrix} 8 & 0 \\ 0 & 8 \end{pmatrix}$	B1 (1)
	(b) $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$	B1 (1)
	(c) $\mathbf{T} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 8 & 0 \\ 0 & 8 \end{pmatrix} = \begin{pmatrix} 8 & 0 \\ 0 & -8 \end{pmatrix}$	M1 A1 (2)
	(d) $\mathbf{AB} = \begin{pmatrix} 6 & 1 \\ 4 & 2 \end{pmatrix} \begin{pmatrix} k & 1 \\ c & -6 \end{pmatrix} = \begin{pmatrix} 6k+c & 0 \\ 4k+2c & -8 \end{pmatrix}$	M1 A1 A1 (3)
	(e) “ $6k + c = 8$ ” and “ $4k + 2c = 0$ ” Form equations and solve simultaneously $k = 2$ and $c = -4$	M1 A1 (2) 9 marks
	<p>Alternative method for (e) M1: $\mathbf{AB} = \mathbf{T} \Rightarrow \mathbf{B} = \mathbf{A}^{-1}\mathbf{T}$ = and compare elements to find k and c. Then A1 as before.</p>	
	<p><u>Notes</u></p> <p>(c) M1: Accept multiplication of their matrices either way round (this can be implied by correct answer) A1: cao</p> <p>(d) M1: Correct matrix multiplication method implied by one or two correct terms in correct positions. A1: for three correct terms in correct positions 2nd A1: for all four terms correct and simplified</p> <p>(e) M1: follows their previous work but must give two equations from which k and c can be found and there must be attempt at solution getting to $k =$ or $c =$. A1: is cao (but not cso - may follow error in position of $4k + 2c$ earlier).</p>	

Question Number	Scheme		Marks
7.	(a) LHS = $f(k+1) = 2^{k+1} + 6^{k+1}$ $= 2(2^k) + 6(6^k)$ $= 6(2^k + 6^k) - 4(2^k) = 6f(k) - 4(2^k)$	OR RHS = $= 6f(k) - 4(2^k) = 6(2^k + 6^k) - 4(2^k)$ $= 2(2^k) + 6(6^k)$ $= 2^{k+1} + 6^{k+1} = f(k+1)$ (*)	M1 A1 A1 (3)
	OR $f(k+1) - 6f(k) = 2^{k+1} + 6^{k+1} - 6(2^k + 6^k)$		M1
	$= (2-6)(2^k) = -4 \cdot 2^k$, and so $f(k+1) = 6f(k) - 4(2^k)$		A1, A1 (3)
	(b) $n = 1$: $f(1) = 2^1 + 6^1 = 8$, which is divisible by 8		B1
	Either Assume $f(k)$ divisible by 8 and try to use $f(k+1) = 6f(k) - 4(2^k)$ Show $4(2^k) = 4 \times 2(2^{k-1}) = 8(2^{k-1})$ or $8(\frac{1}{2}2^k)$ Or valid statement Deduction that result is implied for $n = k + 1$ and so is true for positive integers by induction (may include $n = 1$ true here)	Or Assume $f(k)$ divisible by 8 and try to use $f(k+1) - f(k)$ or $f(k+1) + f(k)$ including factorising $6^k = 2^k 3^k$ $= 2^3 2^{k-3} (1 + 5 \cdot 3^k)$ or $= 2^3 2^{k-3} (3 + 7 \cdot 3^k)$ o.e. Deduction that result is implied for $n = k + 1$ and so is true for positive integers by induction (must include explanation of why $n = 2$ is also true here)	M1 A1 A1cso (4) 7 marks
Notes (a) M1: for substitution into LHS (or RHS) or $f(k+1) - 6f(k)$ A1: for correct split of the two separate powers cao A1: for completion of proof with no error or ambiguity (needs (for example) to start with one side of equation and reach the other or show that each side separately is $2(2^k) + 6(6^k)$ and conclude LHS = RHS) (b) B1: for substitution of $n = 1$ and stating “true for $n = 1$ ” or “divisible by 8” or tick. (This statement may appear in the concluding statement of the proof) M1: Assume $f(k)$ divisible by 8 and consider $f(k+1) = 6f(k) - 4(2^k)$ or equivalent expression that could lead to proof – not merely $f(k+1) - f(k)$ unless deduce that 2 is a factor of 6 (see right hand scheme above). A1: Indicates each term divisible by 8 OR takes out factor 8 or 2^3 A1: Induction statement . Statement $n = 1$ here could contribute to B1 mark earlier. NB: $f(k+1) - f(k) = 2^{k+1} - 2^k + 6^{k+1} - 6^k = 2^k + 5 \cdot 6^k$ only is M0 A0 A0 (b) “ Otherwise ” methods Could use: $f(k+1) = 2f(k) + 4(6^k)$ or $f(k+2) = 36f(k) - 32(6^k)$ or $f(k+2) = 4f(k) + 32(2^k)$ in a similar way to given expression and Left hand mark scheme is applied. Special Case: Otherwise Proof not involving induction : This can only be awarded the B1 for checking $n = 1$.			

Question Number	Scheme	Marks						
8.	(a) $\frac{c}{3}$	B1 (1)						
	(b) $y = \frac{c^2}{x} \Rightarrow \frac{dy}{dx} = -c^2x^{-2}$, or $y + x \frac{dy}{dx} = 0 \Rightarrow \frac{dy}{dx} = -\frac{y}{x}$ or $\dot{x} = c, \dot{y} = -\frac{c}{t^2}$ so $\frac{dy}{dx} = -\frac{1}{t^2}$ and at A $\frac{dy}{dx} = -\frac{c^2}{(3c)^2} = -\frac{1}{9}$ so gradient of normal is 9 Either $y - \frac{c}{3} = 9(x - 3c)$ or $y = 9x + k$ and use $x = 3c, y = \frac{c}{3}$ $\Rightarrow 3y = 27x - 80c$ (*)	B1 M1 A1 M1 A1 (5)						
	<table border="0" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; border-right: 1px solid black; padding: 5px;"> (c) $\frac{c^2}{x} = \frac{27x - 80c}{3}$ $3c^2 = 27x^2 - 80cx$ </td> <td style="width: 33%; border-right: 1px solid black; padding: 5px;"> $\frac{c^2}{y} = \frac{3y + 80c}{27}$ $27c^2 = 3y^2 + 80cy$ </td> <td style="width: 33%; padding: 5px;"> $3\frac{c}{t} = 27ct - 80c$ $3c = 27ct^2 - 80ct$ </td> </tr> <tr> <td style="border-right: 1px solid black; padding: 5px;"> $(x - 3c)(27x + c) = 0$ so $x =$ $x = -\frac{c}{27}, y = -27c$ </td> <td style="border-right: 1px solid black; padding: 5px;"> $(y + 27c)(3y - c) = 0$ so $y =$ $x = -\frac{c}{27}, y = -27c$ </td> <td style="padding: 5px;"> $(t - 3)(27t + 1) = 0$ so $t =$ $(t = -\frac{1}{27}$ and so) $x = -\frac{c}{27}, y = -27c$ </td> </tr> </table>	(c) $\frac{c^2}{x} = \frac{27x - 80c}{3}$ $3c^2 = 27x^2 - 80cx$	$\frac{c^2}{y} = \frac{3y + 80c}{27}$ $27c^2 = 3y^2 + 80cy$	$3\frac{c}{t} = 27ct - 80c$ $3c = 27ct^2 - 80ct$	$(x - 3c)(27x + c) = 0$ so $x =$ $x = -\frac{c}{27}, y = -27c$	$(y + 27c)(3y - c) = 0$ so $y =$ $x = -\frac{c}{27}, y = -27c$	$(t - 3)(27t + 1) = 0$ so $t =$ $(t = -\frac{1}{27}$ and so) $x = -\frac{c}{27}, y = -27c$	M1 A1 M1 A1, A1 (5) 11 marks
(c) $\frac{c^2}{x} = \frac{27x - 80c}{3}$ $3c^2 = 27x^2 - 80cx$	$\frac{c^2}{y} = \frac{3y + 80c}{27}$ $27c^2 = 3y^2 + 80cy$	$3\frac{c}{t} = 27ct - 80c$ $3c = 27ct^2 - 80ct$						
$(x - 3c)(27x + c) = 0$ so $x =$ $x = -\frac{c}{27}, y = -27c$	$(y + 27c)(3y - c) = 0$ so $y =$ $x = -\frac{c}{27}, y = -27c$	$(t - 3)(27t + 1) = 0$ so $t =$ $(t = -\frac{1}{27}$ and so) $x = -\frac{c}{27}, y = -27c$						
	Notes (b) B1: Any valid method of differentiation but must get to correct expression for $\frac{dy}{dx}$ M1 : Substitutes values and uses negative reciprocal (needs to follow calculus) A1: 9 cao (needs to follow calculus) M1: Finds equation of line through A with any gradient (other than 0 and ∞) A1: Correct work throughout – obtaining printed answer . (c) M1: Obtains equation in one variable (x, y or t) A1: Writes as correct three term quadratic (any equivalent form) M1: Attempts to solve three term quadratic to obtain $x =$ or $y =$ or $t =$ A1: x coordinate, A1: y coordinate. (cao but allow recovery following slips)							

Question Number	Scheme	Marks
9.	<p>(a) If $n = 1$, $\sum_{r=1}^n r^2 = 1$ and $\frac{1}{6}n(n+1)(2n+1) = \frac{1}{6} \times 1 \times 2 \times 3 = 1$, so true for $n = 1$.</p> <p>Assume result true for $n = k$</p> $\sum_{r=1}^{k+1} r^2 = \frac{1}{6}k(k+1)(2k+1) + (k+1)^2$ $= \frac{1}{6}(k+1)(2k^2 + 7k + 6) \text{ or } = \frac{1}{6}(k+2)(2k^2 + 5k + 3) \text{ or } = \frac{1}{6}(2k+3)(k^2 + 3k + 2)$ $= \frac{1}{6}(k+1)(k+2)(2k+3) = \frac{1}{6}(k+1)(\{k+1\}+1)(2\{k+1\}+1) \text{ or equivalent}$ <p>True for $n = k + 1$ if true for $n = k$, (and true for $n = 1$) so true by induction for all n.</p>	<p>B1 M1 M1 A1 dM1 A1cso (6)</p>
	<p>Alternative for (a) After first three marks B M M1 as earlier :</p> <p>May state RHS = $\frac{1}{6}(k+1)(\{k+1\}+1)(2\{k+1\}+1) = \frac{1}{6}(k+1)(k+2)(2k+3)$ for third M1</p> <p>Expands to $\frac{1}{6}(k+1)(2k^2 + 7k + 6)$ and show equal to $\sum_{r=1}^{k+1} r^2 = \frac{1}{6}k(k+1)(2k+1) + (k+1)^2$ for A1</p> <p>So true for $n = k + 1$ if true for $n = k$, and true for $n = 1$, so true by induction for all n.</p>	<p>B1M1M1 dM1 A1 A1cso (6)</p>
	<p>(b) $\sum_{r=1}^n (r^2 + 5r + 6) = \sum_{r=1}^n r^2 + 5\sum_{r=1}^n r + (\sum_{r=1}^n 6)$</p> $\frac{1}{6}n(n+1)(2n+1) + \frac{5}{2}n(n+1), + 6n$ $= \frac{1}{6}n[(n+1)(2n+1) + 15(n+1) + 36]$ $= \frac{1}{6}n[2n^2 + 18n + 52] = \frac{1}{3}n(n^2 + 9n + 26) \quad \text{or } a = 9, b = 26$	<p>M1 A1, B1 M1 A1 (5)</p>
	<p>(c) $\sum_{r=n+1}^{2n} (r+2)(r+3) = \frac{1}{3}2n(4n^2 + 18n + 26) - \frac{1}{3}n(n^2 + 9n + 26)$</p> $\frac{1}{3}n(8n^2 + 36n + 52 - n^2 - 9n - 26) = \frac{1}{3}n(7n^2 + 27n + 26) \quad (*)$	<p>M1 A1ft A1cso (3) 14 marks</p>
	<p>Notes:</p> <p>(a) B1: Checks $n = 1$ on both sides and states true for $n = 1$ here or in conclusion M1: Assumes true for $n = k$ (should use one of these two words) M1: Adds $(k+1)$th term to sum of k terms A1: Correct work to support proof M1: Deduces $\frac{1}{6}n(n+1)(2n+1)$ with $n = k + 1$ A1: Makes induction statement. Statement true for $n = 1$ here could contribute to B1 mark earlier</p>	

Question 9 Notes continued:

(b) M1: Expands and splits (but allow 6 rather than sigma 6 for this mark)

A1: first two terms correct

B1: for $6n$

M1: Take out factor $n/6$ or $n/3$ correctly – no errors factorising

A1: for correct factorised cubic or for identifying a and b

(c) M1: Try to use $\sum_1^{2n} (r+2)(r+3) - \sum_1^n (r+2)(r+3)$ with previous result used **at least once**

A1ft Two correct expressions for their a and b values

A1: Completely correct work to printed answer

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email publications@linneydirect.com

Order Code UA023925 Summer 2010

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Edexcel Limited. Registered in England and Wales no.4496750
Registered Office: One90 High Holborn, London, WC1V 7BH