



Pearson

# Mark Scheme (Results)

Summer 2017

Pearson Edexcel GCE

In Further Pure Mathematics FP1 (6667/01)

## Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2017

Publications Code 6667\_01\_1706\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2017

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

**PEARSON EDEXCEL GCE MATHEMATICS****General Instructions for Marking**

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
- ft – follow through
- the symbol  $\surd$  will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- d... or dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper or ag- answer given
- $\square$  or d... The second mark is dependent on gaining the first mark

4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft.
5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

**General Principles for Further Pure Mathematics Marking**

*(But note that specific mark schemes may sometimes override these general principles).*

**Method mark for solving 3 term quadratic:****1. Factorisation**

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

**2. Formula**

Attempt to use the correct formula (with values for a, b and c).

**3. Completing the square**

Solving  $x^2 + bx + c = 0$ :  $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$ ,  $q \neq 0$ , leading to  $x = \dots$

**Method marks for differentiation and integration:****1. Differentiation**

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

**2. Integration**

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

Question Number	Scheme	Marks
1.	$f(x) = \frac{1}{3}x^2 + \frac{4}{x^2} - 2x - 1, \quad x > 0$	
(a)	$f(6) = -0.88888888...$ $f(7) = 1.414965986...$ Sign change or $f(6) = -ve$ and $f(7) = +ve$ or $f(6) \times f(7) = -ve$ o.e. (and $f(x)$ is continuous) therefore a root $\alpha$ (exists between $x = 6$ and $x = 7$ ) o.e.	Either any one of $f(6) = \text{awrt } -0.9$ or $f(7) = \text{awrt } 1.4$ Both $f(6) = \text{awrt } -0.9$ and $f(7) = \text{awrt } 1.4$ , sign change and conclusion. Allow $f(6) = -\frac{8}{9}$ and $f(7) = \frac{208}{147}$ .
(b)	$f'(x) = \frac{2}{3}x - \frac{8}{x^3} - 2$ $\{f'(6) = 1.962962963...\}$ $\alpha \approx 6 - \left( \frac{"-0.88888888..."}{"1.962962963..."} \right)$ $= 6.452830189...$ $= 6.45 \text{ (2dp)}$	$\frac{1}{3}x^2 \rightarrow \pm Ax \text{ or } \frac{4}{x^2} \rightarrow \pm Bx^{-3} \text{ or } -2x - 1 \rightarrow -2$ At least two of these terms differentiated correctly. Correct derivative. $f'(6) = \frac{53}{27}$ Correct application of Newton-Raphson using their values. Exact form of $\alpha$ is $\frac{342}{53}$ 6.45
		M1 A1 A1 M1 A1 <b>cso</b> <b>[5]</b> <b>7</b>
<b>Question 1 Notes</b>		
1. (a)	<b>Note</b> Accept at least 'sign change therefore root' o.e. for A1. Any incorrect statements made in the conclusion award A0.	
(b)	<b>Note</b> Denominator in NR calculation may contain evidence for first 3 marks. Correct answer of 6.45 with minimal working will imply earlier marks for elements not explicitly stated. However, incorrect values leading to a correct final answer should be marked accordingly.	

Question Number	Scheme	Marks
<b>2.</b>	$\mathbf{A} = \begin{pmatrix} 2 & -1 \\ 4 & 3 \end{pmatrix}, \mathbf{P} = \begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix}$	
(a)	$\mathbf{A}^{-1} = \frac{1}{10} \begin{pmatrix} 3 & 1 \\ -4 & 2 \end{pmatrix}$	<p>Either <math>\frac{1}{10}</math> or <math>\begin{pmatrix} 3 &amp; 1 \\ -4 &amp; 2 \end{pmatrix}</math> M1</p> <p>Correct matrix seen. A1</p>
(b)	$\mathbf{P} = \mathbf{AB}$	
<b>Way 1</b>	$\Rightarrow \mathbf{A}^{-1}\mathbf{P} = \mathbf{A}^{-1}\mathbf{AB} \Rightarrow \mathbf{B} = \mathbf{A}^{-1}\mathbf{P}$ $\mathbf{B} = \frac{1}{10} \begin{pmatrix} 3 & 1 \\ -4 & 2 \end{pmatrix} \begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix}$ $= \begin{pmatrix} 2 & 1 \\ 1 & -4 \end{pmatrix}$	<p>Multiplies <b>their</b> <math>\mathbf{A}^{-1}</math> by <math>\mathbf{P}</math> in correct order. M1  This substituted statement is sufficient.</p> <p>At least 2 elements correct or <math>k \begin{pmatrix} 20 &amp; 10 \\ 10 &amp; -40 \end{pmatrix}</math> oe. A1</p> <p>May be unsimplified  Correct simplified matrix. A1</p>
(b)	$\{\mathbf{P} = \mathbf{AB} \Rightarrow\}$	
<b>Way 2</b>	$\begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ $\begin{pmatrix} 3 & 6 \\ 11 & -8 \end{pmatrix} = \begin{pmatrix} 2a-c & 2b-d \\ 4a+3c & 4b+3d \end{pmatrix}$ $\Rightarrow a=2, c=1, b=1, d=-4$ So, $\mathbf{B} = \begin{pmatrix} 2 & 1 \\ 1 & -4 \end{pmatrix}$	<p>Attempt to multiply <math>\mathbf{A}</math> by <math>\mathbf{B}</math> in the correct order and puts equal to <math>\mathbf{P}</math> M1</p> <p>At least 2 elements are correct. A1</p> <p>Correct matrix. A1</p>
		[3]
		[3]
		5



Question Number	Scheme	Marks												
3. (a)	$x = 4t, y = \frac{4}{t}, t \neq 0$ $t = \frac{1}{4} \Rightarrow P(1, 16), t = 2 \Rightarrow Q(8, 2)$ $m(PQ) = \frac{2-16}{8-1} \{ = -2 \}$ $m(l) = \frac{1}{2}$ So, $l: y = \frac{1}{2}x$ or $2y = x$	B1 M1 Finds the gradient of the chord $PQ$ with $\frac{y_2 - y_1}{x_2 - x_1}$ then uses in $y = -\frac{1}{m}x$ . Condone incorrect sign of gradient. A1 oe [3]												
(b)	$xy = 16$ or $y = \frac{16}{x}$ or $x = \frac{16}{y}$	Correct Cartesian equation. Accept $\frac{4}{y} = \frac{x}{4}$ or $xy = 4^2$ B1 oe [1]												
(c)	<table border="1"> <thead> <tr> <th>Way 1</th><th>Way 2</th><th>Way 3</th></tr> </thead> <tbody> <tr> <td><math>\frac{1}{2}x = \frac{16}{x}</math></td><td><math>\frac{4}{t} = \frac{1}{2}(4t)</math></td><td><math>2y = \frac{16}{y}</math></td></tr> <tr> <td><math>\{x^2 = 32\}</math></td><td><math>\{t^2 = 2\}</math></td><td><math>\{y^2 = 8\}</math></td></tr> <tr> <td colspan="3"><math>(4\sqrt{2}, 2\sqrt{2}), (-4\sqrt{2}, -2\sqrt{2})</math></td></tr> </tbody> </table>	Way 1	Way 2	Way 3	$\frac{1}{2}x = \frac{16}{x}$	$\frac{4}{t} = \frac{1}{2}(4t)$	$2y = \frac{16}{y}$	$\{x^2 = 32\}$	$\{t^2 = 2\}$	$\{y^2 = 8\}$	$(4\sqrt{2}, 2\sqrt{2}), (-4\sqrt{2}, -2\sqrt{2})$			Attempts to substitute their $l$ into either their Cartesian equation or parametric equations of $H$ M1 At least one set of coordinates (simplified or un-simplified) or $x = \pm 4\sqrt{2}, y = \pm 2\sqrt{2}$ A1 Both sets of simplified coordinates. Accept written in pairs as $x = 4\sqrt{2}, y = 2\sqrt{2}$ $x = -4\sqrt{2}, y = -2\sqrt{2}$ A1 [3]
Way 1	Way 2	Way 3												
$\frac{1}{2}x = \frac{16}{x}$	$\frac{4}{t} = \frac{1}{2}(4t)$	$2y = \frac{16}{y}$												
$\{x^2 = 32\}$	$\{t^2 = 2\}$	$\{y^2 = 8\}$												
$(4\sqrt{2}, 2\sqrt{2}), (-4\sqrt{2}, -2\sqrt{2})$														
		7												

Question Number	Scheme	Marks
4. (i)	Mark (i)(a) and (i)(b) together.	
(a) Way 1	$w = \frac{p-4i}{2-3i} \quad \arg w = \frac{\pi}{4}$ $w = \frac{(p-4i)}{(2-3i)} \times \frac{(2+3i)}{(2+3i)}$ $= \left( \frac{2p+12}{13} \right) + \left( \frac{3p-8}{13} \right)i$	Multiplies by $\frac{(2+3i)}{(2+3i)}$ M1 At least one of either the real or imaginary part of $w$ is correct. Must be expanded but could be unsimplified e.g. expressed as single fraction. Condone $a+ib$ . A1 Correct $w$ in its simplest form. A1 <b>[3]</b>
(a) Way 2	$(a+ib)(2-3i) = (p-4i)$ $2a+3b = p$ $3a-2b = 4$ $= \left( \frac{2p+12}{13} \right) + \left( \frac{3p-8}{13} \right)i$	Multiplies out to obtain 2 equations in two unknowns. M1 At least one of either the real or imaginary part of $w$ is correct. Must be expanded but could be unsimplified e.g. expressed as single fraction. Condone $a+ib$ . A1 Correct $w$ in its simplest form. A1 <b>[3]</b>
(b)	$\left\{ \arg w = \frac{\pi}{4} \Rightarrow \right\} \quad 2p+12=3p-8 \text{ o.e. seen anywhere.}$ $\Rightarrow p=20$	Sets the numerators of the real part of their $w$ equal to the imaginary part of their $w$ or if arctan used, require evidence of $\tan \frac{\pi}{4} = 1$ $p=20$ A1 <b>[2]</b>
(ii) Way 1	$z = (1-\lambda i)(4+3i) \text{ and }  z  = 45$ $\sqrt{1+\lambda^2} \sqrt{4^2+3^2}$ $\sqrt{1+\lambda^2} \sqrt{4^2+3^2} = 45$ $\{\lambda^2 = 9^2 - 1 \Rightarrow \} \quad \lambda = \pm 4\sqrt{5}$	Attempts to apply $ (1-\lambda i)(4+3i)  = \sqrt{1+\lambda^2} \sqrt{4^2+3^2}$ M1 Correct equation. A1 $\lambda = \pm 4\sqrt{5}$ A1 <b>[3]</b>
Way 2	$z = (4+3\lambda) + (3-4\lambda)i$ $\sqrt{(4+3\lambda)^2 + (3-4\lambda)^2}$ $(4+3\lambda)^2 + (3-4\lambda)^2 = 45^2 \text{ or}$ $\sqrt{(4+3\lambda)^2 + (3-4\lambda)^2} = 45$ $\{16+24\lambda+9\lambda^2+9-24\lambda+16\lambda^2 = 2025\}$ $\{25\lambda^2 = 2000 \Rightarrow \} \quad \lambda = \pm 4\sqrt{5}$	Attempt to multiply out, group real and imaginary parts and apply the modulus. M1 Correct equation. A1 Condone if middle terms in expansions not explicitly stated. A1 <b>[3]</b>
<b>Question 4 Notes</b>		
(ii)	<b>M1</b> Also allow $(1+\lambda^2)(4^2+3^2)$ for M1. <b>M1</b> Also allow $(4+3\lambda)^2 + (3-4\lambda)^2$ for M1.	<b>[3]</b> <b>8</b>

Question Number	Scheme	Marks
5. (i)	$\mathbf{A} = \begin{pmatrix} p & 2 \\ 3 & p \end{pmatrix}, \mathbf{B} = \begin{pmatrix} -5 & 4 \\ 6 & -5 \end{pmatrix}, \mathbf{M} = \begin{pmatrix} a & -9 \\ 1 & 2 \end{pmatrix}$ $p, a$ are constants.	
(a)	$\{\mathbf{AB}\} = \begin{pmatrix} -5p+12 & 4p-10 \\ -15+6p & 12-5p \end{pmatrix}$ At least 2 elements are correct. Correct matrix.	M1 A1 [2]
(b)	$\{\mathbf{AB} + 2\mathbf{A} = k\mathbf{I}\}$ $\begin{pmatrix} -5p+12 & 4p-10 \\ -15+6p & 12-5p \end{pmatrix} + 2\begin{pmatrix} p & 2 \\ 3 & p \end{pmatrix} = k\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ If 'simultaneous equations' used, allocate marks as below. $\begin{pmatrix} -3p+12 & 4p-6 \\ -9+6p & 12-3p \end{pmatrix} = \begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$ $"4p-10" + 4 = 0$ or $"-15+6p" + 6 = 0$ Forms an equation in $p$ or $"-9+6p" = "4p-6"$ $\Rightarrow p = \frac{3}{2}$ $p = \frac{3}{2}$ o.e. $k = -5\left(\frac{3}{2}\right) + 12 + 2\left(\frac{3}{2}\right) \Rightarrow k = \dots$ Substitutes their $p = \frac{3}{2}$ into "their $(-5p+12)" + 2p$ to find a value for $k$ or eliminates $p$ to find $k$ . $k = \frac{15}{2}$ $k = \frac{15}{2}$ oe	M1 A1 M1 A1 [4]
(ii) Way 1	$\pm \frac{270}{15} \{ = \pm 18 \}$ Can be implied from calculations. $\det \mathbf{M} = (a)(2) - (-9)(1)$ Applies $ad - bc$ to $\mathbf{M}$ . Require clear evidence of correct formula being used for M1 if errors seen. $\Rightarrow 2a+9 = 18$ or $2a+9 = -18$ Equates their $\det \mathbf{A}$ to either 18 or -18 $\Rightarrow a = 4.5$ or $a = -13.5$ At least one of either $a = 4.5$ or $a = -13.5$ Both $a = 4.5$ and $a = -13.5$	B1 M1 M1 A1 A1 [5]
(ii) Way 2	Consider vertices of triangle with area 15 units e.g. (0,0), (15,0) and (0,2) and attempting 2 values of $a$ . e.g. $\begin{pmatrix} a & -9 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 0 & 15 & 0 \\ 0 & 0 & 2 \end{pmatrix} = \begin{pmatrix} 0 & 15a & -18 \\ 0 & 15 & 4 \end{pmatrix}$ Pre-multiplies their matrix by $\mathbf{M}$ and obtains single matrix e.g. $\frac{1}{2} \begin{vmatrix} 0 & 15a & -18 & 0 \\ 0 & 15 & 4 & 0 \end{vmatrix} = 270$ Equates their determinant to 270 and attempts to solve. $\Rightarrow a = 4.5$ or $a = -13.5$ At least one of either $a = 4.5$ or $a = -13.5$ Both $a = 4.5$ and $a = -13.5$	M1 M1 A1 A1 [5] 11

Question Number	Scheme	Marks
<b>6.</b>	$x^3 + ax^2 + bx - 52 = 0$ , $a, b \in \mathbb{R}$ , 4 and $2i - 3$ are roots	
(a)	$-2i - 3$ <span style="float: right;"><math>-2i - 3</math> seen anywhere in solution for Q6.</span>	B1
<b>Way 1</b>	$(x - (2i - 3))(x - "(-2i - 3)"); = x^2 + 6x + 13$ or $x = -3 \pm 2i \Rightarrow (x + 3)^2 = -4; = x^2 + 6x + 13 (= 0)$ $(x - 4)(x - (2i - 3)); = x^2 - (1 + 2i)x + 4(2i - 3)$ $(x - 4)(x - "(-2i - 3)"); = x^2 - (1 - 2i)x + 4(-2i - 3)$  $(x - 4)(x^2 + 6x + 13) \{ = x^3 + ax^2 + bx - 52 \}$ $a = 2, b = -11$ or $x^3 + 2x^2 - 11x - 52$	<p>Must follow from their part (a). Any incorrect signs for their part (a) in initial statement award M0; accept any equivalent expanded expression for A1.</p> <p><math>(x - 3^{\text{rd}} \text{ root})(\text{their quadratic})</math>.          Could be found by comparing coefficients from long division.          At least one of <math>a = 2</math> or <math>b = -11</math>          Both <math>a = 2</math> and <math>b = -11</math></p>
<b>Way 2</b>	Sum = $(2i - 3) + "(-2i - 3)" = -6$ Product = $(2i - 3) \times "(-2i - 3)" = 13$ So quadratic is $x^2 + 6x + 13$	<p>Attempts to apply either  <math>x^2 - (\text{sum roots})x + (\text{product roots}) = 0</math>          or <math>x^2 - 2\text{Re}(\alpha)x +  \alpha^2  = 0</math>  <math>x^2 + 6x + 13</math></p>
<b>Way 3</b>	$(2i - 3)^3 + a(2i - 3)^2 + b(2i - 3) - 52 = 0$ $5a - 3b = 43$ (real parts) and $6a - b = 23$ (imaginary parts) or uses $f(4) = 0$ and $f(\text{a complex root}) = 0$ to form equations in $a$ and $b$ .  So $a = 2, b = -11$ or $x^3 + 2x^2 - 11x - 52$	<p>Substitutes <math>2i - 3</math> into the displayed equation and equates both real and imaginary parts.  <math>5a - 3b = 43</math> and <math>6a - b = 23</math> or  <math>16a + 4b = -12</math> and  <math>(2i - 3)^3 + a(2i - 3)^2 + b(2i - 3) - 52 = 0 /</math>  <math>(-2i - 3)^3 + a(-2i - 3)^2 + b(-2i - 3) - 52 = 0</math>          Solves these equations simultaneously to find at least one of either <math>a = \dots</math> or <math>b = \dots</math>          At least one of <math>a = 2</math> or <math>b = -11</math>          Both <math>a = 2</math> and <math>b = -11</math></p>
<b>Way 4</b>	$b = \text{sum of product pairs}$ $= 4(2i - 3) + 4"(-2i - 3)" + (2i - 3)"(-2i - 3)"$ $a = -(\text{sum of 3 roots}) = -(4 + 2i - 3 - 2i - 3)"$ $a = 2, b = -11$ or $x^3 + 2x^2 - 11x - 52$	<p>Attempts sum of product pairs.          All pairs correct o.e.          Adds up all 3 roots          At least one of <math>a = 2</math> or <math>b = -11</math>          Both <math>a = 2</math> and <math>b = -11</math></p>

(b) <b>Way 5</b>	Uses $f(4) = 0$		M1
	$16a + 4b = -12$		A1
	$a = -(\text{sum of 3 roots}) = -(4 + 2i - 3 - 2i - 3)$	Adds up all 3 roots	M1
	$a = 2, b = -11$ or $x^3 + 2x^2 - 11x - 52$	At least one of $a = 2$ or $b = -11$	A1
		Both $a = 2$ and $b = -11$	A1
			<b>[5]</b>
			<b>6</b>

Question Number	Scheme	Marks
7.	$y^2 = 4ax$ , at $Q(aq^2, 2aq)$	
(a)	$y = 2\sqrt{a}x^{\frac{1}{2}} \Rightarrow \frac{dy}{dx} = \sqrt{a}x^{-\frac{1}{2}}$ or $2y\frac{dy}{dx} = 4a$ or $\frac{dy}{dx} = 2a \times \frac{1}{2aq}$	$\frac{dy}{dx} = \pm kx^{-\frac{1}{2}}$ or $ky\frac{dy}{dx} = c$ or their $\frac{dy}{dq}$ their $\frac{dx}{dq}$ A1
	When $x = aq^2$ , $m_T = \frac{dy}{dx} = \frac{\sqrt{a}}{\sqrt{aq^2}} = \frac{\sqrt{a}}{\sqrt{a}q} = \frac{1}{q}$ or when $y = 2aq$ , $m_T = \frac{dy}{dx} = \frac{4a}{2(2aq)} = \frac{1}{q}$	$\frac{dy}{dx} = \frac{1}{q}$ A1
	<b>T:</b> $y - 2aq = \frac{1}{q}(x - aq^2)$	Applies $y - 2aq = (\text{their } m_T)(x - aq^2)$ or $y = (\text{their } m_T)x + c$ and an attempt to find $c$ with gradient from calculus. dM1
	<b>T:</b> $qy - 2aq^2 = x - aq^2$ <b>T:</b> $qy = x + aq^2$	cso A1 * [4]
(b)	$X(-\frac{1}{4}a, 0) \Rightarrow 0 = -\frac{1}{4}a + aq^2$ $\Rightarrow \left\{ q^2 = \frac{1}{4} \Rightarrow q = -\frac{1}{2} \text{ (reject)} \right\} q = \frac{1}{2}$ So, $\frac{1}{2}y = -a + a\left(\frac{1}{2}\right)^2$	Substitutes $x = -\frac{1}{4}a$ and $y = 0$ into <b>T</b> $q = \frac{1}{2}$ oe A1 Substitutes their " $q = \frac{1}{2}$ " and $x = -a$ in <b>T</b> or finds $y_D = \frac{1}{q}(-a + aq^2)$ $D(-a, -\frac{3}{2}a)$ o.e. A1 [4]
(c)	{focus $F(a, 0)$ }	
Way 1	$\text{Area}(FXD) = \frac{1}{2}\left(\frac{5a}{4}\right)\left(\frac{3a}{2}\right) = \frac{15a^2}{16}$	Applies $\frac{1}{2}(\text{their }  FX )(\text{their }  y_D )$ . If their $\left y_D = \frac{1}{q}(-a + aq^2)\right $ then require an attempt to sub for $q$ to award M. $\frac{15a^2}{16}$ or $0.9375a^2$ A1 cso [2]

(c) <b>Way 2</b>	$\text{Area}(FXD) = \frac{1}{2} \begin{vmatrix} a & -\frac{1}{4}a & -a & a \\ 0 & 0 & -\frac{3}{2}a & 0 \end{vmatrix}$ $= \frac{1}{2} \left  \left( 0 + \frac{3}{8}a^2 + 0 \right) - \left( 0 + 0 - \frac{3}{2}a^2 \right) \right  = \frac{15}{16}a^2$	<p>A correct attempt to apply the shoelace method.</p> $\frac{15a^2}{16} \text{ or } 0.9375a^2$	<p>M1</p> <p>A1cao</p> <p>[2]</p>
(c) <b>Way 3</b>	<p>Rectangle – triangle 1 – triangle 2</p> $= 2a \cdot \frac{3a}{2} - \frac{1}{2} \cdot \frac{3a}{4} \cdot \frac{3a}{2} - \frac{1}{2} \cdot 2a \cdot \frac{3a}{2} = 3a^2 - \frac{9a^2}{16} - \frac{3a^2}{2}$	$\frac{15a^2}{16} \text{ or } 0.9375a^2$	<p>M1</p> <p>A1cao</p>
(c) <b>Way 4</b>	<p>Attempts sine rule using appropriate choice from</p> $FX = \frac{5a}{4}, FD = \frac{5a}{2}, DX = \frac{3\sqrt{5}a}{4}, \sin F = \frac{3}{5}, \sin X = \frac{2}{\sqrt{5}}$	<p>Uses Area = <math>\frac{1}{2}ab \sin C</math></p> $\frac{15a^2}{16} \text{ or } 0.9375a^2$	<p>M1</p> <p>A1cao</p> <p><b>10</b></p>

Question 7 Notes	
(c) <b>Way 1</b>	<p>Do not award M1 if area of wrong triangle found e.g. <math>\frac{1}{2} \cdot 2a \cdot \frac{3a}{2} = \frac{3a^2}{2}</math></p>

Question Number	Scheme	Marks
8. (a)	$\sum_{r=1}^n (3r^2 + 8r + 3)$ $= \frac{3}{6}n(n+1)(2n+1) + \frac{8}{2}n(n+1) + 3n$ $= \frac{1}{2}n(n+1)(2n+1) + 4n(n+1) + 3n$ $= \frac{1}{2}n((2n+1)(n+1) + 8(n+1) + 6)$ $= \frac{1}{2}n(2n^2 + 3n + 1 + 8n + 8 + 6)$ $= \frac{1}{2}n(2n^2 + 11n + 15)$ $= \frac{1}{2}n(2n+5)(n+3) \quad (*)$ <p>An attempt to use at least one of the correct standard formulae for first two terms. Correct first two terms. <math>3 \rightarrow 3n</math></p> <p>Factorise out at least <math>n</math> from all terms at any point. There must be a factor of <math>n</math> in every term.</p> <p>Achieves the correct answer, no errors seen.</p>	<p>M1 A1 B1</p> <p>M1</p> <p>A1*cso [5]</p>
(b)	$\sum_{r=1}^{12} (3r^2 + 8r + 3 + k(2^{r-1})) = 3520$ $\sum_{r=1}^{12} (3r^2 + 8r + 3) = \frac{1}{2}(12)(29)(15) \{= 2610\}$ $\sum_{r=1}^{12} (2^{r-1}) = \frac{1(1-2^{12})}{1-2} \{= 4095\}$ <p>Attempt to evaluate <math>\sum_{r=1}^{12} (3r^2 + 8r + 3)</math></p> <p>Attempt to apply the sum to 12 terms of a GP or adds up all 12 terms.</p> $\frac{1(1-2^{12})}{1-2} \text{ o.e. or } 4095.$ <p>So, <math>2610 + 4095k = 3520 \Rightarrow 4095k = 910</math></p> <p>giving, <math>k = \frac{2}{9}</math></p> <p><math>k = \frac{2}{9}</math> or 0.2</p>	<p>M1 M1 A1 A1 A1 [4] 9</p>
<b>Question 8 Notes</b>		
8. (b)	<b>Note</b> 2 <sup>nd</sup> M1 1 <sup>st</sup> A1: These two marks can be implied by seeing 4095 or 4095k	



Question Number	Scheme	Marks
9.		
(i)	$u_{n+2} = 6u_{n+1} - 9u_n, n \geq 1, u_1 = 6, u_2 = 27; u_n = 3^n(n+1)$ $n=1; u_1 = 3(2) = 6$ $n=2; u_2 = 3^2(2+1) = 27$ So $u_n$ is true when $n=1$ and $n=2$ . Assume that $u_k = 3^k(k+1)$ and $u_{k+1} = 3^{k+1}(k+2)$ are true.  Then $u_{k+2} = 6u_{k+1} - 9u_k$ $= 6(3^{k+1})(k+2) - 9(3^k)(k+1)$  $= 2(3^{k+2})(k+2) - (3^{k+2})(k+1)$ $= (3^{k+2})(2k+4-k-1)$ $= (3^{k+2})(k+3)$ $= (3^{k+2})(k+2+1)$ If the result is <b>true for <math>n=k</math> and <math>n=k+1</math></b> then it is now <b>true for <math>n=k+2</math></b> . As it is <b>true for <math>n=1</math> and <math>n=2</math></b> then it is <b>true for all <math>n</math></b> ( $\in \mathbb{Z}^+$ ).	Check that $u_1 = 6$ and $u_2 = 27$  Could assume for $n=k, n=k-1$ and show for $n=k+1$  Substituting $u_k$ and $u_{k+1}$ into $u_{k+2} = 6u_{k+1} - 9u_k$ Correct expression Achieves an expression in $3^{k+2}$  $(3^{k+2})(k+2+1)$ or $(3^{k+2})(k+3)$ Correct conclusion seen at the end. Condone true for $n=1$ and $n=2$ seen anywhere. This should be compatible with assumptions.
(ii)	$f(n) = 3^{3n-2} + 2^{3n+1}$ is divisible by 19 <b>In all ways, first M is for applying <math>f(k+1)</math> with at least 1 power correct. The second M is dependent on at least one accuracy being awarded and making <math>f(k+1)</math> the subject and the final A is correct solution only.</b>	
(ii)	$f(1) = 3^1 + 2^4 = 19$ {which is divisible by 19}. $\therefore f(n)$ is divisible by 19 when $n=1$ } { Assume that for $n=k$ , $f(k) = 3^{3k-2} + 2^{3k+1}$ is divisible by 19 for $k \in \mathbb{Z}^+$ . } $f(k+1) - f(k) = 3^{3(k+1)-2} + 2^{3(k+1)+1} - (3^{3k-2} + 2^{3k+1})$ $f(k+1) - f(k) = 27(3^{3k-2}) + 8(2^{3k+1}) - (3^{3k-2} + 2^{3k+1})$ $f(k+1) - f(k) = 26(3^{3k-2}) + 7(2^{3k+1})$ $= 7(3^{3k-2} + 2^{3k+1}) + 19(3^{3k-2})$ <b>or</b> $= 26(3^{3k-2} + 2^{3k+1}) - 19(2^{3k+1})$ $= 7f(k) + 19(3^{3k-2})$ <b>or</b> $= 26f(k) - 19(2^{3k+1})$ $\therefore f(k+1) = 8f(k) + 19(3^{3k-2})$ <b>or</b> $f(k+1) = 27f(k) - 19(2^{3k+1})$  $\therefore f(k+1) = 8f(k) + 19(3^{3k-2})$ is divisible by 19 as both $8f(k)$ and $19(3^{3k-2})$ are both divisible by 19 }	Shows $f(1) = 19$  Applies $f(k+1)$ with at least 1 power correct  Either $7(3^{3k-2} + 2^{3k+1})$ or $7f(k); 19(3^{3k-2})$ or $26(3^{3k-2} + 2^{3k+1})$ or $26f(k); -19(2^{3k+1})$  Dependent on at least one of the previous accuracy marks being awarded. Makes Applies $f(k+1)$ with at least 1 power correct the subject
Way 1		

	<p>If the result is <b>true for <math>n = k</math></b>, then it is now <b>true for <math>n = k + 1</math></b>. As the result has shown to be <b>true for <math>n = 1</math></b>, then the result is <b>true for all <math>n \in \mathbb{Z}^+</math></b>.</p> <p>(ii) <b>Way 2</b></p> <p><math>f(1) = 3^1 + 2^4 = 19</math> { which is divisible by 19 }.</p> <p>{ <math>\therefore f(n)</math> is divisible by 19 when <math>n = 1</math> }</p> <p>Assume that for <math>n = k</math>,</p> <p><math>f(k) = 3^{3k-2} + 2^{3k+1}</math> is divisible by 19 for <math>k \in \mathbb{Z}^+</math>.</p> <p><math>f(k+1) = 3^{3(k+1)-2} + 2^{3(k+1)+1}</math></p> <p><math>f(k+1) = 27(3^{3k-2}) + 8(2^{3k+1})</math></p> <p><math>= 8(3^{3k-2} + 2^{3k+1}) + 19(3^{3k-2})</math></p> <p><b>or</b> <math>= 27(3^{3k-2} + 2^{3k+1}) - 19(2^{3k+1})</math></p> <p><math>\therefore f(k+1) = 8f(k) + 19(3^{3k-2})</math></p> <p><b>or</b> <math>f(k+1) = 27f(k) - 19(2^{3k+1})</math></p> <p>{ <math>\therefore f(k+1) = 8f(k) + 19(3^{3k-2})</math> is divisible by 19 as both <math>8f(k)</math> and <math>19(3^{3k-2})</math> are both divisible by 19 }</p> <p>If the result is <b>true for <math>n = k</math></b>, then it is now <b>true for <math>n = k + 1</math></b>. As the result has shown to be <b>true for <math>n = 1</math></b>, then the result is <b>true for all <math>n \in \mathbb{Z}^+</math></b>.</p> <p>(ii) <b>Way 3</b></p> <p><math>f(n) = 3^{3n-2} + 2^{3n+1}</math> is divisible by 19</p> <p><math>f(1) = 3^1 + 2^4 = 19</math> { which is divisible by 19 }.</p> <p>{ <math>\therefore f(n)</math> is divisible by 19 when <math>n = 1</math> }</p> <p>Assume that for <math>n = k</math>,</p> <p><math>f(k) = 3^{3k-2} + 2^{3k+1}</math> is divisible by 19 for <math>k \in \mathbb{Z}^+</math>.</p> <p><math>f(k+1) - \alpha f(k) = 3^{3(k+1)-2} + 2^{3(k+1)+1} - \alpha(3^{3k-2} + 2^{3k+1})</math></p> <p><math>f(k+1) - \alpha f(k) = (27 - \alpha)(3^{3k-2}) + (8 - \alpha)2^{3k+1}</math></p> <p><math>= (8 - \alpha)(3^{3k-2} + 2^{3k+1}) + 19(3^{3k-2})</math></p> <p><b>or</b> <math>= (27 - \alpha)(3^{3k-2} + 2^{3k+1}) - 19(2^{3k+1})</math></p> <p><math>\therefore f(k+1) = 8f(k) + 19(3^{3k-2})</math></p> <p><b>or</b> <math>f(k+1) = 27f(k) - 19(2^{3k+1})</math></p> <p>{ <math>\therefore f(k+1) = 27f(k) - 19(2^{3k+1})</math> is divisible by 19 as both <math>27f(k)</math> and <math>19(2^{3k+1})</math> are both divisible by 19 }</p> <p>If the result is <b>true for <math>n = k</math></b>, then it is now <b>true for <math>n = k + 1</math></b>. As the result has shown to be <b>true for <math>n = 1</math></b>, then the result is <b>true for all <math>n \in \mathbb{Z}^+</math></b>.</p>	<p>Correct conclusion seen at the end. Condoned true for <math>n = 1</math> stated earlier.</p> <p>Shows <math>f(1) = 19</math></p> <p>Applies <math>f(k+1)</math> with at least 1 power correct</p> <p>Either <math>8(3^{3k-2} + 2^{3k+1})</math> or <math>8f(k); 19(3^{3k-2})</math> or <math>27(3^{3k-2} + 2^{3k+1})</math> or <math>27f(k); -19(2^{3k+1})</math></p> <p><b>Dependent on at least one of the previous accuracy marks being awarded.</b></p> <p>Correct conclusion seen at the end. Condoned true for <math>n = 1</math> stated earlier.</p> <p>Shows <math>f(1) = 19</math></p> <p>Applies <math>f(k+1)</math> with at least 1 power correct</p> <p><math>(8 - \alpha)(3^{3k-2} + 2^{3k+1})</math> or <math>(8 - \alpha)f(k); 19(3^{3k-2})</math> NB choosing <math>\alpha = 8</math> makes first term disappear. <math>(27 - \alpha)(3^{3k-2} + 2^{3k+1})</math> or <math>(27 - \alpha)f(k); -19(2^{3k+1})</math> NB choosing <math>\alpha = 27</math> makes first term disappear.</p> <p><b>Dependent on at least one of the previous accuracy marks being awarded.</b> Makes <math>f(k+1)</math> the subject.</p> <p>Correct conclusion seen at the end. Condoned true for <math>n = 1</math> stated earlier.</p>	<p>A1 cso</p> <p>[6]</p> <p>B1</p> <p>M1</p> <p>A1; A1 dM1</p> <p>A1 cso</p> <p>[6]</p> <p>B1</p> <p>M1</p> <p>A1; A1</p> <p>dM1</p> <p>A1 cso</p> <p>[6] 12</p>
	<p align="center"><b>Question 9 Notes</b></p>		
<p>(ii)</p>	<p>Accept use of <math>f(k) = 3^{3k-2} + 2^{3k+1} = 19m</math> o.e. and award method and accuracy as above.</p>		

