

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	6	/	0	1	Signature	

Paper Reference(s)

**6666/01**

# Edexcel GCE

## Core Mathematics C4

### Advanced

Tuesday 16 June 2015 – Afternoon  
Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

### Materials required for examination

---

### Mathematical Formulae (Pink)

### Items included with question papers

Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.**

### Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer for each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 32 pages in this question paper. Any blank pages are indicated.

## Advice to Candidates

---

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.  
©2015 Pearson Education Ltd

Printer's Log. No.

Printer's Log. No.  
P44827A

5/1/1/1/



*Turn over*

PEARSON

1. (a) Find the binomial expansion of

$$(4 + 5x)^{\frac{1}{2}}, \quad |x| < \frac{4}{5}$$

(5)

- (b) Find the exact value of  $(4 + 5x)^{\frac{1}{2}}$  when  $x = \frac{1}{10}$

(1)

- (c) Substitute  $x = \frac{1}{10}$  into your binomial expansion from part (a) and hence find an approximate value for  $\sqrt{2}$

Give your answer in the form  $\frac{p}{q}$  where  $p$  and  $q$  are integers.

(2)



Q1

**(Total 8 marks)**



**2.** The curve  $C$  has equation

$$x^2 - 3xy - 4y^2 + 64 = 0$$

- (a) Find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .

(5)

- (b) Find the coordinates of the points on  $C$  where  $\frac{dy}{dx} = 0$

*(Solutions based entirely on graphical or numerical methods are not acceptable.)*

(6)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Leave  
blank



3.




Figure 1

Figure 1 shows a sketch of part of the curve with equation  $y = 4x - xe^{\frac{1}{2}x}$ ,  $x \geq 0$

The curve meets the  $x$ -axis at the origin  $O$  and cuts the  $x$ -axis at the point  $A$ .

(a) Find, in terms of  $\ln 2$ , the  $x$  coordinate of the point  $A$ .

(2)

(b) Find

$$\int xe^{\frac{1}{2}x} dx$$

(3)

The finite region  $R$ , shown shaded in Figure 1, is bounded by the  $x$ -axis and the curve with equation

$$y = 4x - xe^{\frac{1}{2}x}, \quad x \geq 0$$

(c) Find, by integration, the exact value for the area of  $R$ .  
Give your answer in terms of  $\ln 2$

(3)

---

---

---

---

---

---

---

---

---

---



Leave  
blank

Question 3 continued

Lined area for writing answers.





**Q3**

1

**(Total 8 marks)**



4. With respect to a fixed origin  $O$ , the lines  $l_1$  and  $l_2$  are given by the equations

$$l_1: \mathbf{r} = \begin{pmatrix} 5 \\ -3 \\ p \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 1 \\ -3 \end{pmatrix}, \quad l_2: \mathbf{r} = \begin{pmatrix} 8 \\ 5 \\ -2 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ 4 \\ -5 \end{pmatrix}$$

where  $\lambda$  and  $\mu$  are scalar parameters and  $p$  is a constant.

The lines  $l_1$  and  $l_2$  intersect at the point  $A$ .

- (a) Find the coordinates of  $A$ . (2)

- (b) Find the value of the constant  $p$ . (3)

- (c) Find the acute angle between  $l_1$  and  $l_2$ , giving your answer in degrees to 2 decimal places.
- (3)**

The point  $B$  lies on  $l_2$  where  $\mu = 1$

- (d) Find the shortest distance from the point  $B$  to the line  $l_1$ , giving your answer to 3 significant figures.
- (3)**



Leave  
blank

Question 4 continued

Lined area for writing the answer to Question 4.



Leave  
blank



5. A curve  $C$  has parametric equations

(a) Find the value of  $\frac{dy}{dx}$  at the point on  $C$  where  $t = 2$ , giving your answer as a fraction in its simplest form.

(3)

(b) Show that the cartesian equation of the curve  $C$  can be written in the form

where  $a$  and  $b$  are integers to be determined.

(3)

[illegible]






Leave  
blank

[illegible]

**Q5**

**(Total 6 marks)**





### Figure 2

Figure 2 shows a sketch of the curve with equation  $y = \sqrt{(3-x)(x+1)}$ ,  $0 \leq x \leq 3$

The finite region  $R$ , shown shaded in Figure 2, is bounded by the curve, the  $x$ -axis, and the  $y$ -axis.

- (a) Use the substitution  $x = 1 + 2 \sin \theta$  to show that

$$\int_0^3 \sqrt{(3-x)(x+1)} \, dx = k \int_{-\frac{\pi}{6}}^{\frac{\pi}{2}} \cos^2 \theta \, d\theta$$

where  $k$  is a constant to be determined.

(5)

- (b) Hence find, by integration, the exact area of  $R$ .

(3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Leave  
blank

## This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire width, providing a template for writing or drawing. The margins are consistent on all sides.



**Q6**

**(Total 8 marks)**



7. (a) Express  $\frac{2}{P(P-2)}$  in partial fractions.

A team of biologists is studying a population of a particular species of animal.

The population is modelled by the differential equation

$$\frac{dP}{dt} = \frac{1}{2}P(P - 2)\cos 2t, \quad t \geq 0$$

where  $P$  is the population in thousands, and  $t$  is the time measured in years since the start of the study.

Given that  $P = 3$  when  $t = 0$ ,

(b) solve this differential equation to show that

$$P = \frac{6}{3 - e^{\frac{1}{2}\sin 2t}}$$

(7)

(c) find the time taken for the population to reach 4000 for the first time. Give your answer in years to 3 significant figures.

(3)











8.




Figure 3

Figure 3 shows a sketch of part of the curve  $C$  with equation

$$y = 3^x$$

The point  $P$  lies on  $C$  and has coordinates  $(2, 9)$ .

The line  $l$  is a tangent to  $C$  at  $P$ . The line  $l$  cuts the  $x$ -axis at the point  $Q$ .

(a) Find the exact value of the  $x$  coordinate of  $Q$ .

(4)

The finite region  $R$ , shown shaded in Figure 3, is bounded by the curve  $C$ , the  $x$ -axis, the  $y$ -axis and the line  $l$ . This region  $R$  is rotated through  $360^\circ$  about the  $x$ -axis.

(b) Use integration to find the exact value of the volume of the solid generated.

Give your answer in the form  $\frac{p}{q}$  where  $p$  and  $q$  are exact constants.

[You may assume the formula  $V = \frac{1}{3}\pi r^2 h$  for the volume of a cone.]

(6)

---

---

---

---

---

---

---

---

---

---





Leave  
blank

[illegible]

Leave  
blank

[illegible]

Leave  
blank

**(Total 10 marks)**

**TOTAL FOR PAPER: 75 MARKS**

**END**

**Q8**

