

Write your name here

Surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Mechanics M1

Advanced/Advanced Subsidiary

Tuesday 20 January 2015 – Morning

Time: 1 hour 30 minutes

Paper Reference

WME01/01**You must have:**

Mathematical Formulae and Statistical Tables (Blue)

Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P45061A

©2015 Pearson Education Ltd.

1/5/5/5/



PEARSON

1. A railway truck A of mass m and a second railway truck B of mass $4m$ are moving in opposite directions on a smooth straight horizontal track when they collide directly. Immediately before the collision the speed of truck A is $3u$ and the speed of truck B is $2u$. In the collision the trucks join together. Modelling the trucks as particles, find


(c) the magnitude of the impulse exerted by A on B in the collision. (3)

Q1

(Total 7 marks)



Leave
blank



A diagram showing a rectangular block labeled "50 kg" resting on an inclined plane. The plane makes an angle α with the horizontal. A vertical line is drawn from the top of the block, perpendicular to the inclined plane.

Figure 1

A block of mass 50 kg lies on a rough plane which is inclined to the horizontal at an angle α , where $\tan \alpha = \frac{7}{24}$. The block is held at rest by a vertical rope, as shown in

Figure 1, and is on the point of sliding down the plane. The block is modelled as a particle and the rope is modelled as a light inextensible string. Given that the friction force acting on the block has magnitude 65.8 N, find

- (a) the tension in the rope, (4)
- (b) the coefficient of friction between the block and the plane. (4)



Leave
blank

Question 2 continued



Leave
blank

Leave
blank

This image shows a full page of blank, lined paper. It features approximately 28 horizontal grey lines spaced evenly apart, typical of notebook paper. The lines extend across the entire width of the page, leaving small margins at the top and bottom. There are no vertical lines, text, or other markings present.

(Total 8 marks)

Q2







Leave
blank

Question 3 continued

Lined area for writing the answer to Question 3.



Leave
blank

[illegible]

(Total 7 marks)

Q3



4. The points P and Q are at the same height h metres above horizontal ground. A small stone is dropped from rest from P . Half a second later a second small stone is thrown vertically downwards from Q with speed 7.35 m s^{-1} . Given that the stones hit the ground at the same time, find the value of h .

(7)





Leave
blank

Question 4 continued

Lined area for writing the answer to Question 4.



Leave
blank

[illegible]

(Total 7 marks)

Q4



Leave
blank

5.




Figure 2

A particle P of mass 2 kg is pushed up a line of greatest slope of a rough plane by a horizontal force of magnitude X newtons, as shown in Figure 2. The force acts in the vertical plane which contains P and a line of greatest slope of the plane. The plane is

inclined to the horizontal at an angle α , where $\tan \alpha = \frac{3}{4}$

The coefficient of friction between P and the plane is 0.5

Given that the acceleration of P is 1.45 m s^{-2} , find the value of X .

(10)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Leave
blank

6. A uniform rod AC , of weight W and length $3l$, rests horizontally on two supports, one at A and one at B , where $AB = 2l$. A particle of weight $2W$ is placed on the rod at a distance x from A . The rod remains horizontal and in equilibrium.

(5)

(5)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Leave
blank

Question 6 continued

Lined area for writing the answer to Question 6.



Leave
blank

7. A train travels along a straight horizontal track between two stations A and B . The train starts from rest at A and moves with constant acceleration until it reaches its maximum speed of 108 km h^{-1} . The train then travels at this speed before it moves with constant deceleration coming to rest at B . The journey from A to B takes 8 minutes.

- (2)

- (2)

(6)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Leave
blank

(Total 10 marks)



Leave
blank

8.




Figure 3

A particle A of mass $3m$ is held at rest on a rough horizontal table. The particle is attached to one end of a light inextensible string. The string passes over a small smooth pulley P which is fixed at the edge of the table. The other end of the string is attached to a particle B of mass $2m$, which hangs freely, vertically below P . The system is released from rest, with the string taut, when A is 1.3 m from P and B is 1 m above the horizontal floor, as shown in Figure 3.

Given that B hits the floor 2 s after release and does not rebound,

- (a) find the acceleration of A during the first two seconds, (2)
- (b) find the coefficient of friction between A and the table, (8)
- (c) determine whether A reaches the pulley. (6)



Leave
blank

Question 8 continued

Lined area for writing the answer to Question 8.



Leave
blank

Question 8 continued

Handwriting practice area with horizontal lines.





Leave
blank

(Total 16 marks)

Q8

TOTAL FOR PAPER: 75 MARKS

END

