

Write your name here

Surname	Other names
---------	-------------

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Mechanics M1

Advanced/Advanced Subsidiary

Tuesday 20 January 2015 – Morning
Time: 1 hour 30 minutes

Paper Reference
WME01/01

You must have:
Mathematical Formulae and Statistical Tables (Blue)

Total Marks

--

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P45061A

©2015 Pearson Education Ltd.

1/5/15/5/



PEARSON

Leave
blank

Question 3 continued

Lined area for writing the answer to Question 3.

Q3

(Total 7 marks)



P 4 5 0 6 1 A 0 1 1 3 2

Question 4 continued

Lined writing area for question 4 continued.



5.




Figure 2

A particle P of mass 2 kg is pushed up a line of greatest slope of a rough plane by a horizontal force of magnitude X newtons, as shown in Figure 2. The force acts in the vertical plane which contains P and a line of greatest slope of the plane. The plane is

inclined to the horizontal at an angle α , where $\tan \alpha = \frac{3}{4}$

The coefficient of friction between P and the plane is 0.5

Given that the acceleration of P is 1.45 m s^{-2} , find the value of X .

(10)



Leave blank

6. A uniform rod AC , of weight W and length $3l$, rests horizontally on two supports, one at A and one at B , where $AB = 2l$. A particle of weight $2W$ is placed on the rod at a distance x from A . The rod remains horizontal and in equilibrium.

(a) Find the greatest possible value of x .

(5)

The magnitude of the reaction of the support at A is R . Due to a weakness in the support at A , the greatest possible value of R is $2W$,

(b) find the least possible value of x .

(5)



Leave
blank

Question 6 continued

Lined writing area for the answer to Question 6.



P 4 5 0 6 1 A 0 2 1 3 2

7. A train travels along a straight horizontal track between two stations *A* and *B*. The train starts from rest at *A* and moves with constant acceleration until it reaches its maximum speed of 108 km h^{-1} . The train then travels at this speed before it moves with constant deceleration coming to rest at *B*. The journey from *A* to *B* takes 8 minutes.

(a) Change 108 km h^{-1} into m s^{-1} . (2)

(b) Sketch a speed-time graph for the motion of the train between the two stations *A* and *B*. (2)

Given that the distance between the two stations is 12 km and that the time spent decelerating is three times the time spent accelerating,

(c) find the acceleration, in m s^{-2} , of the train. (6)



Leave blank

8.




Figure 3

A particle A of mass $3m$ is held at rest on a rough horizontal table. The particle is attached to one end of a light inextensible string. The string passes over a small smooth pulley P which is fixed at the edge of the table. The other end of the string is attached to a particle B of mass $2m$, which hangs freely, vertically below P . The system is released from rest, with the string taut, when A is 1.3 m from P and B is 1 m above the horizontal floor, as shown in Figure 3.

Given that B hits the floor 2 s after release and does not rebound,

(a) find the acceleration of A during the first two seconds, (2)

(b) find the coefficient of friction between A and the table, (8)

(c) determine whether A reaches the pulley. (6)



Leave blank

Question 8 continued

Lined writing area for the answer to Question 8 continued.



Leave
blank

Question 8 continued

Q8

(Total 16 marks)

TOTAL FOR PAPER: 75 MARKS

END

