Examiner's use only

Team Leader's use only

Duestion

1

2

3

4

5

6

7

Leave

6677

Past Paper

This resource was created and owned by Pearson Edexcel

Centre No.	·					Pape	r Refer	ence			Surname		Initial(s)
Candidate No.				6	6	7	7	/	0	1	Signature		

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Friday 12 January 2007 - Morning

Time: 1 hour 30 minutes

Materials required for examination

Mathematical Formulae (Green)

Items included with question papers

Nil

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature.

Check that you have the correct question paper.

You must write your answer for each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 16 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

©2007 Edexcel Limited.

N23560A W850/R6677/57570 3/3/3/3/3/14,400



Turn over

Total

edexcel

advancing learning, changing lives




blank

₽ast Paper

This resource was created and owned by Pearson Edexcel

1. Figure 1



A particle of weight 24 N is held in equilibrium by two light inextensible strings. One string is horizontal. The other string is inclined at an angle of 30° to the horizontal, as shown in Figure 1. The tension in the horizontal string is Q newtons and the tension in the other string is P newtons. Find

(a) the value of P,

(3)

(b) the value of Q.

(3)

			•		
		*			-

The second secon			AV		
			·		
	The second secon	777 - 1178-2		To the first the state of the s	

Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

Mathematics M1

•		•	•	•
	6	6	7	7

	Leave blank
Question 1 continued	
· · · · · · · · · · · · · · · · · · ·	
· · · · · · · · · · · · · · · · · · ·	
	Q1
	\bigcap
(Total 6 marks)	[] [




This resource was created and owned by Pearson Edexcel

6677

Leave blank

2.

Figure 2



A uniform plank AB has weight 120 N and length 3 m. The plank rests horizontally in equilibrium on two smooth supports C and D, where AC = 1 m and CD = x m, as shown in Figure 2. The reaction of the support on the plank at D has magnitude 80 N. Modelling the plank as a rod,

(a) show that x = 0.75

(3)

A rock is now placed at B and the plank is on the point of tilting about D. Modelling the rock as a particle, find

(b) the weight of the rock,

(4)

(c) the magnitude of the reaction of the support on the plank at D.

(2)

(d) State how you have used the model of the rock as a particle.

(1)

.1



www.mystudybro.com

dybro.com Mathematics M1

	,	•		
ast Paper	This resource was created	d and ow	ned by Pearson	Edexce

AL MANAGEME	



Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

6	6	7	7

	Leave blank
3. A particle P of mass 2 kg is moving under the action of a constant force \mathbf{F} newtons. When $t = 0$, P has velocity $(3\mathbf{i} + 2\mathbf{j})$ m s ⁻¹ and at time $t = 4$ s, P has velocity $(15\mathbf{i} - 4\mathbf{j})$ m s ⁻¹ . Find	
(a) the acceleration of P in terms of \mathbf{i} and \mathbf{j} ,	
(2)	
(b) the magnitude of F ,	
(4)	
(c) the velocity of P at time $t = 6$ s.	
(3)	
	İ

Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

Mathematics M1

_		•	•	•
	_	_	_	_

Question 3 continued	
Anestron 2 continued	
	and the best of the
	,, ————
en e	



Mathematics M1

Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

		Leave
4.	A particle P of mass 0.3 kg is moving with speed u m s ⁻¹ in a straight line on a smooth horizontal table. The particle P collides directly with a particle Q of mass 0.6 kg, which is at rest on the table. Immediately after the particles collide, P has speed 2 m s ⁻¹ and Q has speed 5 m s ⁻¹ . The direction of motion of P is reversed by the collision. Find	
	(a) the value of u ,	
	(4)	
	(b) the magnitude of the impulse exerted by P on Q .	
	(2)	
	Immediately after the collision, a constant force of magnitude R newtons is applied to Q in the direction directly opposite to the direction of motion of Q . As a result Q is brought to rest in $1.5 \mathrm{s}$.	
	(c) Find the value of R .	
	(4)	
	-	
	en de la comp etition de la competition della c	
rene s um e ener _e so		

Mathematics M1

Past	Paper

Γ

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

Question 4 continued	
	The state of the s
· · · · · · · · · · · · · · · · · · ·	-
	erman er euro
	Not desirable.



Turn over

Mathematics M1

Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

(a) the greatest height above A reached by the ball,	د
	(3)
(b) the speed of the ball as it reaches the ground,	(3)
(c) the time between the instant when the ball is projected from A and the	instant when
the ball reaches the ground.	
	(4)
	· made at a 1 after 10 to 10 - 10 at the 20 m at 10 to 10 at the attention of the 10 minutes.
	en e
MALE TO THE PROPERTY OF THE PROPERTY AND CONTRACT OF THE PROPERTY OF THE PROPE	

Mathematics M1

Pact	Paner

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

667	7
ave ank	

	Lear blar
Question 5 continued	
	-
· · · · · · · · · · · · · · · · · · ·	
	Q
(Total 10 marks)	



www.mystudybro.com

Past Paper


This resource was created and owned by Pearson Edexcel

Leave

blank

6.

Figure 3



A box of mass 30 kg is being pulled along rough horizontal ground at a constant speed using a rope. The rope makes an angle of 20° with the ground, as shown in Figure 3. The coefficient of friction between the box and the ground is 0.4. The box is modelled as a particle and the rope as a light, inextensible string. The tension in the rope is P newtons.

(a) Find the value of P.

(8)

The tension in the rope is now increased to 150 N.

(b) Find the acceleration of the box.

(6)

·

Winter 2007	www.mystudybro.com	Mathematics M1
Past Paper	This resource was created and owned by Pearson Edexcel	6677

Add data hash Affir Allin Ayring a suprimper a transfer on property property and a suprimper a post	
And the state of t	
The decision of the Control of the C	



Turn over

This resource was created and owned by Pearson Edexcel

Leave blank

7.

Figure 4




Figure 4 shows two particles P and Q, of mass 3 kg and 2 kg respectively, connected by a light inextensible string. Initially P is held at rest on a fixed smooth plane inclined at 30° to the horizontal. The string passes over a small smooth light pulley A fixed at the top of the plane. The part of the string from P to A is parallel to a line of greatest slope of the plane. The particle Q hangs freely below A. The system is released from rest with the string taut.

(a) Write down an equation of motion for P and an equation of motion for Q.

(4)

(b) Hence show that the acceleration of Q is 0.98 m s⁻².

(2)

(c) Find the tension in the string.

(2)

(d) State where in your calculations you have used the information that the string is inextensible.

(1)

On release, Q is at a height of 0.8 m above the ground. When Q reaches the ground, it is brought to rest immediately by the impact with the ground and does not rebound. The initial distance of P from A is such that in the subsequent motion P does not reach A. Find

(e) the speed of Q as it reaches the ground,

(2)

(f) the time between the instant when Q reaches the ground and the instant when the string becomes taut again.

(5)

Past Paper

www.mystudybro.comThis resource was created and owned by Pearson Edexcel

Mathematics M1

Overtion 7 continued	bl
Question 7 continued	



Past Paper

www.mystudybro.com

Mathematics M1

This resource was	created and	owned by	/ Pearson	Edexcel

Question 7 continued
zuestion / continued

- · · · · · · · · · · · · · · · · · · ·
(Total 16 marks)