





Leave  
blank

Question 1 continued

Lined area for student response, consisting of multiple horizontal lines for writing.

(Total 6 marks)

Q1



Leave  
blank

2. A firework rocket starts from rest at ground level and moves vertically. In the first 3 s of its motion, the rocket rises 27 m. The rocket is modelled as a particle moving with constant acceleration  $a$  m s<sup>-2</sup>. Find

(a) the value of  $a$ , (2)

(b) the speed of the rocket 3 s after it has left the ground. (2)

After 3 s, the rocket burns out. The motion of the rocket is now modelled as that of a particle moving freely under gravity.

(c) Find the height of the rocket above the ground 5 s after it has left the ground. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave  
blank

**Question 2 continued**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**(Total 8 marks)**

Q2



M 2 6 3 3 1 A 0 5 2 8

Leave  
blank

3. A car moves along a horizontal straight road, passing two points  $A$  and  $B$ . At  $A$  the speed of the car is  $15 \text{ m s}^{-1}$ . When the driver passes  $A$ , he sees a warning sign  $W$  ahead of him,  $120 \text{ m}$  away. He immediately applies the brakes and the car decelerates with uniform deceleration, reaching  $W$  with speed  $5 \text{ m s}^{-1}$ . At  $W$ , the driver sees that the road is clear. He then immediately accelerates the car with uniform acceleration for  $16 \text{ s}$  to reach a speed of  $V \text{ m s}^{-1}$  ( $V > 15$ ). He then maintains the car at a constant speed of  $V \text{ m s}^{-1}$ . Moving at this constant speed, the car passes  $B$  after a further  $22 \text{ s}$ .

(a) Sketch, in the space below, a speed-time graph to illustrate the motion of the car as it moves from  $A$  to  $B$ .

(3)

(b) Find the time taken for the car to move from  $A$  to  $B$ .

(3)

The distance from  $A$  to  $B$  is  $1 \text{ km}$ .

(c) Find the value of  $V$ .

(5)












Leave  
blank

4.

**Figure 1**

A particle  $P$  of mass  $6\text{ kg}$  lies on the surface of a smooth plane. The plane is inclined at an angle of  $30^\circ$  to the horizontal. The particle is held in equilibrium by a force of magnitude  $49\text{ N}$ , acting at an angle  $\theta$  to the plane, as shown in Figure 1. The force acts in a vertical plane through a line of greatest slope of the plane.

(a) Show that  $\cos \theta = \frac{3}{5}$ . **(3)**

(b) Find the normal reaction between  $P$  and the plane. **(4)**

The direction of the force of magnitude  $49\text{ N}$  is now changed. It is now applied horizontally to  $P$  so that  $P$  moves up the plane. The force again acts in a vertical plane through a line of greatest slope of the plane.

(c) Find the initial acceleration of  $P$ . **(4)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







5.




Figure 2

A beam  $AB$  has mass 12 kg and length 5 m. It is held in equilibrium in a horizontal position by two vertical ropes attached to the beam. One rope is attached to  $A$ , the other to the point  $C$  on the beam, where  $BC = 1$  m, as shown in Figure 2. The beam is modelled as a uniform rod, and the ropes as light strings.

- (a) Find
- (i) the tension in the rope at  $C$ ,
  - (ii) the tension in the rope at  $A$ .
- (5)**

A small load of mass 16 kg is attached to the beam at a point which is  $y$  metres from  $A$ . The load is modelled as a particle. Given that the beam remains in equilibrium in a horizontal position,

- (b) find, in terms of  $y$ , an expression for the tension in the rope at  $C$ .
- (3)**

The rope at  $C$  will break if its tension exceeds 98 N. The rope at  $A$  cannot break.

- (c) Find the range of possible positions on the beam where the load can be attached without the rope at  $C$  breaking.
- (3)**

---

---

---

---

---

---

---

---

---

---

---

---











Leave blank

6. [In this question, the unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are due east and due north respectively.]

A particle  $P$  is moving with constant velocity  $(-5\mathbf{i} + 8\mathbf{j}) \text{ m s}^{-1}$ . Find

(a) the speed of  $P$ , (2)

(b) the direction of motion of  $P$ , giving your answer as a bearing. (3)

At time  $t = 0$ ,  $P$  is at the point  $A$  with position vector  $(7\mathbf{i} - 10\mathbf{j}) \text{ m}$  relative to a fixed origin  $O$ . When  $t = 3 \text{ s}$ , the velocity of  $P$  changes and it moves with velocity  $(u\mathbf{i} + v\mathbf{j}) \text{ m s}^{-1}$ , where  $u$  and  $v$  are constants. After a further  $4 \text{ s}$ , it passes through  $O$  and continues to move with velocity  $(u\mathbf{i} + v\mathbf{j}) \text{ m s}^{-1}$ .

(c) Find the values of  $u$  and  $v$ . (5)

(d) Find the total time taken for  $P$  to move from  $A$  to a position which is due south of  $A$ . (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave blank

**Question 6 continued**

Handwritten area with horizontal lines for student response.







7.



Figure 3

Two particles  $A$  and  $B$ , of mass  $m$  and  $2m$  respectively, are attached to the ends of a light inextensible string. The particle  $A$  lies on a rough horizontal table. The string passes over a small smooth pulley  $P$  fixed on the edge of the table. The particle  $B$  hangs freely below the pulley, as shown in Figure 3. The coefficient of friction between  $A$  and the table is  $\mu$ . The particles are released from rest with the string taut. Immediately after release, the magnitude of the acceleration of  $A$  and  $B$  is  $\frac{4}{9}g$ . By writing down separate equations of motion for  $A$  and  $B$ ,

(a) find the tension in the string immediately after the particles begin to move, (3)

(b) show that  $\mu = \frac{2}{3}$ . (5)

When  $B$  has fallen a distance  $h$ , it hits the ground and does not rebound. Particle  $A$  is then a distance  $\frac{1}{3}h$  from  $P$ .

(c) Find the speed of  $A$  as it reaches  $P$ . (6)

(d) State how you have used the information that the string is light. (1)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave blank

### Question 7 continued

*(This section contains horizontal lines for writing the answer to Question 7.)*



M 2 6 3 3 1 A 0 2 3 2 8





Leave blank

Question 7 continued

Blank area for writing the answer to Question 7, consisting of 30 horizontal lines.

(Total 15 marks)

Q7

TOTAL FOR PAPER: 75 MARKS

END



**BLANK PAGE**



**BLANK PAGE**



M 2 6 3 3 1 A 0 2 7 2 8

**BLANK PAGE**

