

Mark Scheme (Results)

June 2011

GCE Mechanics M1 (6677) Paper 1



Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at <u>www.edexcel.com</u>.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link: http://www.edexcel.com/Aboutus/contact-us/

June 2011 Publications Code UA028437 All the material in this publication is copyright © Edexcel Ltd 2011



EDEXCEL GCE MATHEMATICS

General Instructions for Marking

- 1. The total number of marks for the paper is 75.
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
 - M marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - B marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
- 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod benefit of doubt
- ft follow through
- the symbol will be used for correct ft
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- dep dependent
- indep independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- The second mark is dependent on gaining the first mark



June 2011 Mechanics M1 6677 Mark Scheme

		
Question Number	Scheme	Marks
1. (a)	$0^{2} = u^{2} - 2x9.8x40$ $u = 28 \text{ m s}^{-1} ** \text{ GIVEN ANSWER}$	M1 A1 A1 (3)
(b)	$33.6 = 28t - \frac{1}{2}9.8t^{2}$ $4.9t^{2} - 28t + 33.6 = 0$ $t = \frac{28 \pm \sqrt{28^{2} - 4x4.9x33.6}}{9.8}$ $= 4 \text{ s or } (1.7 \text{ s or } 1.71 \text{ s})$	M1 A1 M1 A1 A1 (5) 8
2. (a)	$\frac{3}{v} \stackrel{2}{\longrightarrow} CLM: \ 3x3 - 2x2 = 3v + 2(v+1)$ $\xrightarrow{v} \stackrel{*}{\longrightarrow} v_{P} = 0.6 \text{ m s}^{-1}; \ v_{Q} = 1.6 \text{ m s}^{-1}$	M1 A1 M1A1 (A1 ft) (5)
(b)	3(v-3) OR $2(v+12)= 7.2 Ns = 7.2 Ns$	M1 A1 ft A1 (3) 8



Question Number	Scheme	Marks
3. <u>OR</u>	$4\cos \alpha + F = W \sin \alpha$ $R = 4\sin \alpha + W \cos \alpha$ $F = 0.5R$ $\cos \alpha = 0.8 \text{ or } \sin \alpha = 0.6$ $R = 20N ** \text{ GIVEN ANSWER}$ $W = 22N$ $R \sin \alpha = 4 + F \cos \alpha$ $R \cos \alpha + F \sin \alpha = W$ $F = 0.5R$ $\cos \alpha = 0.8 \text{ or } \sin \alpha = 0.6$ $R = 20N ** \text{ GIVEN ANSWER}$ $W = 22N$	M1 A1 M1 A1 B1 B1 M1 A1 A1 (9) M1 A1 B1 B1 B1 B1 M1 A1 A1 (9) 9
4. (a)	$ \begin{array}{c} 5 \\ \nu \\ 0 \\ 4 \\ 64 \\ 84 \end{array} $	B1 shape B1 figs (2)
(b)	$(\frac{1}{2}x4x5) + 60 \times 5$ = 310	M1 A1 A1 (3)
(c)	$\frac{(5+V)}{2} \ge 20 = (400-310)$ $V = 4$	M1 A2 ft DM1 A1 (5)
(d)	$\frac{5-4}{20} = 0.05 \text{ ms}^{-2}$	M1 A1 (2) 12

GCE Mechanics M1 (6677) June 2011



Question Number	Scheme	Marks
5. (a)	$P \xrightarrow{2 \text{ m} 2 \text{ m} 2 \text{ m} 2 \text{ m} Q 2 \text{ m}}_{X 40g 20g X Mg} R$	
(i)	EITHER M(<i>R</i>), $8X + 2X = 40g \ge 6 + 20g \ge 4$ solving for <i>X</i> , $X = 32g = 314$ or 310 N	M1 A2 M1 A1
(ii)	(1) $X + X = 40g + 20g + Mg$ (or another moments equation)	M1 A2
(i)	Solving for $M, M = 4$ OR $M(P), 6X = 40g \ge 2 + 20g \ge 4 + Mg \ge 8$ Solving for $X, X = 32g = 314$ or 310 N (\uparrow) $X + X = 40g + 20g + Mg$ (or another moments)	M1 A1 M1 A2 M1 A1
(ii)	equation) solving for $M, M = 4$	M1 A2 M1 A1 (10
(b)	Masses concentrated at a point or weights act at a point	B1 (1
6. (a)	$R = 0.3g \cos \alpha$ = 0.24g = 2.35 (3sf)=2.4 (2sf)	M1 A1 (2
(b)	$mg - T = 1.4m$ $T - 0.3g \sin \alpha - F = 0.3 \times 1.4$ $F = 0.5R$ Eliminating R and T $m = 0.4$	M1 A1 M1 A2 M1 DM 1 A1
(c)	$v = 1.4 \times 0.5$ -0.3g sin α - F = 0.3a a = -9.8 0 = 0.7 - 9.8t t = 0.071 s or 0.0714 s (1/14 A0)	(8) B1 M1 A1 A1 M1 A1 (6) 10



		advancing learning, changing li
Question Number	Scheme	Marks
7. (a)	$\tan\theta = \frac{3}{4}$; bearing is 37° (nearest degree)	M1; A1
(b) (i) (ii) (iii)	p = (i + j) + t(2i - 3j) q = (-2j) + t(3i + 4j) PQ = q - p = (-i - 3j) + t(i + 7j)	(2) M1 A1 A1 M1 A1
(c) (i)	-1+t = 0 t = 1 or 3pm -1+t = -(-3+7t)	(5) M1 A1 M1
(ii)	$t = \frac{1}{2}$ or 2.30 pm	M1 A1 (4) 11

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467 Fax 01623 450481 Email <u>publication.orders@edexcel.com</u> Order Code UA028437 June 2011

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE







