

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	7	/	0	1	Signature	

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Wednesday 16 May 2012 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2012 Pearson Education Ltd

Printer's Log. No. _____

Printer's Log. No.
P40689A

W850/R6677/57570 6/5/3



Turn over

PEARSON

1. Two particles A and B , of mass $5m$ kg and $2m$ kg respectively, are moving in opposite directions along the same straight horizontal line. The particles collide directly. Immediately before the collision, the speeds of A and B are 3 m s^{-1} and 4 m s^{-1} respectively. The direction of motion of A is unchanged by the collision. Immediately after the collision, the speed of A is 0.8 m s^{-1} .

(3)

(b) Find the value of m .

(3)



Leave
blank

Question 1 continued

Q1

(Total 6 marks)






Figure 1

A non-uniform rod AB has length 3 m and mass 4.5 kg. The rod rests in equilibrium, in a horizontal position, on two smooth supports at P and at Q , where $AP = 0.8$ m and $QB = 0.6$ m, as shown in Figure 1. The centre of mass of the rod is at G . Given that the magnitude of the reaction of the support at P on the rod is twice the magnitude of the reaction of the support at Q on the rod, find

- (a) the magnitude of the reaction of the support at Q on the rod,

(3)

- (b) the distance AG .

(4)



Leave
blank

Question 2 continued

Q2

(Total 7 marks)



Leave
blank

A diagram showing a black rectangular block on a grey inclined plane. The plane is angled upwards from left to right, with an angle of 30° indicated at the bottom left. A horizontal arrow points to the right, labeled "20 N", representing a force applied to the block.

Figure 2

Find

- (a) the magnitude of the normal reaction of the plane on the box, (4)
- (b) the coefficient of friction between the box and the plane. (5)



Leave
blank

Question 3 continued

Lined area for writing answers.



Leave
blank

(Total 9 marks)

Q3



4. A car is moving on a straight horizontal road. At time $t = 0$, the car is moving with speed 20 m s^{-1} and is at the point A . The car maintains the speed of 20 m s^{-1} for 25 s. The car then moves with constant deceleration 0.4 m s^{-2} , reducing its speed from 20 m s^{-1} to 8 m s^{-1} . The car then moves with constant speed 8 m s^{-1} for 60 s. The car then moves with constant acceleration until it is moving with speed 20 m s^{-1} at the point B .

(a) Sketch a speed-time graph to represent the motion of the car from A to B .

(3)

(b) Find the time for which the car is decelerating.

(2)

Given that the distance from A to B is 1960 m,

(c) find the time taken for the car to move from A to B .

(8)



Leave
blank

Question 4 continued

Lined area for writing answers.



Leave
blank

(Total 13 marks)

Q4



5. A particle P is projected vertically upwards from a point A with speed $u \text{ m s}^{-1}$. The point A is 17.5 m above horizontal ground. The particle P moves freely under gravity until it reaches the ground with speed 28 m s^{-1} .

- (3)

(5)

(4)

Leave
blank

Question 5 continued

Lined area for writing the answer to Question 5.



Leave
blank

Leave
blank

(Total 12 marks)

Q5



Leave
blank

- A ship S is moving with constant velocity $(-12\mathbf{i} + 7.5\mathbf{j}) \text{ km h}^{-1}$.

- At time t hours after noon, the position vector of S is \mathbf{s} km. When $t = 0$, $\mathbf{s} = 40\mathbf{i} - 6\mathbf{j}$.

- A fixed beacon B is at the point with position vector $(7\mathbf{i} + 12.5\mathbf{j})$ km.

- (d) Find the distance of S from B when S is due north of B . (4)



Leave
blank

Question 6 continued

Lined area for writing answers.



Leave
blank



Leave
blank

(Total 13 marks)

Q6





Figure 3

At time $t = 0$, a constant force \mathbf{F} of magnitude 4 N is applied to Q in the direction PQ , as shown in Figure 3. The system moves under the action of this force until $t = 6$ s. During the motion, the resistance to the motion of P has constant magnitude 1 N and the resistance to the motion of Q has constant magnitude 2 N.

Find

- (a) the acceleration of the particles as the system moves under the action of \mathbf{F} ,

- (b) the speed of the particles at $t = 6$ s,

- (c) the tension in the rod as the system moves under the action of \mathbf{F} . (3)

At $t = 6$ s, \mathbf{F} is removed and the system decelerates to rest. The resistances to motion are unchanged. Find

- (d) the distance moved by P as the system decelerates, (4)

- (e) the thrust in the rod as the system decelerates. (3)





Leave
blank



Leave
blank

Leave
blank

Question 7 continued

Handwriting practice area with 30 horizontal lines.



Leave
blank

(Total 15 marks)

Q7

TOTAL FOR PAPER: 75 MARKS

END

