

Leave blank

1. Two particles A and B , of mass $5m$ kg and $2m$ kg respectively, are moving in opposite directions along the same straight horizontal line. The particles collide directly. Immediately before the collision, the speeds of A and B are 3 m s^{-1} and 4 m s^{-1} respectively. The direction of motion of A is unchanged by the collision. Immediately after the collision, the speed of A is 0.8 m s^{-1} .

(a) Find the speed of B immediately after the collision.

(3)

In the collision, the magnitude of the impulse exerted on A by B is 3.3 N s .

(b) Find the value of m .

(3)



Leave blank

Question 1 continued

Lined area for writing the answer to Question 1.

(Total 6 marks)

Q1



Leave blank

2.




Figure 1

A non-uniform rod AB has length 3 m and mass 4.5 kg. The rod rests in equilibrium, in a horizontal position, on two smooth supports at P and at Q , where $AP = 0.8$ m and $QB = 0.6$ m, as shown in Figure 1. The centre of mass of the rod is at G . Given that the magnitude of the reaction of the support at P on the rod is twice the magnitude of the reaction of the support at Q on the rod, find

(a) the magnitude of the reaction of the support at Q on the rod, (3)

(b) the distance AG . (4)



3.




Figure 2

A box of mass 5 kg lies on a rough plane inclined at 30° to the horizontal. The box is held in equilibrium by a horizontal force of magnitude 20 N , as shown in Figure 2. The force acts in a vertical plane containing a line of greatest slope of the inclined plane. The box is in equilibrium and on the point of moving down the plane. The box is modelled as a particle.

Find

(a) the magnitude of the normal reaction of the plane on the box, (4)

(b) the coefficient of friction between the box and the plane. (5)



Leave blank

Question 3 continued

[Lined area for writing answers]



Leave blank

Question 3 continued

A series of horizontal lines for writing the answer to Question 3.



Leave
blank

4. A car is moving on a straight horizontal road. At time $t = 0$, the car is moving with speed 20 m s^{-1} and is at the point A . The car maintains the speed of 20 m s^{-1} for 25 s. The car then moves with constant deceleration 0.4 m s^{-2} , reducing its speed from 20 m s^{-1} to 8 m s^{-1} . The car then moves with constant speed 8 m s^{-1} for 60 s. The car then moves with constant acceleration until it is moving with speed 20 m s^{-1} at the point B .

(a) Sketch a speed-time graph to represent the motion of the car from A to B .

(3)

(b) Find the time for which the car is decelerating.

(2)

Given that the distance from A to B is 1960 m,

(c) find the time taken for the car to move from A to B .

(8)



Leave
blank

Question 5 continued

Blank lined area for writing the answer to Question 5.

Q5

(Total 12 marks)

--	--



Leave blank

6. [In this question \mathbf{i} and \mathbf{j} are horizontal unit vectors due east and due north respectively and position vectors are given with respect to a fixed origin.]

A ship S is moving with constant velocity $(-12\mathbf{i} + 7.5\mathbf{j})$ km h⁻¹.

(a) Find the direction in which S is moving, giving your answer as a bearing. (3)

At time t hours after noon, the position vector of S is \mathbf{s} km. When $t = 0$, $\mathbf{s} = 40\mathbf{i} - 6\mathbf{j}$.

(b) Write down \mathbf{s} in terms of t . (2)

A fixed beacon B is at the point with position vector $(7\mathbf{i} + 12.5\mathbf{j})$ km.

(c) Find the distance of S from B when $t = 3$ (4)

(d) Find the distance of S from B when S is due north of B . (4)



7.



Figure 3

Two particles P and Q , of mass 0.3 kg and 0.5 kg respectively, are joined by a light horizontal rod. The system of the particles and the rod is at rest on a horizontal plane. At time $t = 0$, a constant force F of magnitude 4 N is applied to Q in the direction PQ , as shown in Figure 3. The system moves under the action of this force until $t = 6$ s. During the motion, the resistance to the motion of P has constant magnitude 1 N and the resistance to the motion of Q has constant magnitude 2 N.

Find

- (a) the acceleration of the particles as the system moves under the action of F , (3)
- (b) the speed of the particles at $t = 6$ s, (2)
- (c) the tension in the rod as the system moves under the action of F . (3)

At $t = 6$ s, F is removed and the system decelerates to rest. The resistances to motion are unchanged. Find

- (d) the distance moved by P as the system decelerates, (4)
- (e) the thrust in the rod as the system decelerates. (3)



Leave
blank

Question 7 continued

Blank lined area for writing the answer to Question 7.

Q7

(Total 15 marks)

TOTAL FOR PAPER: 75 MARKS

END

