

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	7	/	0	1	Signature	

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Wednesday 3 June 2015 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$ and give your answer to either two significant figures or three significant figures.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2015 Pearson Education Ltd

Printer's Log. No.

Printer's Log. No.
P44835A

5/1/1/1/1/



Turn over

PEARSON

Leave
blank

Find

- (a) the value of k , (3)
- (b) the magnitude of the impulse exerted on P by Q in the collision. (3)

This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire width, providing a guide for handwriting or typing. The paper itself is a clean, off-white color.

Q1

104

This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire surface, typical of notebook or school paper. There are no margins, text, or other markings present.

(Total 6 marks)



Leave
blank

- (5)



Q2

1

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

(Total 7 marks)



3.




Figure 1

A particle of mass 2 kg is suspended from a horizontal ceiling by two light inextensible strings, PR and QR . The particle hangs at R in equilibrium, with the strings in a vertical plane. The string PR is inclined at 55° to the horizontal and the string QR is inclined at 35° to the horizontal, as shown in Figure 1.

Find

- (i) the tension in the string PR ,
- (ii) the tension in the string QR .

(7)



Q3

(Total 7 marks)



Leave
blank

A diagram showing a person standing on a platform. A vertical line extends upwards from the center of the platform, representing the person's head and torso. The platform is a shaded rectangular area labeled "55 kg".

Figure 2

A lift of mass 200 kg is being lowered into a mineshaft by a vertical cable attached to the top of the lift. A crate of mass 55 kg is on the floor inside the lift, as shown in Figure 2. The lift descends vertically with constant acceleration. There is a constant upwards resistance of magnitude 150 N on the lift. The crate experiences a constant normal reaction of magnitude 473 N from the floor of the lift.

- (a) Find the acceleration of the lift. **(3)**
- (b) Find the magnitude of the force exerted on the lift by the cable. **(4)**


This image shows a single sheet of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Leave
blank

Q4

(Total 7 marks)



5.




Figure 3

A beam AB has length 5 m and mass 25 kg. The beam is suspended in equilibrium in a horizontal position by two vertical ropes. One rope is attached to the beam at A and the other rope is attached to the point C on the beam where $CB = 0.5$ m, as shown in Figure 3. A particle P of mass 60 kg is attached to the beam at B and the beam remains in equilibrium in a horizontal position. The beam is modelled as a uniform rod and the ropes are modelled as light strings.

(a) Find

- (i) the tension in the rope attached to the beam at A ,
- (ii) the tension in the rope attached to the beam at C .

(6)

Particle P is removed and replaced by a particle Q of mass M kg at B . Given that the beam remains in equilibrium in a horizontal position,

(b) find

- (i) the greatest possible value of M ,
- (ii) the greatest possible tension in the rope attached to the beam at C .

(6)



Leave
blank

Question 5 continued

Lined area for writing the answer to Question 5.



Leave
blank

6. A particle P is moving with constant velocity. The position vector of P at time t seconds ($t \geq 0$) is \mathbf{r} metres, relative to a fixed origin O , and is given by

(a) Find the initial position vector of P .

(1)

(b) Find the value of T .

(3)

(c) Find the speed of P .

(4)



Leave
blank

Question 6 continued

Lined area for writing answers.



Leave
blank

Q6

104

(Total 8 marks)



7. A train travels along a straight horizontal track between two stations, A and B . The train starts from rest at A and moves with constant acceleration 0.5 m s^{-2} until it reaches a speed of $V \text{ ms}^{-1}$, ($V < 50$). The train then travels at this constant speed before it moves with constant deceleration 0.25 m s^{-2} until it comes to rest at B .

- (a) Sketch in the space below a speed-time graph for the motion of the train between the two stations A and B .

(2)

The total time for the journey from A to B is 5 minutes.

- (b) Find, in terms of V , the length of time, in seconds, for which the train is

- (i) accelerating,
- (ii) decelerating,
- (iii) moving with constant speed.

(5)

Given that the distance between the two stations A and B is 6.3 km,


- (c) find the value of V .

(6)





Leave
blank



Two particles P and Q have mass 4 kg and 0.5 kg respectively. The particles are attached to the ends of a light inextensible string. Particle P is held at rest on a fixed rough plane, which is inclined to the horizontal at an angle α where $\tan \alpha = \frac{4}{3}$. The coefficient of friction between P and the plane is 0.5. The string lies along the plane and passes over a small smooth light pulley which is fixed at the top of the plane. Particle Q hangs freely at rest vertically below the pulley. The string lies in the vertical plane which contains the pulley and a line of greatest slope of the inclined plane, as shown in Figure 4. Particle P is released from rest with the string taut and slides down the plane.

Given that Q has not hit the pulley, find

- (a) the tension in the string during the motion, (11)
- (b) the magnitude of the resultant force exerted by the string on the pulley. (4)





Leave
blank



Leave
blank

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

(Total 15 marks)

Q8

TOTAL FOR PAPER: 75 MARKS

END

