				-		
This	resource	was	created an	d owned	by Pearson	Edexcel

Vrite your name here Surname	Other	names
Pearson Edexcel nternational Advanced Level	Centre Number	Candidate Number
Chemistry Advanced Subsidiar		
Unit 1: The Core Prir	•	nistry
	nciples of Cher	Daper Reference

Instructions

- Use **black** ink or ball-point pen.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided there may be more space than you need.

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets - use this as a guide as to how much time to spend on each question.
- Questions labelled with an asterisk (*) are ones where the quality of your written communication will be assessed
 you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.



Turn over 🕨

PEARSON



O	n thi	S S	ALL the questions in this section. You should aim to spend no more than 20 minutes ection. For each question, select one answer from A to D and put a cross in the box \boxtimes . hange your mind, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .
1	Wh	nich	of the following species has 50 neutrons?
	\times	A	⁵⁰ ₂₃ V
	\times	B	⁸⁶ ₃₇ Rb ⁻
	×	C	⁸⁹ ₃₉ Y ⁺
	×	D	$^{91}_{40}$ Zr ⁺
			(Total for Question 1 = 1 mark)
2	Wh	nich	of the following statements is correct about all isotopes of an element? They have
	×	A	the same mass number.
	\times	В	the same number of neutrons.
	\times	C	more protons than neutrons.
	×	D	the same electronic configuration.
			(Total for Question 2 = 1 mark)
3			ement rhenium has two naturally-occurring isotopes, ¹⁸⁵ Re and ¹⁸⁷ Re. The e atomic mass of rhenium is 186.2.
	Fro	mt	this information, the percentage abundances of these two isotopes are
	\times	A	12% ¹⁸⁵ Re and 88% ¹⁸⁷ Re
	\mathbf{X}	В	40% ¹⁸⁵ Re and 60% ¹⁸⁷ Re
	\times	C	60% ¹⁸⁵ Re and 40% ¹⁸⁷ Re
	\times	D	88% ¹⁸⁵ Re and 12% ¹⁸⁷ Re
			(Total for Question 3 = 1 mark)
	2		

P 4 2 9 7 6 A 0 2 2 4

4	In whi	ch of the following pairs	does the second elem	ent have a lower 1st ionization
7		than the first element?		
		First element	Second element	
	A 🖂	Si	С	
	🖾 B	Na	Mg	
	🛛 C	Ве	В	
	🖾 D	Ar	Ne	
			'	(Total for Question 4 = 1 mark)
5		de of nitrogen contains s the empirical formula		.0 g of oxygen.
	Δ	NO		
	B	NO ₃		
	🖾 C	N ₂ O ₃		
	⊠ D	N ₂ O ₅		
				(Total for Question 5 = 1 mark)
_				
6		ate the total number of	atoms present in 1.8 g	or water, H_2O .
	DATA • The	e molar mass of H ₂ O is 1	8 g mol⁻¹	
	• The	e Avogadro constant is	$6.0 \times 10^{23} \text{ mol}^{-1}$	
	Α	6.0×10^{22}		
	B	6.0×10^{23}		
	○ C	1.8×10^{23}		
	D 🛛	1.8×10^{24}		
				(Total for Question 6 = 1 mark)



e the mass of hydrated sodium thiosulfate, $Na_2S_2O_3$, $5H_2O$, required to prepare of a 0.100 mol dm ⁻³ solution. • that the molar mass of $Na_2S_2O_3$, $5H_2O$ is 248 g mol ⁻¹] • 124 g • 96 g • 4.8 g • 960 g (Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, C_xH_y , was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. • hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon? CH_x
of a 0.100 mol dm ⁻³ solution. that the molar mass of Na ₂ S ₂ O ₃ ,5H ₂ O is 248 g mol ⁻¹] 124 g .96 g 4.8 g .960 g (Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, C _x H _y , was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : CO ₂ = 44; H ₂ O = 18] f the following is a possible formula of the unknown hydrocarbon?
1.124 g 3.96 g 4.8 g 9960 g (Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, C_xH_y , was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
4.8 g 4.8 g 4.960 g (Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, C_xH_y , was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
44.8 g 4960 g (Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, C_xH_y , was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
(Total for Question 7 = 1 mark) (Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, $C_x H_y$, was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
(Total for Question 7 = 1 mark) sample of an unknown hydrocarbon, $C_x H_y$, was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
sample of an unknown hydrocarbon, $C_x H_y$, was burned completely in excess to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
to form 88.0 g of carbon dioxide and 27.0 g of water. hasses / g mol ⁻¹ : $CO_2 = 44$; $H_2O = 18$] f the following is a possible formula of the unknown hydrocarbon?
f the following is a possible formula of the unknown hydrocarbon?
ΞΗ ₄
-
E_2H_6
E_4H_6
(Total for Question 8 = 1 mark)
gadro constant is equal to the number of
rams of any element which contains 6.0 \times 10 ²³ atoms of that element.
toms contained in one mole of any element in its standard state.
particles (atoms, ions or molecules) required to make one gram of a ubstance.
toms contained in one mole of any monatomic element.
(Total for Question 9 = 1 mark)

10 Nitrogen monoxide, NO, can be made by the catalytic oxidation of ammonia, N	IH ₃ .
10 Nitrogen monoxide, NO, can be made by the catalytic oxidation of ammonia, N	lΗ ₃ .
$4NH_3 + 5O_2 \rightarrow 4NO + 6H_2O$	
In an experiment, 8.5 g of ammonia reacted to form 15.0 g of nitrogen monoxi The percentage yield of nitrogen monoxide in this experiment is	de.
▲ 50%	
■ B 57%	
C 100%	
D 176%	
(Total for Question 10	0 = 1 mark)
 Calculate the mass, in grams, of silver chloride, AgCl, formed when excess silver nitrate solution is added to 55.0 cm³ of a 0.200 mol dm⁻³ solution of potassium chloride. [The molar mass of AgCl = 143.4 g mol⁻¹] 	
$AgNO_{3}(aq) + KCI(aq) \rightarrow AgCI(s) + KNO_{3}(aq)$	
▲ A 1.10 g	
B 1.58 g	
C 7.89 g	
□ 11.0 g	
(Total for Question 11	l = 1 mark)
12 Element X is in Group 3 and element Y is in Group 6 of the Periodic Table.Which of the following is the most likely formula of the compound formed whe and Y react together?	en X
\square A X_2Y_3	
$\blacksquare \mathbf{B} X_{3}Y_{2}$	
\Box C X ₂ Y	
\square D XY ₂	
(Total for Question 12	2 = 1 mark)
	5

ασιΓά	me i aper	r 20	14 www.mystudybro.com This resource was created and owned by Pearson Edexcel	Chemistry Unit 7 WCH0
17	N 4 -	4 - 11 ³		
			ic bonding is best described as the electrostatic attraction between	
	\mathbf{X}		positive ions and delocalized electrons.	
	\mathbf{X}	B	protons and electrons.	
	\mathbf{X}	C		
	X	D	nuclei and shared pairs of electrons.	
			(Total for Question 13 =	- 1 mark)
14	Wł	nich	of the following molecules contains a double bond?	
	X	Α	F ₂	
	\mathbf{X}	В	F ₂ O	
	X	С	C ₂ F ₄	
	\mathbf{X}	D	C ₂ F ₆	
			(Total for Question 14 =	= 1 mark)
	X	A	Breaking covalent bonds requires energy and making ionic bonds require energy.	?S
	X	В	energy. Bond breaking is endothermic whereas bond making is exothermic.	
		c	Bond breaking is exothermic whereas bond making is endothermic.	
		D	Breaking ionic bonds releases energy whereas making covalent bonds	
		U	requires energy.	

Sumn Past Pa		Www.mystudybro.com This resource was created and owned by Pearson Edexcel	Chemistry Unit
16	Consid	der the two equations given below.	
		$2H_2(g) + O_2(g) \rightarrow 2H_2O(I)$ $\Delta H = -572 \text{ kJ mol}^{-1}$	
		$2H_2(g) + O_2(g) \rightarrow 2H_2O(g)$ $\Delta H = -484 \text{ kJ mol}^{-1}$	
1	From 1	this information, calculate the enthalpy change for the following proc	cess
		$H_2O(g) \rightarrow H_2O(I)$	
	🖾 A	–44 kJ mol ⁻¹	
	B	+44 kJ mol ⁻¹	
	🛛 C	–88 kJ mol ⁻¹	
	D 🛛	+88 kJ mol ⁻¹	
		(Total for Question	n 16 = 1 mark)
17	How r	nany structural isomers have the molecular formula C ₆ H ₁₄ ?	
	A 🛛	Four	
	B	Five	
	🛛 C	Six	
	D	Seven	
		(Total for Question	n 17 = 1 mark)
	comb	lition to water, which of the following could be formed during the inc ustion of a hydrocarbon? Carbon, carbon monoxide and hydrogen	complete
	B	Carbon and hydrogen	
		Carbon monoxide and hydrogen	
		Carbon and carbon monoxide	
		(Total for Question	n 18 = 1 mark)



19			thene) is a plastic material made by polymerizing the hydrocarbon ethene. of the following is not true?
	Pu	re p	oly(ethene) is
	\times	Α	solidified ethene.
	\mathbf{X}	В	composed of carbon and hydrogen only.
	\mathbf{X}	С	a long-chain compound.
	\times	D	non-biodegradable.
			(Total for Question 19 = 1 mark)
20			of the following statements correctly describes an environmental problem d by the burning of hydrocarbon fuels?
	\times	Α	The carbon dioxide produced is toxic and kills plants.
	×	В	The smoke produced obscures sunlight and leads to global warming.
	×	С	The water produced results in a damaging increase in rainfall.
	×	D	The carbon dioxide produced traps heat radiated from the Earth and leads to global warming.
			(Total for Question 20 = 1 mark)
_			
			(Total for Question 20 = 1 mark) TOTAL FOR SECTION A = 20 MARKS
_			
-			



SECTION B	
Answer ALL the questions. Write your answers in the spaces provided.	
21 Bromine, Br ₂ , can react with both alkanes and alkenes. The type of reaction that occurs depends on whether the Br—Br bond breaks by homolytic or heterolytic fission.	
(a) (i) Write an equation to show the homolytic fission of the Br—Br bond. Do not include curly arrows or state symbols.	(1)
(ii) Write an equation to show the heterolytic fission of the Br—Br bond. Do not include curly arrows or state symbols.	(1)
(iii) Choosing from the products you have given in (a)(i) and (a)(ii), write the formula of a free radical and an electrophile.	(2)
Free radical	
Electrophile	



according to the equation

(b) The compound hexane, C_6H_{14} , can react with bromine, in the presence of UV light,

 $C_6H_{14} + Br_2 \rightarrow C_6H_{13}Br + HBr$ (i) Give the displayed formulae of the three structural isomers of $C_6 H_{13}$ Br that could be formed in the above reaction. (3) **First isomer** Second isomer **Third isomer** (ii) The bromoalkanes and the hydrogen bromide formed in this reaction are hazardous. The bromoalkanes would be labelled as 'flammable'. Suggest a suitable hazard warning for the hydrogen bromide. (1) 10 P 4 2 9 7 6 A 0 1 0 2 4

$ \begin{array}{c} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$	11 Turn over
Suggest an explanation for this prediction.	(2)
*(ii) On the basis of comparing the relative sizes of the fluorine and chlorine atoms, it might be predicted that the F—F bond energy would be greater than the CI—CI bond energy.	
(i) Write an equation for the reaction between methane and fluorine, assuming they react in a 1:1 mole ratio. State symbols are not required.	(1)
(c) Fluorine, F ₂ , and chlorine, Cl ₂ , react with methane , CH ₄ , by a similar mechanism, although the rates of reaction are very different.	
sum of the molar masses of all products	(2)
Use the expression atom economy = $\frac{\text{molar mass of the desired product}}{\text{sum of the molar masses of all products}} \times 100\%$	
Give your answer to three significant figures.	
(iii) Calculate the percentage atom economy by mass for the formation of $C_6^{}H_{_{13}}^{}Br$	

WCH01 (iii) Draw a dot and cross diagram to show the arrangement of the outermost electrons in a fluorine molecule, F₂. (2) (iv) The actual bond energies are shown below. Bond Bond energy / kJ mol⁻¹ F—F 158 CI-CI 243 By referring to your dot and cross diagram in your answer to (c)(iii), suggest an explanation for the fact that the F—F bond energy is **less** than that of the Cl—Cl bond energy. (2) (v) Suggest why a mixture of methane and chlorine requires exposure to UV light, or heat, before a reaction occurs, whereas methane reacts rapidly with fluorine at room temperature in the absence of UV light or heat. (1)

> 2 9 7 6 A 0 1

Summer 2014

Past Paper

(d) The alkene hex-3-ene reacts with bromine to produce 3,4-dibromohexane. Complete the mechanism below by adding curly arrows to show the movement of electron pairs in both steps and by giving the structural formula of the intermediate carbocation. (3) н н \longrightarrow H₅C₂--C--C₂H₂ Br Br Br Br 3,4-dibromohexane (e) The mechanism shown in (d) shows Z-hex-3-ene reacting with bromine. E-hex-3-ene also reacts with bromine to form 3,4-dibromohexane. (i) Draw the structure of *E*-hex-3-ene. (1) (ii) Explain why both Z-hex-3-ene and E-hex-3-ene react with bromine to produce the **same** structural isomer. (1) (Total for Question 21 = 23 marks) 13

2 9 7 6 A 0 1 3 2

22 Lattice energy can be used as a measure of ionic bond strength. Born-Haber cycles can be used to determine experimental values of lattice energies.

The table below shows the energy changes that are needed to determine the lattice energy of lithium fluoride, LiF.

Energy change	ΔH / kJ mol ⁻¹
Enthalpy change of atomization of lithium	+159
First ionization energy of lithium	+520
Enthalpy change of atomization of fluorine, $\frac{1}{2}F_2$	+79
First electron affinity of fluorine	-328
Enthalpy change of formation of lithium fluoride	-616

(a) Define the term **lattice energy**.

(2)



(4)

- (b) The diagram below shows an incomplete Born-Haber cycle for the formation of lithium fluoride from lithium and fluorine.
 - (i) Complete the diagram by writing the formulae of the correct species, including state symbols, in the four empty boxes.



(ii) Calculate the lattice energy of lithium fluoride, in kJ mol⁻¹.

(2)





*(c) The lattice energies of sodium fluoride, sodium chloride and magnesium fluoride are shown in the table below.

Compound	Lattice energy / kJ mol ⁻¹
Sodium fluoride, NaF	-918
Sodium chloride, NaCl	-780
Magnesium fluoride, MgF ₂	-2957

Explain, in terms of the sizes and charges of the ions involved, the differences between the lattice energy values of

(i) NaF and NaCl

(2)

(ii) NaF and MgF₂

(2)

(Total for Question 22 = 12 marks)



ast P	aper This resource was created and owned by Pearson Edexcel		WCH0
23	Alkanes are used as fuels in homes and in industry. It is, therefore, important that the enthalpy changes involving alkanes are known.		
	(a) Define the term standard enthalpy change of formation of a compound.		
	Give the conditions of temperature and pressure that are used when measuring a standard enthalpy change.	(3)	
De	finition		
Sta	andard temperature is		
Sta	andard pressure is		
	(b) Write the equation, with state symbols, that accompanies the enthalpy change of formation of hexane, C ₆ H ₁₄ (I).	(2)	
	$\begin{array}{ $	Turn	17 over

(c) Enthalpy changes can be calculated using enthalpy changes of combustion. Values for some standard enthalpy changes of combustion are shown in the table below. Substance $\Delta H_{c}^{\leftrightarrow}$ / kJ mol⁻¹ C(s) -394 $H_2(g)$ -286 $CH_4(g)$ -890 Use these data to complete the Hess cycle below for the reaction and then calculate the standard enthalpy change for the reaction, in kJ mol⁻¹. C(s) 2H₂(g) $CH_4(g)$ + \rightarrow (3) Space for working standard enthalpy change for the reaction = kJ mol⁻¹ 18 P 4 2 9 7 6 A 0 1 8 2 4

Summer 2014

st Paper	2014	This resource	was created and c	wined by Pearson Edexcel	Chemistry	WC
(d)				carbon atoms and gaseous hane, C_2H_6 , are shown below.		
			$\rightarrow CH_4(g)$			
				$\Delta H = -2825 \text{ kJ mol}^{-1}$		
	Use these da	ata to calculate				
	(i) the mear					
					(1)	
	(ii) the bond	d enthalpy of a C—	-C bond, in kJ m	ol ⁻¹ , clearly showing your wor	king. (2)	
					(~)	
				(Total for Question 23 = 1	11 marks)	
				(
				AN AANAN NANA NANA NANA MAANA		1

- **24** This question is about atomic structure.
 - (a) Draw diagrams to show the shape of an s-orbital and of a p-orbital.

(2)

s-orbital	p-orbital

(b) Complete the table to show the number of electrons that **completely** fill the following regions.

(3)

Region	Number of electrons present when completely filled
a d-orbital	
a p sub-shell	
the third shell $(n = 3)$	



(2)

- (c) The energy diagram below is for the eight electrons present in an oxygen atom. Complete the diagram for an oxygen atom by adding
 - labels to identify the other occupied sub-shells
 - arrows to show how the remaining six electrons are arranged in the orbitals.



(d) Successive ionization energies provide evidence for the arrangement of electrons in atoms. The eight successive ionization energies of oxygen are shown in the table below.

lonization number	1st	2nd	3rd	4th	5th	6th	7th	8th
lonization energy / kJ mol ⁻¹	1314	3388	5301	7469	10989	13327	71337	84080

(i) Define the term **first ionization energy**.

(3)



(ii) Write an equation, with state symbols, to show the third ionization energy oxygen.	of (2)
*(iii) Explain how the data in the table provide evidence that there are two occupied electron shells in an oxygen atom.	(2)
(Total for Question 24 = 14	
TOTAL FOR SECTION B = 60 TOTAL FOR PAPER = 80	



BLANK PAGE



	0 (8)	(18) 4.0 hetium	2 70 2	Ne Ne	neon 10	39.9	Ar argon 18	83.8	ĸ	krypton 36	131.3	Xe	xenon 54	[222]	Rn	radon 86		_						
	7 0	L.			fluorine 9		chlorine		-	bromine ku 35	126.9 1		iodine >	[210] [astatine 85		Elements with atomic numbers 112-116 have been reported but not fully authenticated	175	Ľ	T1 71	[257]	٦	103 103
	6	(717)	14.0	0.0	oxygen 8	32.1	sulfur 16	79.0		34 t	127.6	Te	tellurium 52	[209]		polonium 84		116 have be ticated	173		ytterbium 1 70	[254]		102
	2	(15)	((1)	0.4 0.4	nitrogen 7	31.0	P phosphorus 15	74.9	As	arsenic 33	121.8	Sb	antimony 51	209.0	Bi	bismuth 83		tomic numbers 112-116 hav but not fully authenticated	169		thutium 69	[256]		mendelevium 101
	4		12 0	C C	carbon 6	28.1	Si Silicon 14	72.6	Ge	germanium 32	118.7	Sn	tin 50	207.2	PP	lead 82		atomic nul but not f	167	٦ E	erbium 68	[253]		100
	m	1617	0.01	0.0 B	boron 5	27.0	Al aluminium 13	69.7	Ga	gallium 31	114.8	드	indium 49	204.4		thallium 81		nents with	165		holmium 67	[254]	ES	97 98 99 99
The Periodic Table of Elements							(12)	65.4	Zn	zinc 30	112.4	B	cadmium 48	200.6	Hg	mercury 80			163	-		[251]	ັບ	catifornium 98
Elen							(11)	63.5	J	copper 29	107.9	Ag		197.0	Au	gold 79	[272]	Rg roentgenium 111	159	10.0		[245]	BK	perketium 97
le of							(10)	58.7	ï	nickel 28	106.4	Pd	palladium 46	195.1	¥.	platinum 78	[271]	DS damstadtium 110	157		gadolinium 64	[247]		ounum 96
c Tab			-				(6)	58.9	ပိ	cobalt 27	102.9		rhodium 45	192.2	<u>ہ</u>	77	[268]	Mt meitnerium 109	152		europium 63	[243]	Am	amencum 95
riodi		1.0 H hydrogen					(8)	55.8	Fe		1	Ru	5	190.2	S	osmum 76	[277]	Hs hassium 108	150		samarıum 62	[242]	Pu	plutonium 94
he Pe							(2)	54.9	Mn	chromium manganese 24 25	[98]	Mo Tc	technetium 43	186.2	Re	75	[264]	Bh bohrium 107	[147]	E E	prometinum 61	[237]	Np Pu Am	93
F				mass bol	number		(9)	52.0	֊	chromium 24	95.9	Wo	molybdenum 42	183.8	3	tungsten 74	[266]	Sg seaborgium 106	144	PZ	59 60 61 61 61	238		uranıum 92
		Kev	Ney .	relative atomic mass atomic symbol	name atomic (proton) number		(2)	50.9	>	vanadium 23	92.9	qN	niobium 41	180.9	Ta	tantalum 73	[262]	dubnium 105	141	۲.	praseodymum 59	[231]	Pa	protactinium 91
			and here	ato	atomic		(4)	47.9	ï	titanium 22	91.2	Zr	zirconium 40	178.5		natnium 72	[261]	Rf nutherfordium 104	140	e.	cerium 58	232	Ę	90
							(3)	45.0	Sc	scandium 21	88.9		yttrium 39	138.9	La*	tantnanum 57	[227]	Ac* actinium 89		es				
	2	0	(7)	Be	beryllium 4	24.3	Mg magnesium 12	40.1		calcium 20	87.6	Sr	strontium 38	137.3	Ba	56	[226]	Ra radium 88		* Lanthanide series	* Actinide series			
	-	(E)	(1)	Li	lithium 3	23.0	sodium 11	39.1	¥	potassium 19	85.5	å	rubidium 37	132.9	ک	55	[223]	Fr francium 87		* Lanth	* Actin			

P 4 2 9 7 6 A 0 2 4 2 4