

Write your name here

Surname	Other names
---------	-------------

**Pearson Edexcel**  
International  
Advanced Level

Centre Number	Candidate Number
<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>

**Chemistry**  
**Advanced Subsidiary**  
**Unit 1: The Core Principles of Chemistry**

Friday 22 May 2015 – Morning <b>Time: 1 hour 30 minutes</b>	Paper Reference <b>WCH01/01</b>
--	------------------------------------

Candidates may use a calculator.	Total Marks
----------------------------------	-------------

### Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided – *there may be more space than you need.*

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*
- Questions labelled with an **asterisk** (\*) are ones where the quality of your written communication will be assessed – *you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.*
- A Periodic Table is printed on the back cover of this paper.

### Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P44880A

©2015 Pearson Education Ltd.

6/6/5/1/



**PEARSON**

## SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box . If you change your mind, put a line through the box  and then mark your new answer with a cross .

1 In which order do the electrons fill the orbitals of an atom?

- A 1s 2s 2p 3s 3p 4s 4p 3d
- B 1s 2s 2p 3s 3d 3p 4s 4p
- C 1s 2s 2p 3s 3p 3d 4s 4p
- D 1s 2s 2p 3s 3p 4s 3d 4p

(Total for Question 1 = 1 mark)

2 Ions are separated in the mass spectrometer by

- A a vacuum pump.
- B a magnetic field.
- C an ionization chamber.
- D electron bombardment.

(Total for Question 2 = 1 mark)

3 Which of the following contains one mole of neutrons?

- A 1 g of  ${}^1_1\text{H}$
- B 1 g of  ${}^{12}_6\text{C}$
- C 2 g of  ${}^{24}_{12}\text{Mg}$
- D 2 g of  ${}^{22}_{10}\text{Ne}$

(Total for Question 3 = 1 mark)



- 4 Solutions of barium chloride and silver nitrate are mixed together. The reaction that takes place is an example of
- A displacement.
  - B neutralization.
  - C oxidation.
  - D precipitation.

(Total for Question 4 = 1 mark)

- 5 The Avogadro constant is numerically equal to the number of
- A ions in 1 mol of sodium chloride, NaCl
  - B atoms in 1 mol of hydrogen gas, H<sub>2</sub>
  - C electrons in 1 mol of helium gas, He
  - D molecules in 1 mol of oxygen gas, O<sub>2</sub>

(Total for Question 5 = 1 mark)

- 6 10 g of magnesium is added to 1 dm<sup>3</sup> of 1 mol dm<sup>-3</sup> copper(II) sulfate solution and the mixture is stirred until no further reaction occurs.

Which of the following is a result of this reaction?

- A The resulting solution is colourless.
- B 10 g of copper is displaced.
- C 63.5 g of copper is displaced.
- D All the magnesium reacts.

(Total for Question 6 = 1 mark)



- 7 Which of the following gas samples has the same volume as 7.0 g of carbon monoxide?

All volumes are measured at the same temperature and pressure.

- A 1.0 g of hydrogen
- B 3.5 g of nitrogen
- C 10.0 g of argon
- D 35.5 g of chlorine

(Total for Question 7 = 1 mark)

- 8 Which of the following aqueous solutions contains the greatest number of **negative** ions?

- A 500 cm<sup>3</sup> of 0.10 mol dm<sup>-3</sup> Na<sub>2</sub>SO<sub>4</sub>(aq)
- B 250 cm<sup>3</sup> of 0.12 mol dm<sup>-3</sup> BaCl<sub>2</sub>(aq)
- C 250 cm<sup>3</sup> of 0.15 mol dm<sup>-3</sup> KI(aq)
- D 500 cm<sup>3</sup> of 0.10 mol dm<sup>-3</sup> Zn(NO<sub>3</sub>)<sub>2</sub>(aq)

(Total for Question 8 = 1 mark)

- 9 In an experiment carried out at 200°C and 1 atm pressure, 20 cm<sup>3</sup> of ammonia gas reacted with an excess of heated copper(II) oxide.




If all measurements were made at 200°C and 1 atm pressure, what would be the total volume, in cm<sup>3</sup>, of gaseous products?

- A 10
- B 20
- C 30
- D 40

(Total for Question 9 = 1 mark)



10 Ammonia is manufactured from hydrogen and nitrogen in the Haber process.



If 60 tonnes of hydrogen produces 80 tonnes of ammonia, what is the percentage yield in the reaction?

- A  $\frac{80}{170} \times 100\%$
- B  $\frac{80}{340} \times 100\%$
- C  $\frac{30}{80} \times 100\%$
- D  $\frac{60}{80} \times 100\%$

(Total for Question 10 = 1 mark)

11 Which of the following compounds has the greatest ionic character?

- A Caesium fluoride
- B Caesium iodide
- C Potassium fluoride
- D Potassium iodide

(Total for Question 11 = 1 mark)

12 Which species has a dative covalent bond?

- A  $\text{H}_3\text{O}^+$
- B  $\text{H}_2\text{O}$
- C  $\text{OH}^-$
- D  $\text{O}_2$

(Total for Question 12 = 1 mark)




13 The atomic radius of potassium is larger than that of sodium because potassium has

- A a larger nuclear charge.
- B a larger nucleus.
- C more occupied electron shells.
- D a smaller first ionization energy.

(Total for Question 13 = 1 mark)

14 Consider the following data.



What is the relationship between a, b, c and d?

- A  $a = b + c - d$
- B  $a = d - b - c$
- C  $a = b - c - d$
- D  $a = d + c - b$

(Total for Question 14 = 1 mark)



15 In which of the following does **X** represent the mean bond enthalpy for the O–H bond in water?

- A  $\text{H}_2\text{O}(\text{g}) \rightarrow \text{O}(\text{g}) + \text{H}_2(\text{g}) \quad \Delta H = 2\text{X}$
- B  $\text{H}_2\text{O}(\text{g}) \rightarrow \text{O}(\text{g}) + 2\text{H}(\text{g}) \quad \Delta H = 2\text{X}$
- C  $\text{H}_2\text{O}(\text{g}) \rightarrow \text{O}(\text{g}) + \text{H}_2(\text{g}) \quad \Delta H = \text{X}$
- D  $\text{H}_2\text{O}(\text{g}) \rightarrow \text{O}(\text{g}) + 2\text{H}(\text{g}) \quad \Delta H = \text{X}$


(Total for Question 15 = 1 mark)

16 Which of the following is a step in the propagation stage of the chlorination of methane?

- A  $\text{Cl}_2 \rightarrow \text{Cl}\cdot + \text{Cl}\cdot$
- B  $\text{CH}_3\cdot + \text{Cl}\cdot \rightarrow \text{CH}_3\text{Cl}$
- C  $\text{CH}_3\cdot + \text{Cl}_2 \rightarrow \text{CH}_3\text{Cl} + \text{Cl}\cdot$
- D  $\text{CH}_4 + \text{Cl}\cdot \rightarrow \text{CH}_3\text{Cl} + \text{H}\cdot$

(Total for Question 16 = 1 mark)

17 A molecule of **Z** has the following structure:



Molecule of **Z**

What are the total numbers of  $\sigma$ -bonds and  $\pi$ -bonds in a molecule of **Z**?


	Number of $\sigma$ -bonds	Number of $\pi$ -bonds
<input type="checkbox"/> A	3	11
<input type="checkbox"/> B	8	3
<input type="checkbox"/> C	11	3
<input type="checkbox"/> D	14	6

(Total for Question 17 = 1 mark)




18 Which is the structure of Z-1,2-dibromoprop-1-ene?


A




B



C



D




(Total for Question 18 = 1 mark)






19 The skeletal formula of 3-methylcyclobut-1-ene is shown below.




What is the skeletal formula of 1-ethyl-3-methylcyclopent-1-ene?


A




B



C



D



(Total for Question 19 = 1 mark)



20 Which of the following fuels, when burned, would make no significant contribution to climate change?

- A Hydrogen
- B Methane
- C Petrol
- D Coal

(Total for Question 20 = 1 mark)

---

**TOTAL FOR SECTION A = 20 MARKS**



BLANK PAGE







P 4 4 8 8 0 A 0 1 1 2 8

## SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

21 A propellant for a rocket consists of a fuel, kerosene, and an oxidizer, liquid oxygen.

(a) The formulae of some hydrocarbons present in kerosene are shown in the table below.

Hydrocarbon	Formula
<b>A</b>	
<b>B</b>	$\text{CH}_3(\text{CH}_2)_{10}\text{CH}_3$
<b>C</b>	
<b>D</b>	
<b>E</b>	

(i) Name the homologous series to which the compounds **A**, **B**, **C** and **E** belong.

(1)



(ii) Name the compound **A**.

(1)

(iii) Explain the term **structural isomers**, by reference to two molecules selected from the table in part (a).

(3)

(iv) Give the **molecular** formula of the compound **D**.

(2)



- (b) In the petrochemical industry, other fuels are obtained by the cracking and reforming of kerosene.

Using appropriate letters, **A** to **D**, identify a molecule listed in the table that could be formed from **E** by

- (i) cracking alone

(1)

- (ii) cracking and then reforming

(1)

- (c) Suggest how engine performance is improved by using a fuel containing the molecule that you have identified in (b)(ii).

(1)

- (d) The **energy density** of a fuel is defined as the energy produced per kilogram of fuel.

Calculate the energy density of dodecane,  $C_{12}H_{26}$ , in  $\text{kJ kg}^{-1}$ . Give your answer to **two** significant figures.

The enthalpy change of combustion of dodecane is  $-8086 \text{ kJ mol}^{-1}$ .

[Molar mass:  $C_{12}H_{26} = 170 \text{ g mol}^{-1}$ ]

(3)

energy density = .....  $\text{kJ kg}^{-1}$

(Total for Question 21 = 13 marks)




BLANK PAGE



P 4 4 8 8 0 A 0 1 5 2 8

22 Born-Haber cycles can be used to determine experimental values of lattice energies.

(a) The diagram below shows a Born-Haber cycle for the formation of strontium chloride from strontium and chlorine.



Using symbols from  $\Delta H_1$  to  $\Delta H_7$  as appropriate, identify the

(i) enthalpy change of atomization of strontium

(1)

(ii) bond enthalpy of chlorine

(1)

(iii) first electron affinity of chlorine

(1)

(iv) enthalpy change of formation of strontium chloride

(1)





- (b) The table below shows the energy changes that are needed to determine the lattice energy of strontium chloride,  $\text{SrCl}_2$ .

Energy change	$\Delta H / \text{kJ mol}^{-1}$
enthalpy change of atomization of strontium	+164
first ionization energy of strontium	+550
second ionization energy of strontium	+1064
enthalpy change of atomization of chlorine, $\frac{1}{2}\text{Cl}_2$	+122
first electron affinity of chlorine	-349
enthalpy change of formation of strontium chloride	-829

- (i) Define the term **lattice energy**.

(2)

.....

.....

.....

.....

- (ii) Calculate the lattice energy of strontium chloride, in  $\text{kJ mol}^{-1}$ .

(2)

lattice energy = .....  $\text{kJ mol}^{-1}$



\*(c) The lattice energies of sodium fluoride and magnesium fluoride are shown in the table below.

Compound	Lattice energy / $\text{kJ mol}^{-1}$
Sodium fluoride, NaF	-918
Magnesium fluoride, $\text{MgF}_2$	-2957

Explain, in terms of the sizes and charges of the ions involved, why the lattice energy of  $\text{MgF}_2$  is more negative than that of NaF.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 22 = 11 marks)



BLANK PAGE



P 4 4 8 8 0 A 0 1 9 2 8

23 This question is about alkenes.

(a) Give the general formula for the homologous series of alkenes.

(1)

---

(b) Give the **structural** formula of the organic product formed when **ethene**,  $\text{CH}_2=\text{CH}_2$ , reacts with

(i) hydrogen

(1)

(ii) chlorine

(1)

(iii) acidified aqueous potassium manganate(VII)

(1)

(iv) bromine **water**

(1)



(c) When **propene**,  $\text{CH}_3\text{CH}=\text{CH}_2$ , reacts with hydrogen chloride, there are **two** possible products, a major product and a minor product.

(i) Draw the **displayed** formulae of these products.

(2)

Major product	Minor product

(ii) Give the mechanism for the reaction of **propene** with hydrogen chloride which forms the major product.

(3)



(d) Propene can be polymerized.

- (i) Write a balanced equation for the polymerization of propene to form poly(propene), drawing the **displayed** formula of the repeat unit of poly(propene).

(3)

- (ii) State a problem associated with the disposal of waste poly(propene).

(1)

.....

.....



- (e) Standard enthalpy changes of combustion can be used to calculate the standard enthalpy change of formation of propene.



Values for some standard enthalpy changes of combustion,  $\Delta H_c^\ominus$ , are shown in the table below.

Substance	$\Delta H_c^\ominus / \text{kJ mol}^{-1}$
C(s)	-394
H <sub>2</sub> (g)	-286
C <sub>3</sub> H <sub>6</sub> (g)	-2058

- (i) Complete the Hess cycle below to enable you to calculate  $\Delta H_f^\ominus$  from combustion data.

(1)



- (ii) Calculate  $\Delta H_f^\ominus$ , in kJ mol<sup>-1</sup>.


(2)

standard enthalpy change of formation of propene = ..... kJ mol<sup>-1</sup>

**(Total for Question 23 = 17 marks)**



24 The diagram below shows the pattern in the first ionization energies of the first 18 elements.



(a) Give the equation, including state symbols, for the first ionization energy of fluorine.

(2)

\*(b) Explain why there is a **general** increase in the first ionization energies from sodium to argon.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





(c) \*(i) Explain why the first ionization energy of aluminium ( $Z = 13$ ) is less than that of magnesium ( $Z = 12$ ).

(2)

\*(ii) Explain why the first ionization energy of sulfur ( $Z = 16$ ) is less than that of phosphorus ( $Z = 15$ ).

(2)

(d) The table below, which is incomplete, refers to the elements sodium to sulfur.

Element	Na	Mg	Al	Si	P	S
Melting temperature	low	high				
Structure		giant				
Electrical conductivity		high		X		

(i) Complete the **melting temperature** row by using only the words 'high' or 'low'.

(2)

(ii) Complete the **structure** row by using only the words 'giant' or 'molecular'.


(2)

(iii) Complete the **electrical conductivity** row by using only the words 'high' or 'low'.

(1)



- (e) In an experiment, 2.76 g of sodium completely reacted with water to form 500 cm<sup>3</sup> of aqueous sodium hydroxide.



- (i) Calculate the number of moles of sodium that reacted. (1)

- (ii) Calculate the maximum volume, in dm<sup>3</sup>, of hydrogen that can be formed at room temperature and pressure.

[1 mol of any gas occupies 24 dm<sup>3</sup> at room temperature and pressure.] (2)

- (iii) Calculate the concentration, in mol dm<sup>-3</sup>, of the sodium hydroxide solution, NaOH(aq), formed in the experiment. (2)

---

(Total for Question 24 = 19 marks)

---

**TOTAL FOR SECTION B = 60 MARKS**  
**TOTAL FOR PAPER = 80 MARKS**



**BLANK PAGE**



P 4 4 8 8 0 A 0 2 7 2 8

The Periodic Table of Elements

	1	2											3	4	5	6	7	0 (8)		
	(1)	(2)											(13)	(14)	(15)	(16)	(17)	(18)		
													Key							
													relative atomic mass							
													atomic symbol							
													name							
													atomic (proton) number							
	6.9	9.0											10.8	12.0	14.0	16.0	19.0	4.0		
	Li	Be											B	C	N	O	F	He		
	lithium	beryllium											boron	carbon	nitrogen	oxygen	fluorine	helium		
	3	4											5	6	7	8	9	2		
	23.0	24.3											27.0	28.1	31.0	32.1	35.5	20.2		
	Na	Mg											Al	Si	P	S	Cl	Ne		
	sodium	magnesium											aluminium	silicon	phosphorus	sulfur	chlorine	neon		
	11	12											13	14	15	16	17	10		
	39.1	40.1											69.7	72.6	74.9	79.0	79.9	83.8		
	K	Ca											Ga	Ge	As	Se	Br	Kr		
	potassium	calcium											gallium	germanium	arsenic	selenium	bromine	krypton		
	19	20											31	32	33	34	35	36		
	85.5	87.6											114.8	118.7	121.8	127.6	126.9	131.3		
	Rb	Sr											In	Sn	Sb	Te	I	Xe		
	rubidium	strontium											indium	tin	antimony	tellurium	iodine	xenon		
	37	38											49	50	51	52	53	54		
	132.9	137.3											204.4	207.2	209.0	209.0	[210]	[222]		
	Cs	Ba											Tl	Pb	Bi	Po	At	Rn		
	caesium	barium											thallium	lead	bismuth	polonium	astatine	radon		
	55	56											81	82	83	84	85	86		
	[223]	[226]											80	82	83	84	85	86		
	Fr	Ra											Hg	Pb	Bi	Po	At	Rn		
	francium	radium											mercury	lead	bismuth	polonium	astatine	radon		
	87	88											80	82	83	84	85	86		
	[227]	[226]											80	82	83	84	85	86		
	La*	Ac*											Cu	Ni	Co	Fe	Mn	Zn		
	lanthanum	actinium											copper	nickel	cobalt	iron	manganese	zinc		
	57	89											29	28	27	26	25	30		
	[227]	[227]											107.9	106.4	102.9	101.1	[98]	112.4		
													Ag	Pd	Rh	Ru	Tc	Cd		
													silver	palladium	rhodium	ruthenium	technetium	cadmium		
													47	46	45	44	43	48		
													197.0	195.1	192.2	190.2	186.2	200.6		
													Au	Pt	Ir	Os	Re	Hg		
													gold	platinum	iridium	osmium	rhenium	mercury		
													79	78	77	76	75	80		
													[272]	[271]	[268]	[277]	[264]	[200.6]		
													Rg	Ds	Mt	Hs	Bh	[200.6]		
													roentgenium	darmstadtium	meitnerium	hassium	bohrium	[200.6]		
													111	110	109	108	107	80		
													159	157	152	150	[147]	163		
													Tb	Gd	Eu	Sm	Pm	Dy		
													terbium	gadolinium	europium	samarium	promethium	dysprosium		
													65	64	63	62	61	66		
													[251]	[247]	[243]	[242]	[237]	[251]		
													Cf	Bk	Am	Pu	Np	Cf		
													californium	berkelium	americium	plutonium	neptunium	californium		
													98	97	95	94	93	98		
													[254]	[253]	[256]	[254]	[257]	[254]		
													Fm	No	Md	Lr	Lu	[254]		
													fermium	nobelium	mendeleevium	lawrencium	lutetium	[254]		
													100	102	101	103	71	103		
													100	102	101	103	71	103		
													165	167	169	173	175	175		
													Ho	Er	Tm	Yb	Lu	Lu		
													holmium	erbium	thulium	ytterbium	lutetium	lutetium		
													67	68	69	70	71	71		
													[254]	[253]	[256]	[254]	[257]	[257]		
													Es	Fm	Md	No	Lr	[257]		
													einsteinium	fermium	mendeleevium	nobelium	lawrencium	[257]		
													99	100	101	102	103	103		

Elements with atomic numbers 112-116 have been reported but not fully authenticated

\* Lanthanide series  
\* Actinide series

