

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Tuesday 9 October 2018

Morning (Time: 1 hour 30 minutes)

Paper Reference **WCH01/01**

Chemistry
Advanced Subsidiary
Unit 1: The Core Principles of Chemistry

You must have: Scientific calculator

Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- Questions labelled with an **asterisk** (*) are ones where the quality of your written communication will be assessed
– *you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.*
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P55423A

©2018 Pearson Education Ltd.

5/2/2/2/2/2/




Pearson

SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box . If you change your mind, put a line through the box and then mark your new answer with a cross .

- 1 Silicon dioxide reacts with hydrogen fluoride to form water and a compound with the formula H_2SiF_6 .



The mole ratio of HF to H_2O in the balanced equation is

- A 1:2
 B 3:1
 C 2:1
 D 6:1

(Total for Question 1 = 1 mark)

- 2 For safety reasons, the concentration of lead in paint should not exceed 600 parts per million (ppm) by mass.

Therefore, the mass of lead in one kilogram of paint should not exceed

- A 0.06 g
 B 0.60 g
 C 6.0 g
 D 60 g

(Total for Question 2 = 1 mark)

The solution containing the greatest number of chloride ions is

- A 10 cm³ of $1.00 \times 10^{-2} \text{ mol dm}^{-3} \text{ AlCl}_3$
 B 20 cm³ of $1.50 \times 10^{-2} \text{ mol dm}^{-3} \text{ MgCl}_2$
 C 30 cm³ of $1.50 \times 10^{-2} \text{ mol dm}^{-3} \text{ HCl}$
 D 10 cm³ of $2.50 \times 10^{-2} \text{ mol dm}^{-3} \text{ CaCl}_2$

(Total for Question 3 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4 Which statement is true about the ions $^{55}\text{Mn}^{2+}$ and $^{56}\text{Fe}^{2+}$?

- A $^{55}\text{Mn}^{2+}$ is deflected less in a mass spectrometer than $^{56}\text{Fe}^{2+}$.
- B They have the same number of electrons.
- C $^{55}\text{Mn}^{2+}$ has more protons than $^{56}\text{Fe}^{2+}$.
- D They have the same number of neutrons.

(Total for Question 4 = 1 mark)

5 10cm^3 of a $1.00 \times 10^{-2}\text{ mol dm}^{-3}$ solution needs to be diluted to make the concentration $5.00 \times 10^{-4}\text{ mol dm}^{-3}$.

What volume of water, in cm^3 , should be added?

- A 20
- B 40
- C 190
- D 200

(Total for Question 5 = 1 mark)

6 The Avogadro constant is $6.0 \times 10^{23}\text{ mol}^{-1}$.

The number of atoms in 15 g of nitrogen monoxide, NO, is


- A 3.0×10^{23}
- B 6.0×10^{23}
- C 2.0×10^{24}
- D 9.0×10^{23}

(Total for Question 6 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



- 7 Nitrogen monoxide reacts with oxygen to form nitrogen dioxide.



200 cm³ of nitrogen monoxide is mixed with 350 cm³ of oxygen.

What is the total volume, in cm³, of the gaseous mixture when the reaction is complete?

All volumes are measured at the same temperature and pressure.

- A 200
- B 350
- C 450
- D 550

(Total for Question 7 = 1 mark)

- 8 The first six successive ionisation energies of an element X are given in the table.

Ionisation energy	1st	2nd	3rd	4th	5th	6th
Value/kJ mol ⁻¹	739	1577	3232	4356	16091	19785

The formula of the oxide of X is most likely to be

- A XO₂
- B XO₃
- C X₂O
- D X₂

(Total for Question 8 = 1 mark)

- 9 The total number of occupied orbitals in the **third** quantum shell of a silicon atom in its ground state is

- A 2
- B 3
- C 4
- D 5

(Total for Question 9 = 1 mark)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



10 Which of these statements is correct?

- A The ionic radii of the alkali metals increase down the group.
- B The ionic radii for the ions Na^+ , Mg^{2+} , Al^{3+} increase across this series.
- C The first ionisation energies of the alkali metals increase down the group.
- D The melting temperatures of successive elements in Period 3 always increase across the period.

(Total for Question 10 = 1 mark)

11 Which compound would be expected to show the greatest covalent character?

- A LiBr
- B LiI
- C KF
- D KCl

(Total for Question 11 = 1 mark)

12 Phosphoric(V) acid, H_3PO_4 , can be made from phosphorus in two stages.



Data

Formula	P_4	O_2	P_4O_{10}	H_2O	H_3PO_4
Molar mass / g mol^{-1}	124	32	284	18	98


The percentage atom economy, by mass, for the production of phosphoric(V) acid from phosphorus is

- A 38.0
- B 69.0
- C 72.4
- D 100

(Total for Question 12 = 1 mark)



13 This question is about the reaction of nickel(II) carbonate and hydrochloric acid.



(a) The ionic equation for this reaction is

(1)

- A $\text{NiCO}_3(\text{s}) + 2\text{H}^+(\text{aq}) \rightarrow \text{Ni}^{2+}(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$
- B $\text{Ni}^{2+}(\text{s}) + 2\text{Cl}^-(\text{aq}) \rightarrow \text{NiCl}_2(\text{aq})$
- C $\text{Ni}^{2+}(\text{s}) + 2\text{HCl}(\text{aq}) \rightarrow \text{NiCl}_2(\text{aq}) + 2\text{H}^+(\text{aq})$
- D $\text{NiCO}_3(\text{s}) + 2\text{HCl}(\text{aq}) \rightarrow \text{Ni}^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq}) + \text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{l})$

(b) Excess hydrochloric acid reacts with 0.20 mol of nickel(II) carbonate. What is the volume, in dm^3 , of gas produced at room temperature and pressure? (1 mol of any gas occupies 24 dm^3 at room temperature and pressure)

(1)

- A 1.2
- B 2.4
- C 4.8
- D 9.6

(c) What is the minimum volume of hydrochloric acid with a concentration of 4.0 mol dm^{-3} that reacts with 0.20 mol of nickel carbonate?

(1)

- A 20 cm^3
- B 50 cm^3
- C 100 cm^3
- D 200 cm^3

(Total for Question 13 = 3 marks)

Use this space for any rough working. Anything you write in this space will gain no credit.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- 14 When 100 cm³ of 2.0 mol dm⁻³ sodium hydroxide solution is added to 100 cm³ of 2.0 mol dm⁻³ sulfuric acid (an excess) to form sodium sulfate, the temperature rise is 12.5 °C.


Energy transferred (J) = mass × 4.2 × temperature change

What is the enthalpy change of the reaction in kJ mol⁻¹?

- A $\Delta H = -\frac{200 \times 4.2 \times 12.5}{0.4}$
- B $\Delta H = -100 \times 4.2 \times 12.5 \times 0.4$
- C $\Delta H = -\frac{200 \times 4.2 \times 12.5}{0.2}$
- D $\Delta H = -100 \times 4.2 \times 12.5 \times 0.2$

(Total for Question 14 = 1 mark)

- 15 Hydrogen is manufactured using the reaction



The percentage yield of hydrogen in this process is 90%.

The mass of hydrogen, in tonnes, which can be produced from 160 tonnes of methane is


- A 27
- B 54
- C 60
- D 63

(Total for Question 15 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



16 The repeat unit of a polymer is shown.



What is the systematic name of the monomer which forms this polymer?

- A 2-ethylbut-2-ene
- B 2,3-dimethylbut-1-ene
- C 2-ethylpent-2-ene
- D 3-methylpent-2-ene

(Total for Question 16 = 1 mark)


17 A compound contains 31.25% Ca, 18.75% C and 50.00% O.

Its empirical formula is

- A CaC_2O_4
- B Ca_2CO_3
- C Ca_2CO_2
- D CaCO_3

(Total for Question 17 = 1 mark)

18 Which reagent reacts with but-2-ene to form the compound with the formula shown?



- A Water
- B Sodium hydroxide
- C Hydrogen peroxide
- D Acidified potassium manganate(VII)

(Total for Question 18 = 1 mark)

TOTAL FOR SECTION A = 20 MARKS



SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

- 19 A sample of zinc has the relative atomic mass 65.44. The sample contains four isotopes. The abundance of three of these isotopes is shown.

Relative isotopic mass	64	66	67
Abundance (%)	49.00	27.90	4.50

- (a) (i) Use these data to calculate the relative isotopic mass of the fourth isotope.

Show your working, and give your answer to an appropriate number of significant figures.

(3)

- (ii) State and explain what difference, if any, you would expect between the chemical properties of the lightest and heaviest isotopes of zinc.

(1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) Isotopic masses are determined using a mass spectrometer. The sample under investigation is first converted into gaseous ions.

(i) Ions then pass through slits in a series of electrically charged plates.

Give **two** reasons for this procedure.

(2)

(ii) State how ions of different mass are separated.

(1)

(iii) The ions eventually produce a current in the detector. Data from the detector are used to produce a mass spectrum.

State how the horizontal axis of a mass spectrum is labelled.
Give your answer in **words**, not symbols.

(1)

Complete the electronic configuration of an atom of zinc using s p d notation.

(1)



(d) Describe, with the aid of a diagram, the bonding in a sample of zinc. You should state the attractions which hold the particles together in the solid.

(3)

(Total for Question 19 = 12 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

EXAMPLE



EXAMPLAR

BLANK PAGE


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



20 (a) The **second** ionisation energies of some elements in Period 3 are shown on the grid.



(i) Mark on the grid, with a cross, the value you would expect for sulfur. (1)

(ii) Write an equation, including state symbols, for the **second** ionisation of aluminium. (2)



*(iii) Explain why the **second** ionisation energy of aluminium is greater than both the second ionisation energy of magnesium and the second ionisation energy of silicon.

(4)

(iv) Predict, with reason, which element in Period 3 has the highest **second** ionisation energy.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) Magnesium and sulfur both react with chlorine to form chlorides with a formula XCl_2 .

Magnesium chloride, $MgCl_2$, is ionic. Sulfur dichloride, SCl_2 , consists of covalently bonded molecules.

(i) Describe how the electrical conductivity of these two compounds differs.

(1)

(ii) Draw a dot and cross diagram for sulfur dichloride.

Use crosses (x) for electrons in sulfur and dots (•) for electrons in chlorine.
Only show outer shell electrons.

(2)

(iii) Sketch an electron density map of sulfur dichloride.

(1)

(iv) State how the electron density map of magnesium chloride differs from that of sulfur dichloride.

(1)




- (c) The Born-Haber cycle can be used to determine the lattice energy of magnesium chloride. The table below shows the enthalpy changes that are needed.

Energy change	$\Delta H / \text{kJ mol}^{-1}$
Enthalpy change of atomisation of magnesium	+147.7
First ionisation energy of magnesium	+738
Second ionisation energy of magnesium	+1451
Enthalpy change of atomisation of chlorine ($\frac{1}{2}\text{Cl}_2$)	+121.7
First electron affinity of chlorine	-348.8
Enthalpy change of formation of magnesium chloride	-641.3

- (i) The diagram shows an incomplete Born-Haber cycle for the formation of magnesium chloride from magnesium and chlorine.

Complete the diagram by writing the **formulae** of the correct species, including state symbols, on the five empty horizontal lines.

(4)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(ii) Calculate the lattice energy of magnesium chloride in kJ mol^{-1} .

(2)

(Total for Question 20 = 19 marks)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

EXAMPLE



21 The compound hydrazine, N_2H_4 , is a liquid which is used as a rocket fuel.




It reacts with oxygen to form nitrogen and water.

(a) Complete the Hess cycle and, using data in the table, calculate the enthalpy change for the oxidation of hydrazine, $\Delta H_{\text{reaction}}^{\ominus}$.

Species	Standard enthalpy change of formation / kJ mol^{-1}
$\text{N}_2\text{H}_4(\text{l})$	+50.6
$\text{H}_2\text{O}(\text{l})$	-285.8

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) Some bond enthalpies are given in the table.

Bond	Bond enthalpy/kJ mol ⁻¹
N—N	158
O=O	498
N≡N	945
H—O	464
N—H	391

(i) Calculate the enthalpy change for the oxidation of hydrazine, using the bond enthalpy values in the table.



(3)

(ii) Give two reasons why the enthalpy change calculated using bond enthalpies differs from the enthalpy change calculated from the Hess cycle.

(2)

(Total for Question 21 = 7 marks)



22 One component of petrol is decane, $C_{10}H_{22}$.

(a) Decane reacts with chlorine in the presence of ultraviolet light to form a mixture of products.

(i) Complete the equation for the initiation step, including appropriate curly arrows. (2)




(ii) Write equations, using molecular formulae, for **two** propagation steps. (2)

(iii) Write equations, using molecular formulae, for **two** termination steps, other than the one in which chlorine forms. (2)

(b) The structure of decane can be changed by the process called reforming.

Name the compound shown, which can be produced in this process. (1)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (c) Write an equation, using molecular formulae, for the incomplete combustion reaction in which decane reacts to form carbon monoxide and **one** other product.

State symbols are not required.

(1)

- (d) Decane can be cracked to form a mixture of butane, and two different alkenes which have different molecular formulae.

- (i) Write an equation for this reaction, using molecular formulae.
State symbols are not required.

(1)

- * (ii) Explain why geometric isomerism can occur in alkenes and why alkenes produced by this cracking reaction may not have geometric isomers.

(2)

- iii) Draw the structure of the *trans*, (*E*), isomer of an alkene produced by the cracking reaction in (d)(i).

(1)

(Total for Question 22 = 12 marks)



23 This question is about alkenes.

*(a) Describe in detail the structure of the C=C double bond in alkenes and hence explain why alkenes are more reactive than alkanes.

(3)

(b) Hydrogen bromide reacts with propene to form a mixture of 1-bromopropane and 2-bromopropane.

(i) Draw the mechanism for the formation of the major product in the reaction of propene with hydrogen bromide. You should show relevant dipoles and curly arrows.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(ii) State why the amounts of each product are **not** equal.

(1)

(c) A derivative of propene called allyl bromide, or 3-bromoprop-1-ene, is used to make polymers. The formula of allyl bromide is $\text{CH}_2=\text{CHCH}_2\text{Br}$.

Write the equation for the polymerisation of allyl bromide, showing the structure of the polymer.

(2)

(Total for Question 23 = 10 marks)

TOTAL FOR SECTION B = 60 MARKS

TOTAL FOR PAPER = 80 MARKS



The Periodic Table of Elements

(1)	2	3	4	5	6	7	0 (8)		
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)		
6.9 Li lithium	9.0 Be beryllium	11 Na sodium	12 Mg magnesium	13 B boron	14 C carbon	15 N nitrogen	16 O oxygen	17 F fluorine	18 Ne neon
23.0 Al aluminium	24.3 Mg magnesium	27.0 Al aluminium	28.1 Si silicon	31.0 P phosphorus	32.1 S sulfur	35.5 Cl chlorine	39.9 Ar argon	4.0 He helium	20.2 Ne neon
39.1 K potassium	40.1 Ca calcium	45.0 Sc scandium	47.9 Ti titanium	50.9 V vanadium	52.0 Cr chromium	55.8 Mn manganese	58.7 Fe iron	63.5 Cu copper	65.4 Zn zinc
85.5 Rb rubidium	87.6 Sr strontium	88.9 Y yttrium	91.2 Zr zirconium	92.9 Nb niobium	95.9 Mo molybdenum	98 Tc technetium	101 Ru ruthenium	106.4 Rh rhodium	112.4 Cd cadmium
132.9 Cs caesium	137.3 Ba barium	138.9 La* lanthanum	178.5 Hf hafnium	180.9 Ta tantalum	183.8 W tungsten	186.2 Re rhenium	190.2 Os osmium	195.1 Pt platinum	200.6 Hg mercury
223 Fr francium	226 Ra radium	227 Ac* actinium	261 Rf rutherfordium	262 Db dubnium	266 Sg seaborgium	264 Bh bohrium	277 Hs hassium	272 Ds darmstadtium	285 Uu ununseptium
87	88	89	104	105	106	107	108	109	110
140 Ce cerium	141 Pr praseodymium	144 Nd neodymium	147 Pm promethium	150 Sm samarium	152 Eu europium	157 Gd gadolinium	163 Dy dysprosium	165 Ho holmium	167 Er erbium
58	59	60	61	62	63	64	65	66	68
232 Th thorium	231 Pa protactinium	238 U uranium	237 Np neptunium	242 Pu plutonium	243 Am americium	247 Cm curium	251 Cf californium	254 Es einsteinium	253 Fm fermium
90	91	92	93	94	95	96	98	99	100
90 Th thorium	91 Pa protactinium	92 U uranium	93 Np neptunium	94 Pu plutonium	95 Am americium	96 Cm curium	97 Bk berkelium	98 Cf californium	99 Es einsteinium
103 Lr lawrencium	102 Nb niobium	101 Md mendelevium	100 Fm fermium	99 Es einsteinium	98 Cf californium	97 Bk berkelium	96 Cm curium	95 Am americium	94 Pu plutonium
103 Lr lawrencium	102 Nb niobium	101 Md mendelevium	100 Fm fermium	99 Es einsteinium	98 Cf californium	97 Bk berkelium	96 Cm curium	95 Am americium	94 Pu plutonium
175 Lu lutetium	173 Yb ytterbium	173 Tm thulium	169 Yb ytterbium	167 Er erbium	163 Dy dysprosium	159 Tb terbium	157 Gd gadolinium	152 Eu europium	150 Sm samarium
175 Lu lutetium	173 Yb ytterbium	169 Tm thulium	167 Er erbium	163 Dy dysprosium	159 Tb terbium	157 Gd gadolinium	152 Eu europium	150 Sm samarium	147 Pm promethium
175 Lu lutetium	173 Yb ytterbium	169 Tm thulium	167 Er erbium	163 Dy dysprosium	159 Tb terbium	157 Gd gadolinium	152 Eu europium	150 Sm samarium	147 Pm promethium
257 Lv livermorium	256 Uu ununseptium	255 Uub ununbium	254 Uut ununtrium	253 Uuq ununquadium	252 Uuq ununquadium	251 Uuq ununquadium	250 Uuq ununquadium	249 Uuq ununquadium	248 Uuq ununquadium
257 Lv livermorium	256 Uu ununseptium	255 Uub ununbium	254 Uut ununtrium	253 Uuq ununquadium	252 Uuq ununquadium	251 Uuq ununquadium	250 Uuq ununquadium	249 Uuq ununquadium	248 Uuq ununquadium
257 Lv livermorium	256 Uu ununseptium	255 Uub ununbium	254 Uut ununtrium	253 Uuq ununquadium	252 Uuq ununquadium	251 Uuq ununquadium	250 Uuq ununquadium	249 Uuq ununquadium	248 Uuq ununquadium

Elements with atomic numbers 112-116 have been reported but not fully authenticated

* Lanthanide series

* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

