

Write your name here

Surname

Other names

**Pearson Edexcel**  
**International**  
**Advanced Level**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

# Further Pure Mathematics F2

## Advanced/Advanced Subsidiary

Wednesday 3 June 2015 – Morning

**Time: 1 hour 30 minutes**

Paper Reference

**WFM02/01****You must have:**

Mathematical Formulae and Statistical Tables (Blue)

Total Marks

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

### Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P44832A

©2015 Pearson Education Ltd.

5/1/1/



PEARSON

Leave  
blank

1. Using algebra, find the set of values of  $x$  for which

$$\frac{x}{x+2} < \frac{2}{x+5}$$

(7)



1

**(Total 7 marks)**



2. (a) Express  $\frac{1}{(r+6)(r+8)}$  in partial fractions.

(1)

(b) Hence show that

$$\sum_{r=1}^n \frac{2}{(r+6)(r+8)} = \frac{n(an+b)}{56(n+7)(n+8)}$$

where  $a$  and  $b$  are integers to be found.

(4)





Leave  
blank

Q2

**(Total 5 marks)**









Leave  
blank







Leave  
blank

Leave  
blank

**(Total 9 marks)**

Q4









Leave  
blank

Leave  
blank

**(Total 9 marks)**

**Q5**



Leave  
blank

- $$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} - 3y = 2\sin x \quad (\text{I})$$

Given that  $y = 0$  and  $\frac{dy}{dx} = 1$  when  $x = 0$

- (b) find the particular solution of differential equation (I).






Leave  
blank



7.



### Figure 1

$$r = \sqrt{3} \sin \theta, \quad 0 \leq \theta \leq \pi$$

$$r = 1 + \cos \theta, \quad 0 \leq \theta \leq \pi$$

- (a) Verify that the curves intersect at the point  $P$  with polar coordinates  $\left(\frac{3}{2}, \frac{\pi}{3}\right)$ . **(2)**

The region  $R$ , bounded by the two curves, is shown shaded in Figure 1.

- (b) Use calculus to find the exact area of  $R$ , giving your answer in the form  $a(\pi - \sqrt{3})$ , where  $a$  is a constant to be found.





Leave  
blank



Leave  
blank

Leave  
blank

**(Total 8 marks)**

**Q7**







Leave  
blank



Leave  
blank

**Q8**

**END**

