This resource was created and owned by Pearson Edexcel

Surname		Other names	
Pearson Edexcel International Idvanced Level	Centre Number		Candidate Number
Advanced Friday 12 June 2015 – Morr	ning		C34 Paper Reference
Advanced Friday 12 June 2015 – Morr Time: 2 hours 30 minutes	ning		C34 Paper Reference WMA02/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 125.
- The marks for **each** question are shown in brackets – use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

P44970A ©2015 Pearson Education Ltd. 5/1/1/1/1/



Turn over 🕨



WMA02 Leave

blank

1. A curve has equation

 $4x^2 - y^2 + 2xy + 5 = 0$

The points P and Q lie on the curve.

Given that $\frac{dy}{dx} = 2$ at *P* and at *Q*,

(a) use implicit differentiation to show that y - 6x = 0 at *P* and at *Q*.

(6)

(3)

(b) Hence find the coordinates of P and Q.



Question Number	Scheme	Marks
1. (a)	$\underbrace{\left(\frac{dy}{dx}\right)}_{dx} = \underbrace{8x - 2y\frac{dy}{dx}}_{dx} + \underbrace{2x\frac{dy}{dx} + 2y}_{dx} = 0$	<u>M1B1</u> A1
	Either Way 1: Sets $\frac{dy}{dx} = 2$ in each term in their differentiated expression	dM1
	$\Rightarrow 8x - 4y + 4x + 2y = 0, \Rightarrow y - 6x = 0 *$	ddM1,A1*
	Or Way 2: Obtains $\frac{dy}{dx} = \left(\frac{8x+2y}{2y-2x}\right)$ (ft their differentiated expression)	dM1
	$\frac{8x+2y}{2y-2x} = 2, \text{ so } y - 6x = 0^*$	ddM1,A1*
		(6)
(b)	Put $y = 6x$ or $x = \frac{y}{6}$ into $4x^2 - y^2 + 2xy + 5 = 0$ and obtains $Ay^2 = B$ or $Ax^2 = B$ where A and B are constants	M1
	$x = \pm \frac{1}{2}$ or $y = \pm 3$ or $(\frac{1}{2}, 3)$ or $(-\frac{1}{2}, -3)$	A1
	both $\left(\frac{1}{2},3\right)$ and $\left(-\frac{1}{2},-3\right)$ and no extra solutions	A1
		(9 marks) (3)

(a) M1 Differentiating
$$4x^2 - y^2$$
 with respect to x to obtain $Ax + By \frac{dy}{dx}$

Condone $\frac{dy}{dx} = \dots$ at start. May not have lost the +5

B1 Sight of
$$\frac{d}{dx}(2xy) = 2x\frac{dy}{dx} + 2y$$

- A1 A fully correct derivative. Accept 8xdx 2ydy + 2xdy + 2ydx = 0 (needs = 0)
- dM1 depends on previous M mark and B mark so has at least two terms in $\frac{dy}{dx}$ and at least two other terms.

Way 1: Sets $\frac{dy}{dx} = 2$ in each term in their differentiated function.

- Or Way 2: May see algebra used to give $\frac{dy}{dx}$ = (condone sign slips and slight copying errors but not omission of terms)
- ddM1 Dependent upon **both** previous M's. It is for proceeding to obtain an unsimplified **correct equation** in *x* and *y* equivalent to those in the scheme e.g. $\frac{-(4x+y)}{x-y} = 2$
- A1* cso y-6x = 0 no errors should have been seen the solutions in the mark scheme would gain full marks (extra lines of working are not required) accept y = 6x
- (b)M1 Substitutes y = 6x or $x = \frac{y}{6}$ into the equation of curve C to form an equation in one variable and reaches two term quadratic $Ax^2 = B$ or $Ay^2 = B$ for any values of A or B $(4x^2 - 36x^2 + 12x^2 + 5 = 0 \Longrightarrow)$ $20x^2 = 5$ $\left(4\left(\frac{y}{6}\right)^2 - y^2 + \frac{y^2}{3} + 5 = 0 \Longrightarrow\right)$ $\frac{5}{9}y^2 = 5$ but condone slips in the working
 - A1 Either one correct pair of coordinates or both *x* or both *y* values.
 - A1 **Both** $\left(\frac{1}{2},3\right)$ and $\left(-\frac{1}{2},-3\right)$ (two answers correct with no incorrect working implies M1A1A1) Any extra solutions (obtained by substituting the variable found first into the quadratic, instead of the linear equation for example) result in A0.

Unsimplified answers lose the final A mark e.g. $\left(\frac{\sqrt{5}}{\sqrt{20}}, 3\right)$ and $\left(-\frac{\sqrt{5}}{\sqrt{20}}, -3\right)$

Allow (0.5, 3) and (-0.5, -3) – need not be in brackets if pairing is clear but wrong pairing is A0.

$$x = \frac{1}{2}, y = 3$$
 and $x = -\frac{1}{2}, y = -3$ is acceptable but $x = \pm \frac{1}{2}, y = \pm 3$ is not sufficient

WMA02 Leave

blank

2. Given that

$$\frac{4(x^2+6)}{(1-2x)(2+x)^2} \equiv \frac{A}{(1-2x)} + \frac{B}{(2+x)} + \frac{C}{(2+x)^2}$$

(a) find the values of the constants A and C and show that B = 0

(b) Hence, or otherwise, find the series expansion of

$$rac{4(x^2+6)}{\left(1-2x
ight)(2+x)^2}, \hspace{0.5cm} |x|<rac{1}{2}$$

in ascending powers of x, up to and including the term in x^2 , simplifying each term.

(5)

(4)



Question Number	Scheme	Marks
2(a)	$4(x^{2}+6) = A(2+x)^{2} + B(1-2x)(2+x) + C(1-2x)$	M1
Way 1	Let $x = -2 \Longrightarrow 40 = 5C \Longrightarrow C = 8$	dM1
	Let $x = \frac{1}{2} \Rightarrow 25 = 6.25 A \Rightarrow A = 4$ $A = 4, C = 8$	A1
	Compare constants / terms in <i>x</i> or substitute another value of <i>x</i> into identity and conclude that $B = 0$ e.g. $24 = 4A + 2B + C \Longrightarrow B = 0 *$	A1* (4)
Way 2 (a)	$4(x^{2}+6) = A(2+x)^{2} + B(1-2x)(2+x) + C(1-2x)$	M1
	Compare x^2 : so $4 = A - 2B$, x : so $0 = 4A - 3B - 2C$, constants: so $24 = 4A + 2B + C$	dM1
	So $A = 4, C = 8$, and $B = 0^*$	A1, A1
Way 1 (b)	$\frac{4(x^2+6)}{(1-2x)(2+x)^2} = 4(1-2x)^{-1} + 8(2+x)^{-2} = 4(1-2x)^{-1} + 8 \times \frac{1}{2^2} \left(1+\frac{x}{2}\right)^{-2}$	B1 ft
	See $\left(1 + (-1)(-2x) + \frac{(-1)(-2)(-2x)^2}{2!}\right)$ or $\left(1 + (-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \right)$	M1
	$\dots \left(1 + (-1)(-2x) + \frac{(-1)(-2)(-2x)^2}{2!}\right) \text{ and } \dots \left(1 + (-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \right)$	A1
	$= 4(1+2x+4x^{2}+)+2\left(1-x+\frac{3}{4}x^{2}\right) \qquad = 6+6x+\frac{35}{2}x^{2}$	dM1A1 (5)
Way 2 (b)	Or $\frac{4(x^2+6)}{(1-2x)(2+x)^2} = 4(x^2+6) \times (1-2x)^{-1} \times \frac{1}{2^2} \left(1+\frac{x}{2}\right)^{-2}$	B1
	See $\left(1 + (-1)(-2x) + \frac{(-1)(-2)(-2x)^2}{2!}\right)$ or $\left(1 + (-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \right)$	M1
	$\dots \left(1 + (-1)(-2x) + \frac{(-1)(-2)(-2x)^2}{2!}\right) \text{ and } \dots \left(1 + (-2)\left(\frac{x}{2}\right) + \frac{(-2)(-3)}{2!}\left(\frac{x}{2}\right)^2 + \right)$	A1
	$= 4(x^{2}+6)(1+2x+4x^{2}+)\times\frac{1}{4}\left(1-x+\frac{3}{4}x^{2}\right) = 6+6x+\frac{35}{2}x^{2}$	dM1A1 (5)
L		(9 marks)

- (a) M1 Uses correct form $4(x^2+6) = A(2+x)^2 + B(1-2x)(2+x) + C(1-2x)$ allow sign errors.
 - dM1 Uses correct method, either substitution or equating coefficients of terms to find at least one constant (see Way 1 and Way 2 above)
 - A1 Both A = 4, C = 8 (Correct answers with no working (cover up rule) imply M1M1A1)

A1* This **needs a method** as the answer is given and **needs to conclude that** B = 0. For method: May compare constants i.e. 24 = 4A + 2B + C with A = 4 and C = 8, or terms in x i.e. 0 = 4A - 3B - 2C with A = 4 and C = 8, or term in x^2 : so 4 = A - 2B with A = 4or substitute another value for x such as x = 0 or x = 1 ...with A = 4 and C = 8.

(b) Way 1

B1ft Writes their expression in the form $A(1-2x)^{-1} + C \times \frac{1}{2^2} \left(1 + \frac{x}{2}\right)^{-2}$ ft on values of A and C.

There should be no *B* term.

(This may be awarded for writing or using $A(1-2x)^{-1} + C(2+x)^{-2}$ and writing

separately in their solution $C(2+x)^{-2} = C \times \frac{1}{2^2} \left(1 + \frac{x}{2}\right)^{-2}$) This could appear in the

binomial expansions.

- M1 Uses the binomial expansion correctly for one expansion, with power of 2 outside the second bracket ignored. Allow missing brackets for this mark. Ignore constants outside the bracket for this method mark. One completely correct expansion in the bracket
- A1 'Both' expansions in the brackets correct and unsimplified. Can be awarded for the two completely correct expansions in the brackets without mention of "*A*" and /or "*C*" or adding and with power of 2 outside the second bracket ignored.
- dM1 Multiplies out brackets and collects terms. Dependent upon **previous M** mark. Allow sign slips.

A1
$$6+6x+\frac{35}{2}x^2$$
. May be written as list or in reverse order. Implies previous M mark.

Way 2

B1 Writes
$$\frac{4(x^2+6)}{(1-2x)(2+x)^2} = 4(x^2+6) \times (1-2x)^{-1} \times \frac{1}{4} \left(1+\frac{x}{2}\right)^{-2}$$
. This could be given after

the two binomial expansions. (It may be awarded for writing

$$4(x^{2}+6) \times (1-2x)^{-1} \times (2+x)^{-2}$$
 and writing separately $(2+x)^{-2} = \frac{1}{2^{2}} \left(1+\frac{x}{2}\right)^{-2}$

M1A1: Follow the scheme and the notes for Way 1

dM1 Must multiply out three brackets for this method

A1
$$6+6x+\frac{35}{2}x^2$$
. May be written as list or in reverse order. Implies previous M mark.

N.B.
$$(2+x)^{-2} = 2^{-2} + (-2)2^{-3}(x) + \frac{(-2)(-3)}{2!}2^{-4}(x)^2 + ... = (\frac{1}{4} - \frac{1}{4}x + \frac{3}{16}x^2)$$
 is an alternative

correct expansion and implies the B1 mark as well as contributing to the M1(one correct expansion) A1 (two correct expansions)





Question Number	Scheme	Marks
3 (a)	$f'(x) = e^x \times 2 + (2x - 5)e^x$	M1A1
	$f'(x) = 0 \Longrightarrow (2x-3)e^x = 0 \Longrightarrow x = \frac{3}{2}$	M1A1
	{Coordinates of $A = \left(\frac{3}{2}, -2e^{\frac{3}{2}}\right)$ } obtains $y = -2e^{\frac{3}{2}}$	A1ft
		(5)
(b)	$-2e^{\frac{3}{2}} < k < 0$	M1A1
		(2)
(c)	(0,5) $(0,5)$ $(0,5$	B1
	$\left(\frac{5}{2},0\right)$ only	B1
	(0,5) only	B1
		(3)
		(10 marks)

- (a) M1 For applying the product rule correctly to get a form $Ae^{x} + (2x-5)e^{x}$ with A a constant
 - A1 correct differentiation need not be simplified isw
 - M1 Sets their f'(x) = 0 and proceeds to x = ...
 - A1 Correct answer in any equivalent form. (Ignore extra answers such as ln0 or even 0)
 - A1ft For finding their correct exact *y* coordinate(ft their *x*) .Allow even if positive. May not be given as coordinates.

Allow $y = -2e^{\frac{3}{2}}$ for this mark

(b) M1 Uses *their* minimum y value as lower limit.

Accept $x \ge \text{their} - 2e^{\frac{3}{2}}$ or $x > \text{their} - 2e^{\frac{3}{2}}$ (may use any letter for the M mark)

- A1 $-2e^{\overline{2}} < k < 0$ with both inequalities strictly less than this is cao (so $-2e^{\overline{2}} \le k < 0$ is A0). Need to use k this time, not y and need exact lower limit.
- (c) B1 Correct shape curve lies only in first two quadrants with maximum in first quadrant. Tends to x axis as x becomes large and negative . Crosses y axis and touches x axis with discontinuous gradient –cusp- (not a minimum point). Then gradient becomes steeper as x becomes large and Positive. (Give bod if curve looks like straight line here but must not bend back on itself)
 - B1 Correct x coordinate- may be on sketch or in the text. Diagram takes priority over text if there is a contradiction. Need both coordinates if in the text but on the x axis the 2.5 is sufficient (even allow (0, 2.5) on x axis) **Must be the only crossing point**.
 - B1 Correct *y* coordinate– may be on sketch or in the text. Need both coordinates if in the text but on the *y* axis the 5 is sufficient (even allow (5,0) on *y* axis). Must be only crossing point.

Summer Past Paper	2015 www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics C34 WMA02
4.	A A B Figure 2	Leave blank
	Figure 2 shows the points A and B with position vectors a and b respectively, fixed origin O .	relative to a
	Given that $ a = 5$, $ b = 6$ and $a.b = 20$	
	a) find the cosine of angle <i>AOB</i> ,	(2)
	b) find the exact length of AB .	(2)
	c) Show that the area of triangle <i>OAB</i> is $5\sqrt{5}$	(3)
12		

Question Number	Scheme	Marks	
4 (a)	Uses $\mathbf{a} \cdot \mathbf{b} = \mathbf{a} \mathbf{b} \cos \theta \Rightarrow 20 = 5 \times 6 \cos \theta \Rightarrow \cos \theta = \frac{20}{30} = \left(\frac{2}{3}\right)$	M1A1	
(b)	$c^{2} = a^{2} + b^{2} - 2ab \cos C$ Uses $\Rightarrow c^{2} = 5^{2} + 6^{2} - 2 \times 5 \times 6 \times \frac{2}{3}$ $\Rightarrow c^{2} = 5^{2} + 6^{2} - 2 \times 20$ $\Rightarrow \overline{ AB } = \sqrt{21}$	M1 A1	(2)
(c)	Uses any method (or no method) with their $\cos \theta = \frac{2}{3} \implies \sin \theta = \left(\frac{\sqrt{5}}{3}\right)$ or gives exact height $= \left(\frac{5\sqrt{5}}{3}\right)$	M1	(2)
	Area of triangle $OAB = \frac{1}{2} \times 5 \times 6 \times \sin(AOB)$	M1	
	$\frac{1}{2} \times 5 \times 6 \times \sin(AOB) = 5\sqrt{5}$ (no evidence of calculator and clear working with surds) (See notes for other methods)	A1* (7 marks)	(3)
		· · · · · ·	

(a) M1 Uses
$$\mathbf{a}.\mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$$
 to obtain $\cos \theta = ...$ or writes $\theta = \arccos\left(\frac{20}{6 \times 5}\right)$

A1 Obtains $\cos \theta = \frac{20}{30}$ or $\frac{2}{3}$ then isw (This answer implies M1A1) (isw if they go on to find the angle)

$$\cos\theta = \frac{20}{5 \times 6} \quad \text{earns M1A0}$$

Special case: Uses printed answer to part (c) to find angle then deduces $\cos \theta = \frac{2}{3}$ is M0A0

- (b) M1 Uses correct version of the cosine rule with their $\cos \theta = ..$ or $\theta = ..$ to find $|\overrightarrow{AB}|^2 = or c^2 =$ (may include non exact angles or cosines) May be done by splitting into two right angles triangles correctly and using trigonometry and Pythagoras. Method must be completely correct. A1 For correct **exact** answer only
- (c) M1 Uses any method with their exact numerical $\cos\theta$ to find exact value of $\sin\theta$

Could use $\sin^2 \theta + \cos^2 \theta = 1$ or Pythagoras' theorem on Use of angle = 48.19 or any **non exact** work is M0

Just writes down $\sin \theta = \left(\frac{\sqrt{5}}{3}\right)$ should be given the mark bod

May find height of triangle by correct Pythagoras using $5^2 - 25\cos^2 \theta = h^2$

M1 Uses a correct method - the formula $\frac{1}{2}5 \times 6 \sin C$ is most likely but may find height of

triangle by trigonometry (or Pythagoras see first M1 mark) and use $\frac{1}{2}b \times h$, with

- values for *b* and *h* (usually 6 and $5\sin C$) For this mark **non exact work may be seen**. A1 Completely correct **exact** work (without using calculator approximation) and **states answer** Sight of 0.74535.. for $\sin \theta$ or 3.726.. for height of triangle or the numerical value of the angle (48.2) should be awarded A0
- (c)Alternative Method: A neat method is to use

1

Area = $\frac{1}{2}\sqrt{\{(\mathbf{a}.\mathbf{a})(\mathbf{b}.\mathbf{b}) - (\mathbf{a}.\mathbf{b})^2\}} = \frac{1}{2}\sqrt{\{(25)(36) - (20)^2\}}$ This is M2 then A1 for achieving the printed answer

Alternative method for (c) using Heron's formula – can get M1M1A1

- M1 Attempts to use $A = \sqrt{s(s-a)(s-b)(s-c)}$ with $s = \frac{a+b+c}{2}$ and a = 5, b = 6 and c = their **exact** answer to part (b)
- M1 Attempts to simplify their $\sqrt{\left(\frac{11+\sqrt{21}}{2}\right)\left(\frac{11+\sqrt{21}}{2}-5\right)\left(\frac{11+\sqrt{21}}{2}-6\right)\left(\frac{11+\sqrt{21}}{2}-\sqrt{21}\right)}$ by any means (even using a calculator) so this may be inexact

A1 Needs to show
$$A = \frac{1}{4}\sqrt{(11+\sqrt{21})(1+\sqrt{21})(\sqrt{21}-1)(11-\sqrt{21})} = \frac{1}{4}\sqrt{(121-21)(21-1)}$$

$$=\frac{1}{4}\sqrt{(2000)} = 5\sqrt{5}$$
 (Seeing calculator approximation is A0 e.g. 14.3)

		Leave blank
5. (i)	Find the x coordinate of each point on the curve $y = \frac{x}{x+1}$, $x \neq -1$, at which the	
	gradient is $\frac{1}{4}$ (4)	
(ii) Given that	
	$\int_{a}^{2a} \frac{t+1}{t} \mathrm{d}t = \ln 7 \qquad a > 0$	
	find the exact value of the constant <i>a</i> . (4)	

Question Number	Scheme	Marks	
5.(i)	$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{(x+1)-x}{(x+1)^2} = \frac{1}{(x+1)^2} \text{ or } y = \frac{x}{x+1} = 1 - \frac{1}{x+1} \Longrightarrow \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{(x+1)^2}$ (see notes for further methods)	M1	
	$\frac{1}{(x+1)^2} = \frac{1}{4} or (x+1)^2 = 4 or x^2 + 2x + 1 = 4$	A1	
	x = 1, -3	M1 A1	
			(4)
(ii)	$\int \frac{t+1}{t} dt = \int 1 + \frac{1}{t} dt = t + \ln t (+c) \text{see notes for integration by parts.}$	M1A1	
	$[t+\ln t]_a^{2a} = \ln 7 \Longrightarrow 2a + \ln 2a - a - \ln a = \ln 7$		
	$a + \ln\left(\frac{2a}{a}\right) = \ln 7 \Rightarrow a = \ln\left(\frac{7}{2}\right) \text{ or } a = \ln 7 - \ln 2$	dM1A1	
	(") (2)		(4)
		(8 marks)	

Correct use of quotient, product, implicit differentiation OR chain rules - may not be M1 (i) simplified – accept any correct answer or correct formula quoted followed by slip.

Correct answers are: $-(-1)(x+1)^{-2}$ (chain rule)

$$(x+1)^{-1} + (-1)x(x+1)^{-2}$$
 (product rule)

$$y(x+1) = x$$
 gives $(x+1)\frac{dy}{dx} + y = 1$ so $\frac{dy}{dx} = \frac{1-y}{x+1}$ or $\frac{1}{(x+1)^2}$ (implicit differentiation)

 $\frac{dy}{dx} = \frac{1}{4}$ and proceeds to one of the listed correct equations in x only. A1

i.e.
$$\frac{1}{(x+1)^2} = \frac{1}{4} or (x+1)^2 = 4 or x^2 + 2x + 1 = 4$$

- M1 Solves their quadratic by usual methods (see notes) to obtain two values for x = ...
- Need both answers two correct answers for x with no working implies the method here. A1 Ignore *y* values if they are given as well.
- Writes as a sum of 1 and t^{-1} and integrates this sum giving a sum with one correct term (ii) M1 e.g. $t + t^{-2}$

May use parts (see below for two variants on parts) If parts are used they must be used accurately (as given below). It is not the most efficient method here and usually results in errors and no marks.

- A1 Both terms correct (ignore arbitrary constant) e.g. From parts may obtain $t + \ln t + 1$
- Uses limits correct way round and sets $I(2a) I(a) = \ln 7$ and also uses or states dM1 $\ln 2a - \ln a = \ln 2$ leading to a = ...This is dependent upon the previous M having been scored
- A1 Correct answer. Accept any correct equivalent e.g. ln 3.5. $\ln 7 - \ln 2$ is A1 but if it is followed by $\ln 5$ this is A0

Parts in integration:

Either:
$$\int \frac{t+1}{t} dt = \frac{1}{t} \left(\frac{t^2}{2} + t \right) - \int -\frac{1}{t^2} \left(\frac{t^2}{2} + t \right) dt = \frac{t}{2} + 1 + \int \frac{1}{2} + \frac{1}{t} dt \text{ for M1}$$
$$= \frac{t}{2} + 1 + \frac{t}{2} + \ln t \text{ for A1 then as before}$$
Or:
$$\int \frac{t+1}{t} dt = (t+1)\ln t - \int \ln t \, dt \text{ for M1}$$

 $= (t+1) \ln t - t \ln t + t$ for A1 then as before

I mme st Paper	r 2015 www.mystudybro.com r This resource was created and owned by Pearson Edexcel	Mathematics
6.	The mass, m grams, of a radioactive substance t years after first being modelled by the equation	observed, is
	$m = 25 \mathrm{e}^{1-kt}$	
	where k is a positive constant.	
	(a) State the value of <i>m</i> when the radioactive substance was first observed.	(1)
	Given that the mass is 50 grams, 10 years after first being observed,	
	(b) show that $k = \frac{1}{10} \ln\left(\frac{1}{2}e\right)$	(4)
	(c) Find the value of t when $m = 20$, giving your answer to the nearest year	r. (3)

Question Number		Scheme		Marks	
6(a)	25e or equivalent dec	eimal - Accept awrt 68		B1	
(b)	$50 = 25e^{1-10k}$			M1	(1)
	Way 1 $e^{1-10k} = 2$	Way 2 $e^{-10k} = 2/e$	Way 3 $e^{10k} = e/2$	A1	
	$\Rightarrow 1-10k = \ln 2$	$-10k = \ln(2/e)$	$10k = \ln(e/2)$	M1	
	$\implies k = \frac{\ln e - \ln 2}{10}$	$k = \frac{-\ln\left(\frac{2}{e}\right)}{10}$	No intermediate step needed		
		$\Rightarrow k = \frac{\ln\left(\frac{1}{2}e\right)}{10} *$		A1*	(4)
(c)	Uses $m = 20$ and their	numerical k so $20 = 25e^{1}$	$e^{-k't} \Rightarrow e^{1-k't} = 0.8$ o.e. $\Rightarrow t = \frac{1 - \ln 0.8}{k'}$	M1 dM1	
		=	$\Rightarrow t = awrt 40 (years)$	A1	(3)
				(8 marks)	(0)

- (a) B1 for 25e or for numerical answer, e.g. 67.957 allow awrt 68
- (b) M1 Uses t = 10, m = 50, in $m = 25e^{1-kt}$ to give $50 = 25e^{1-10k}$

A1
$$e^{1-10k} = 2$$
 (way 1) or $e^{-10k} = 2/e$ o.e. (way 2) or $e^{10k} = e/2$ (way 3) or

$$e^{10k-1} = \frac{1}{2}$$
 (variant on Way 1) to give a correct equation $e^{f(k)} = B$. Some solutions will

move from

one of these options to another by sound algebra – this is acceptable.

M1 Taking logs **correctly** to give $f(k) = \ln B$ i.e. $1-10k = \ln 2$ (way 1) or $-10k = \ln(2/e)$ o.e. (way 2) or $10k = \ln(e/2)$ (way 3) or $10k - 1 = \ln(1/2)$ (way 4*)

(There are a number of correct alternatives but this line should follow directly from the previous one) This must be a correct equation.

A1* cso-Needs both M marks, everything should have been correct and exact. Makes k the subject of the Formula. Needs an intermediate step for ways 1 and 2 but not for way 3.

e.g.
$$k = \frac{\ln e - \ln 2}{10}$$
 (Way 1 or Way 4*) or $k = \frac{-\ln(2/e)}{10}$ (Way2) or straight to
 $k = \frac{\ln\left(\frac{1}{2}e\right)}{10}$ (Way 3)
 $\ln\left(\frac{1}{2}e\right) = \ln\left(\frac{e}{2}\right)$

Must conclude with the printed answer $k = \frac{\ln\left(\frac{1}{2}e\right)}{10}$ or $k = \frac{\ln\left(\frac{e}{2}\right)}{10}$ or $k = \frac{1}{10}\ln\left(\frac{e}{2}\right)$ o.e.

Special Case Taking the mass as 50 + 25e in part (b) should be treated as misread. Can earn M1A0M1A0 and obtains $k = \frac{1 - \ln(2 + e)}{10}$

(c) M1 Uses m = 20 and their numerical k in $m = 25e^{1-kt} \implies e^{1-k't} = 0.8$ (NB k = 0.030685...) NB $\implies e^{(1-k')t} = 0.8$ is M0 (usually $e^{0.97t} = 0.8$)

dM1 Use of correct work to reach
$$\Rightarrow t = \frac{1 - \ln 0.8}{k'}$$
 or equivalent e.g. $\Rightarrow t = \frac{\ln(\frac{5e}{4})}{k'}$

A1 Allow awrt 40 (may see 39.86 or 39.9). (Decimals are acceptable in part (c)) Do not allow -40 **Special Case**

If the answer 40 appears with no working or after minimal working where no marks have been scored then award M1M0A0 – special case.

If the first M mark in (c) has been awarded and they give the answer 40 with no further working, then award M1M1A1

Summer 20 Past Paper	15 www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics C34 WMA02
7. (a)	Use the substitution $t = \tan x$ to show that the equation	Leave blank
	$4\tan 2x - 3\cot x \sec^2 x = 0$	
	can be written in the form	
	$3t^4 + 8t^2 - 3 = 0$	
	5i + 6i - 5 = 0	(4)
(b)	Hence solve, for $0 \leq x < 2\pi$,	
	$4\tan 2x - 3\cot x \sec^2 x = 0$	
	Give each answer in terms of π . You must make your method clear.	(4)

Question Number	Scheme	Marks	
7(a)	$4\tan 2x - 3\cot x \sec^2 x = 0 \Longrightarrow 4 \times \frac{2t}{1 - t^2} - 3 \times \frac{1}{t} \times (1 + t^2) = 0$	<u>B1M1A1</u>	
	So $4 \times 2t^2 - 3 \times (1+t^2)(1-t^2) = 0$ and $3t^4 + 8t^2 - 3 = 0$ *	<u>A1*</u>	
(b)	$3t^4 + 8t^2 - 3 = 0 \Rightarrow (3t^2 - 1)(t^2 + 3) = 0$ so $t =$	M1	(4)
	$\tan x(t) = \pm \frac{1}{\sqrt{3}} \text{ or } \pm 0.5774$	A1	
	$x = \frac{\pi}{6}, \frac{7\pi}{6}, \frac{5\pi}{6}, \frac{11\pi}{6}$	M1A1	(4)
		(8 mai	rks)

- (a) B1 Uses $4 \tan 2x = 4 \times \frac{2t}{1-t^2}$ or $4 \tan 2x = 4 \times \frac{2 \tan x}{1-\tan^2 x}$. (B0 for $\tan 2x = \frac{\sin 2x}{\cos 2x}$)
 - M1 Uses either $\cot x = \frac{1}{\tan x}$ or $\frac{1}{t}$ or $\sec^2 x = 1 + \tan^2 x$ or $1 + t^2$ (quoted correctly)
 - A1 Uses **both** $\cot x = \frac{1}{\tan x} \text{ or } \frac{1}{t}$ **and** $\sec^2 x = 1 + \tan^2 x \text{ or } 1 + t^2$ both quoted correctly

[This M1A1 may also arise from the use of
$$\cot x = \frac{\cos x}{\sin x}$$
 with $\sec^2 x = \frac{1}{\cos^2 x}$ and with

 $\cos^2 x + \sin^2 x = 1$ to reach $\frac{1}{t} + t$; so M1 for all three of these identities and A1 for reaching result.] $\cot x = \frac{1}{\tan x}$ may be implied (by $\tan x \cot x = 1$ for example)

A1* As the answer is given, a <u>fully correct intermediate line</u> of working must be seen (Answer may have terms in a different order but must be equivalent correct equation in t)

- (b) M1 Solves given quadratic in t^2 . (may be one slip copying) Accept correct factorisation (shown), formula, completing square. Must then **also square root**. Need to get to t = This may be implied by just one answer for t or tan x (from a graphical calculator)
 - A1 $\tan x(t) = \pm \frac{1}{\sqrt{3}}$ (need both plus and minus) or *awrt* ± 0.5774 (ignore further values of tan x from their quadratic) N.B. $\tan^2 x = \frac{1}{3}$ is not enough. This is M0A0
 - M1 For obtaining two answers for x from their answers for t (must be in different quadrants if following through wrong t) (check on your calculator)
 - A1 All 4 exact and correct and no extra values in the range If all four answers are correct but in degrees 30, 210, 150, 330 lose final A mark If the answers are given as decimals 0.524, 3.67, 2.62 and 5.76 lose final A mark.

This resource was created and owned by Pearson Edexcel Past Paper WMA02 Leave blank (a) Prove by differentiation that 8. $\frac{\mathrm{d}}{\mathrm{d}y}(\ln\tan 2y) = \frac{4}{\sin 4y}, \qquad 0 < y < \frac{\pi}{4}$ (4) (b) Given that $y = \frac{\pi}{6}$ when x = 0, solve the differential equation $\frac{\mathrm{d}y}{\mathrm{d}x} = 2\cos x \sin 4y, \qquad 0 < y < \frac{\pi}{4}$ Give your answer in the form $\tan 2y = Ae^{B\sin x}$, where A and B are constants to be determined. (6) 22 P 4 4 9 7 0 A 0 2 2 4 4

Question Number	Scheme	Marks
8 (a)	$\frac{\mathrm{d}}{\mathrm{d}y}(\ln\tan 2y) = \frac{1}{\tan 2y} \times 2\sec^2 2y$	M1A1
	$=\frac{\cos 2y}{\sin 2y} \times \frac{2}{\cos^2 2y} = -\frac{k}{\sin 2y \cos 2y}$	M1
	$=\frac{4}{2\sin 2v\cos 2v} = \frac{4}{\sin 4v}$	A1* cso
(b)	dy a contract of dy factor	(4)
Way 1	$\frac{1}{dx} = 2\cos x \sin 4y \Rightarrow \int \frac{1}{\sin 4y} = \int 2\cos x dx$	B1
	$\Rightarrow \frac{1}{4} \ln \tan 2y = 2\sin x (+c)$	M1A1
Finds limits first	Put $x = 0, y = \frac{\pi}{6} \Rightarrow \frac{1}{4} \ln \tan 2\frac{\pi}{6} = 2\sin 0 + c \Rightarrow c = \dots \left(\frac{1}{4} \ln \sqrt{3} \text{ or } \frac{1}{8} \ln 3\right)$	M1
	Takes exponentials so $\tan 2y = e^{8\sin x + c}$	M1
Finds limits after removing lns	$\tan 2y = e^{8\sin x + c} (\operatorname{so} \tan 2y = Ae^{8\sin x})$	M1 (bM3 on epen)
	Put $x = 0, y = \frac{\pi}{6}$, so $A = \text{ or } e^{c} =$	M1 (bM2 on epen)
	$\tan 2y = \sqrt{3} e^{8\sin x}$	A1
		(6) (10 marks)
(b) Way 2	$\frac{dy}{dx} = 2\cos x \sin 4y \Rightarrow \int \frac{dy}{\sin 4y} = \int 2\cos x dx$	B1
	$\Rightarrow -\frac{1}{4}\ln(\csc 4y + \cot 4y) = 2\sin x(+c)$	M1A1
	Sub $x = 0, y = \frac{\pi}{6}$	
	$\Rightarrow -\frac{1}{4}\ln(\operatorname{cosec}\frac{2\pi}{3} + \cot\frac{2\pi}{3}) = 2\sin 0 + c \Rightarrow c = \dots \left(-\frac{1}{4}\ln\frac{1}{\sqrt{3}} \text{ or } \frac{1}{4}\ln\sqrt{3}\right)$	M1
	$-\frac{1}{4}\ln(\frac{1+\cos 4y}{\sin 4y}) = \frac{1}{4}\ln(\tan 2y) = 2\sin x + \frac{1}{4}\ln\sqrt{3} \text{so} \tan 2y = \sqrt{3} e^{8\sin x}$	M1A1 (6)
(b) Way 3	Special case: Differentiates the answer. Marks available B0M1A0M1M1A1	
	$\tan 2y = A e^{B \sin x} \rightarrow (2) \sec^2 2y \frac{dy}{dx} = AB \cos x e^{B \sin x}$	M1
	$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{B\cos x \tan 2y \cos^2 2y}{(2)}$	M1
	$\frac{dy}{dx} = \frac{B\cos x 2\sin 2y \cos 2y}{(4)} = \frac{B\cos x \sin 4y}{(4)}$	M1
	$B = 8$ and $A = \sqrt{3}$	A1 4/6

(a) M1 Uses chain rule to obtain
$$\frac{d}{dy}(\ln \tan 2y) = \frac{1}{\tan 2y} \times A \sec^2 2y$$
 (A can even be 1)

A1 Correct answer (unsimplified)

M1 Uses identity for
$$\tan 2y = \frac{\sin 2y}{\cos 2y}$$
 and for $\sec^2 2y = \frac{1}{\cos^2 2y}$ and reaches $\frac{k}{\sin 2y \cos 2y}$

or $k \times \frac{1}{\sin 2y} \times \frac{1}{\cos 2y}$

NB Some use long alternative methods using

$$\sec^{2} 2y = 1 + \tan^{2} 2y = 1 + \frac{\sin^{2} 2y}{\cos^{2} 2y} = \frac{\cos^{2} 2y + \sin^{2} 2y}{\cos^{2} 2y} = \frac{1}{\cos^{2} 2y}$$

There needs to be a complete method leading to $\frac{\kappa}{\sin 2y \cos 2y}$ for the M1

A1* cso Writes their expression in terms of $\sin 4y$ using the identity $\sin 4y = 2\sin 2y \cos 2y$ This is a given answer which must be stated and all aspects of the proof must be correct. (Need to multiply top and bottom of fraction by 2 or use other convincing intermediate step).

- (b) B1 Separate terms. Accept without the integral as long as integration is implied by subsequent working.

 $\int \cos x dx = \pm \sin x \text{ to produce } A \ln \tan 2y = B \sin x (+c) \text{ (or quotes results)}$

- A1 Correct answer, no need for (+c) (Correct answer implies the M1)
- M1 Subs $x = 0, y = \frac{\pi}{6}$ into their integrated expression to find c = (must have c for this mark)
- M1 Uses correct ln work to find an unsimplified expression for $\tan 2y$ (must have *c*) (They may remove lns before finding a value for *c*)

A1 Correct answer and correct solution. Do not accept
$$A = \tan\left(\frac{\pi}{3}\right)$$
 need $A = \sqrt{3}$

Way 2

A scheme is given but most will struggle to complete this method. The first 4 marks are readily accessible but showing the answer is difficult. If in doubt, send to review.

- B1 Separate terms as in first method
- M1 Use standard integral on scheme and obtain $A \ln(\csc 4y + \cot 4y) = 2 \sin x(+c)$
- A1 Correct answer, no need for (+c)

M1 Subs $x = 0, y = \frac{\pi}{6}$ into their integrated expression to find *c* (must have *c* for this mark)

M1 Uses definitions of cosec and cot together with double angle formulae to give $\tan 2y$ (must have *c*)

A1 Correct answer and conclusion with correct values for *A* and *B*

Way 3 (Special Case)

This assumes the answer and differentiates. This is not a complete answer to the question as it merely shows that the given answer satisfies the differential equation. So this is treated as a misread and four of the six marks are available.

- B0 Not available as variables are not separated
- M1 Implicit differentiation (There may be sign errors or wrong factors of 2)
- A0 Not available
- M1 Obtains expression in scheme (may be sign errors or wrong factors of 2)

M1 Uses definitions with double angle formulae to give
$$\frac{B \cos x \sin 4y}{2}$$

- (4)
- A1 Correct answer and conclusion with correct values for *A* and *B*





Question Number	Scheme	Marks
9(a)	A and B are where $y = 0$ so $t^3 - 9t = 0 \Rightarrow t(t^2 - 9) = 0 \Rightarrow t = 3$ (0 and -3)	M1
	When $t = 3$, $x = 15$	A1
	A = (3, 0)	B1
•		(3)
Or	Special case - uses answer - $t^2 + 2t = 15 \Rightarrow t = 3$ (-5)	MI
	when $t = 3$, $y = 0$	Al
	A = (3,0)	
	dv	(5)
(b)	$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\frac{\mathrm{d}y}{\mathrm{d}t}}{\frac{\mathrm{d}x}{\mathrm{d}t}} = \frac{3t^2 - 9}{2t + 2}$	M1A1
	Substitutes $t = 3$ into $\frac{dy}{dx} = \frac{3t^2 - 9}{2t + 2} \Rightarrow \text{gradient} = \left(\frac{9}{4}\right)$	M1
	Uses their $\left(\frac{9}{4}\right)$ and (15,0) to produce tangent equation $9x - 4y - 135 = 0*$	M1 A1*
		(5)
(c)	Substitutes $x = t^2 + 2t$, $y = t^2 - 9t$, into $9x - 4y - 135 = 0$	
	$\Rightarrow 9(t^2 + 2t) - 4(t^3 - 9t) - 135 = 0$	M1
	$\Rightarrow 4t^3 - 9t^2 - 54t + 135 = 0$	
	$\Rightarrow (t^2 - 6t + 9)(4t + 15) = 0 \text{or} \Rightarrow (t - 3)(t - 3)(4t + 15) = 0$	dM1
	$t = -\frac{15}{4}$	A1
	Coordinates of $X \operatorname{are}\left(\frac{105}{16}, -\frac{1215}{64}\right)$ or $\left(6\frac{9}{16}, -18\frac{63}{64}\right)$	ddM1A1cso
	Accept awrt (6.56, -18.98)	
		(5) (13 marks)

(a)	M1	Sets $y = 0$, so $t^3 - 9t = 0 \Longrightarrow t = 3$.
	A1	Uses $t = 3$, to state the x coordinate of B is 15
	B1	States $A = (3,0)$ (need not see working) must have both coordinates
S.C.	M1	Sets $t^2 + 2t = 15 \Longrightarrow t = 3$
	A1	Uses $t = 3$, to state the y coordinate of B is 0
	B1	States $A = (3,0)$ (need not see working)
(b)	M1	Differentiates $x(t)$ and $y(t)$ (allow one error) and calculates $\frac{dy}{dx}$ by using $\frac{dy}{dt}$ -
	A1	$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{3t^2 - 9}{2t + 2}$
	M1	Substitutes 'their' $t=3$ into their $\frac{dy}{dx}$ to find the gradient of the tangent at B
	M1	Uses (15,0) and their non-zero numerical gradient to find an equation of the tangent.
	A1*	Achieves given answer cso $\Rightarrow 9x - 4y - 135 = 0$
(c)	M1	Substitutes $x = t^2 + 2t$, $y = t^3 - 9t$ into $9x - 4y - 135 = 0$ correctly to form a cubic equation in just <i>t</i> . (Need a correct equation) N.B. Any attempts to work in just <i>x</i> or in just <i>y</i> are unlikely to achieve a cubic. If there seems
		to be any success send to review. Equations in a mixture of x , y and t score M0
	dM1	Attempts to solve a cubic polynomial = 0. Division by $(t-3)$ or by $(t-3)^2$ Use of a graphical calculator is acceptable. Just seeing $(4t + 15)$ is enough but any errors e.g. extra solutions or errors factorising) are penalised by the loss of the final A mark. May use other variables than t for this mark, when trying to factorise.
	A1	$t = -\frac{15}{4}$ (This answer with no working implies previous M mark if cubic has been seen)
	ddM1	Uses their value of t to find both the x and y co-ordinates. It is dependent upon both the previous M's having been scored
	Alcso	Coordinates are $\left(\frac{105}{16}, -\frac{1215}{64}\right)$ Accept awrt (6.56, -18.98). Allow for two sets of
		coordinates if (15.0) is not rejected. But lose this mark if the correct point

is rejected in favour of (15,0). Lose this mark for errors factorising cubic earlier, or for extra wrong values of t found earlier. Allow for $x = \frac{105}{16}$, $y = -\frac{1215}{64}$

$$y = -\frac{1}{64}$$

t Paper	This resource was created and owned by Pearson Edexcel	N
10.		
	x $6x$	
	Figure 4	
Figure 4 sho	ows a right circular cylindrical rod which is expanding as it is heate	d.
At time t see	conds the radius of the rod is x cm and the length of the rod is $6x$ cr	n.
Given that the of $\frac{\pi}{20}$ cm ² s ⁻¹	he cross-sectional area of the rod is increasing at a constant rate $^{-1}$, find the rate of increase of the volume of the rod when $x = 2$	
Write your a	answer in the form $k\pi$ cm ³ s ⁻¹ where k is a rational number.	(6)



www.mystudybro.com This resource was created and owned by Pearson Edexcel

Question Number	Scheme	Marks
10.	$A = \pi x^2 \Rightarrow \frac{dA}{dx} = 2\pi x$ (condone use of <i>r</i> throughout instead of <i>x</i>)	B1
	Uses $\frac{dA}{dt} = \frac{dA}{dx} \times \frac{dx}{dt} \Rightarrow \frac{\pi}{20} = 2\pi x \times \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = \frac{1}{40x} = \left(\frac{1}{80}\right)$	M1 A1
	$V = 6\pi x^3 \Longrightarrow \frac{\mathrm{d}V}{\mathrm{d}x} = 18\pi x^2$	B1
	Uses $\frac{dV}{dt} = \frac{dV}{dx} \times \frac{dx}{dt} \Rightarrow \frac{dV}{dt} = 18\pi 2^2 \times \frac{1}{80} = \frac{9}{10}\pi$	dM1 A1
	$(A)^{\frac{3}{2}}$	(6 marks)
Way 2	$V = 6\pi x^3, A = \pi x^2 \Longrightarrow V = 6\pi \left(\frac{A}{\pi}\right)^2$	M1
	$\frac{dV}{dA} = 9\left(\frac{A}{\pi}\right)^{\frac{1}{2}}$	B1 B1
	Uses $\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{\mathrm{d}V}{\mathrm{d}A} \times \frac{\mathrm{d}A}{\mathrm{d}t} \Rightarrow \frac{\mathrm{d}V}{\mathrm{d}t} = 9\left(\frac{4\pi}{\pi}\right)^{\frac{1}{2}} \times \frac{\pi}{20} \Rightarrow \frac{\mathrm{d}V}{\mathrm{d}t}\Big _{A=4\pi} = \frac{9}{10}\pi$	dM1A1 A1 (6 marks)
Way 3	$A = \pi x^2 \Rightarrow \frac{\mathrm{d}A}{\mathrm{d}x} = 2\pi x$	First B1
	$V = 6\pi x^3 \Longrightarrow \frac{\mathrm{d}V}{\mathrm{d}x} = 18\pi x^2$	Second B1
	Uses $\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{\mathrm{d}V}{\mathrm{d}x} \times \frac{\mathrm{d}x}{\mathrm{d}A} \times \frac{\mathrm{d}A}{\mathrm{d}t} \Rightarrow \frac{\mathrm{d}V}{\mathrm{d}t} = 18\pi x^2 \times \frac{1}{2\pi x} \times \frac{\pi}{20}$	M1A1
	Put $x = 2$ to give $\frac{9}{10}\pi$	dM1A1 (6 marks)
	Misunderstands area as Total Surface Area $A = 14\pi x^2 \Rightarrow \frac{dA}{dx} = 28\pi x$	B0
	Uses $\frac{dA}{dt} = \frac{dA}{dx} \times \frac{dx}{dt} \Rightarrow \frac{\pi}{20} = 28\pi x \times \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = \frac{1}{560x} = \left(\frac{1}{1120}\right)$	M1A1
	$V = 6\pi x^3 \Longrightarrow \frac{\mathrm{d}V}{\mathrm{d}x} = 18\pi x^2$	B1
	$\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{\mathrm{d}V}{\mathrm{d}x} \times \frac{\mathrm{d}x}{\mathrm{d}t} \Longrightarrow \frac{\mathrm{d}V}{\mathrm{d}t} = 18\pi 2^2 \times \frac{1}{1120} = \frac{9}{140}\pi$	dM1A0
	Misunderstands area as Curved Surface Area $A = 12\pi x^2 \Rightarrow \frac{dA}{dx} = 24\pi x$	B0
	Similar scheme to above with $\frac{\pi}{20} = 24\pi x \times \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = \frac{1}{480x} = \left(\frac{1}{960}\right)$	M1A1
	$V = 6\pi x^3 \Longrightarrow \frac{\mathrm{d}V}{\mathrm{d}x} = 18\pi x^2$	B1
	$\frac{\mathrm{d}V}{\mathrm{d}t} = \frac{\mathrm{d}V}{\mathrm{d}x} \times \frac{\mathrm{d}x}{\mathrm{d}t} \Longrightarrow \frac{\mathrm{d}V}{\mathrm{d}t} = 18\pi 2^2 \times \frac{1}{960} = \frac{9}{120}\pi \ (or \ \frac{3}{40}\pi)$	dM1A0

Must use calculus.

Way 1 B1 Correct statement
$$\frac{dA}{dx} = 2\pi x$$

M1 Uses correct chain rule $\frac{dA}{dt} = \frac{dA}{dx} \times \frac{dx}{dt}$ or equivalent e.g. $\frac{dx}{dt} = \frac{dA}{dt} \div \frac{dA}{dx}$
with $\frac{dA}{dt} = \frac{\pi}{20}$ and their $\frac{dA}{dx}$ to calculate $\frac{dx}{dt}$. NB If they correctly state the chain rule
 $\frac{dA}{dt} = \frac{dA}{dx} \times \frac{dx}{dt}$ then make an algebraic error they may be awarded this method mark
A1 Obtain correct expression for $\frac{dx}{dt}$ e.g. $= \frac{\pi/20}{2\pi x}$ then isw (award this mark in the two
misread cases described where correct Curved Surface Area or Total Surface Area are used
correctly)
B1 Correct statement $\frac{dV}{dx} = 18\pi x^2$
dM1 Uses $\frac{dV}{dt} = \frac{dV}{dx} \times \frac{dx}{dt}$ with their $\frac{dx}{dt}$ and $\frac{dV}{dx}$ to calculate $\frac{dV}{dt}$, with $x = 2$ substituted.
NB $\frac{dV}{dx}$ should be in terms of one variable x (so of the form kx^2 and not kxh)
A1 $\frac{9}{10}\pi$ or 0.9π or $\frac{18}{20}\pi$ or $k = 0.9$ etc
Way 2 M1 Writes V in terms of A . Accept $V = ...A^{\frac{3}{2}}$ (first M on epen) This indicates way 2.
B2 $\frac{dV}{dt} = 9\left(\frac{A}{\pi}\right)^{\frac{1}{2}}$ (both Bs on epen)
dM1 This is dependent on the first method mark where $V = ...A^{\frac{3}{2}}$.
Uses $\frac{dV}{dt} = \frac{dV}{dA} \times \frac{dA}{dt}$ with their $\frac{dV}{dA}$ and $\frac{dA}{dt} = \frac{\pi}{20}$ to find $\frac{dV}{dt}$ (second M on epen)
NB $\frac{dV}{dt}$ should be in terms of one variable A (so of the form $kA^{\frac{1}{2}}$)
A2 $\frac{9}{10}\pi$ or 0.9π or $\frac{18}{20}\pi$ or $k = 0.9$ etc (both As on epen)
Way 3 Similar to Way 1 but does calculation in one stage, not two – see scheme above

Condone use of r throughout, but if there is a mixture of r, x and h, then final accuracy mark may be with- held if full marks would have been gained. (Send to review if in doubt)

NB: Misreads/ misunderstandings of cross section area- there are two examples in the scheme above- another possible is an open cylinder where area = $13\pi x^2 \Rightarrow \frac{dA}{dx} = 26\pi x - \frac{dA}{dx}$

then
$$\frac{\pi}{20} = 26\pi x \times \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = \frac{1}{520x} = \left(\frac{1}{1040}\right)$$
 and then $V = 6\pi x^3 \Rightarrow \frac{dV}{dx} = 18\pi x^2$ which gives
 $\frac{dV}{dt} = \frac{dV}{dx} \times \frac{dx}{dt} \Rightarrow \frac{dV}{dt} = 18\pi 2^2 \times \frac{1}{1040} = \frac{9}{130}\pi$ These may be also combined with Way 3

This resource was created and owned by Pearson Edexcel **WMA02** Leave blank 11. (a) Express $1.5\sin\theta - 1.2\cos\theta$ in the form $R\sin(\theta - \alpha)$, where R > 0 and $0 < \alpha < \frac{\pi}{2}$ Give the value of *R* and the value of α to 3 decimal places. (3) The height, H metres, of sea water at the entrance to a harbour on a particular day, is modelled by the equation $H = 3 + 1.5 \sin\left(\frac{\pi t}{6}\right) - 1.2 \cos\left(\frac{\pi t}{6}\right), \quad 0 \leqslant t < 12$ where *t* is the number of hours after midday. (b) Using your answer to part (a), calculate the minimum value of H predicted by this model and the value of t, to 2 decimal places, when this minimum occurs. (4) (c) Find, to the nearest minute, the times when the height of sea water at the entrance to the harbour is predicted by this model to be 4 metres. (6)



Question Number	Scheme	Marks	
11(a)	$(R = \sqrt{1.5^2 + 1.2^2}) = \text{awrt } 1.921 \text{ - accept e.g. } \sqrt{3.69} \text{ or } \frac{3\sqrt{41}}{10}$	B1	
	$\tan \alpha = \frac{1.2}{1.5} \Rightarrow \alpha = 0.675 \text{ or } 0.215\pi$	M1A1	
	(πt)		(3)
(b)	$H = 3 + 1.921 \sin\left(\frac{\pi}{6} - 0.675\right)$		
	$H_{\min} = 3 - 1.921' = a \text{ wrt } 1.08$	M1A1	
	$\left(\frac{\pi t}{6} - "0.675"\right) = \frac{3\pi}{2} \Longrightarrow t = 10.29$	M1A1	
			(4)
(c)	$4 = 3 + 1.921\sin\left(\frac{\pi t}{6} - 0.675\right) \Longrightarrow \sin\left(\frac{\pi t}{6} - 0.675\right) = \frac{1}{1.921}$	M1	
	$\frac{\pi t}{6} - 0.675 = 0.548 \Longrightarrow t = \text{awrt } 2.33 \text{ or } 2.34$	dM1A1	
	$\frac{\pi t}{6} - 0.675 = \pi - 0.548 = 2.594 \Longrightarrow t = \text{awrt } 6.24 \text{ or } 6.25$	ddM1A1	
	Times are 2:20pm and 6:15pm or 6.14pm (14:20 and 18:15 or 18:14) – allow 2 hours 20minutes and 6 hours 15 or 14minutes or 140 minutes and 375 or 374 minutes	A1	
	Extra values in the range – lose final A mark.		(6)
		(13 marks	;)

(a) B1
$$R = \operatorname{awrt} 1.921(3 \operatorname{dp}) - \operatorname{allow} \text{ any equivalent e.g. } \sqrt{3.69} \text{ or } \frac{3\sqrt{41}}{10}$$

M1
$$\tan \alpha = \pm \frac{1.2}{1.5} \text{ or } \tan \alpha = \pm \frac{1.5}{1.2}$$

A1 $\alpha = \text{awrt } 0.675 \text{ (3dp) also allow } 0.215\pi \text{ (must be in radians)}$

(b) M1 States or attempts to calculate
$$3-R$$
 with their value of R

A1 $H_{\text{min}} = \text{awrt } 1.08$. Or 3 - $\frac{3\sqrt{41}}{10}$ o.e. Accept this for both marks as long as no incorrect working is seen.

M1 Attempts
$$\left(\frac{\pi t}{6} - \alpha'\right) = \frac{3\pi}{2} \Rightarrow t = \dots$$

(Putting equal to $-\frac{\pi}{2}$ is M1A0 (outside range) Putting equal to $\frac{\pi}{2}$ is M0A0 (wrong)) (Allow method mark for using -90 or 270 degrees (not 90 degrees), if alpha was in degrees earlier)

A1 t = awrt 10.29 (2dp). Accept this for both marks as long as no incorrect working is seen.

(c) M1
$$\sin\left(\frac{\pi t}{6} \pm \alpha'\right) = \frac{4-3}{R}$$
, where $\left|\frac{4-3}{R}\right| < 1$ (allow for degrees)

dM1 Dependent upon the previous M, using the correct order to find **one value** of *t* (allow consistent degrees)

NB: $\sin(1/1.921) = 0.497$ Seeing 0.497 is indication that sin instead of arcsin has been used. This indicates the wrong method and so M0.

- A1 Accept either awrt 2.33 or 2.34 or awrt 6.24 or 6.25 do not need units ignore wrong units e.g. minutes and seconds for this mark. 2 hours 20 minutes is correct here. Note that 2 minutes 20 seconds could get this mark but would lose the final A1.
- ddM1 Dependent upon the previous M, using the correct order to find a second value of t
- A1 Accept awrt 2.33 or 2.34 **and** awrt 6.24 or 6.25- ignore wrong units. So 6 minutes 14 seconds could get this mark but would lose the next.
- A1 Times are 2:20pm and 6:15pm (or 6.14pm) (14:20 and 18:15 (or 18.14)) (Need both times) allow 2 hours 20minutes and 6 hours 15 or 14minutes or 140 minutes and 375 or 374 minutes. Extra values in the range lose final A mark Allow method marks for degrees, and accuracy marks if they converted to

$$\sin(30t \pm 38.65') = \frac{4-3}{R}$$
, where $\left|\frac{4-3}{R}\right| < 1$ and continued to correct answers. Using $\sin\left(\frac{\pi t}{6} \pm 38.65'\right) = \frac{4-3}{R}$, where $\left|\frac{4-3}{R}\right| < 1$ will lose the accuracy marks.

WMA02 Leave

blank

- www.mystudybro.com This resource was created and owned by Pearson Edexcel Past Paper 12. (i) Relative to a fixed origin O, the line l_1 is given by the equation l_1 : $\mathbf{r} = \begin{pmatrix} -5\\1\\6 \end{pmatrix} + \lambda \begin{pmatrix} 2\\-3\\1 \end{pmatrix}$ where λ is a scalar parameter. The point P lies on l_1 . Given that \overrightarrow{OP} is perpendicular to l_1 , calculate the coordinates of P. (ii) Relative to a fixed origin O, the line l_2 is given by the equation
 - l_2 : $\mathbf{r} = \begin{pmatrix} 4 \\ -3 \\ 12 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ -3 \\ 4 \end{pmatrix}$ where μ is a scalar parameter.

The point A does not lie on l_2 . Given that the vector \overrightarrow{OA} is parallel to the line l_2 and $|\overrightarrow{OA}| = \sqrt{2}$ units, calculate the possible position vectors of the point A.

(5)

(5)



Question Number	Scheme	Marks
12(i)	$(\overrightarrow{OP}) = \begin{pmatrix} -5+2\lambda \\ 1-3\lambda \\ 6+1\lambda \end{pmatrix} \text{ or coordinates of } P \text{ are } (-5+2\lambda,1-3\lambda,6+\lambda)$ $\overrightarrow{OP} \cdot \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} = 0 \Longrightarrow \begin{pmatrix} -5+2\lambda \\ 1-3\lambda \\ 6+1\lambda \end{pmatrix} \cdot \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} = 0$	B1
	$\Rightarrow 2(-5+2\lambda) - 3(1-3\lambda) + 1(6+1\lambda) = 0 \Rightarrow 14\lambda = 7 \Rightarrow \lambda =$	M1
	$\lambda = \frac{1}{2}$	A1
	Substitute their $\lambda = \frac{1}{2}$ into their $\overrightarrow{OP} = \begin{pmatrix} -5+2\lambda \\ 1-3\lambda \\ 6+1\lambda \end{pmatrix}$	dM1
	<i>P</i> has coordinates $(-4, -0.5, 6.5)$	A1 (5)
(ii)	Way 1 $\overrightarrow{OA} = k \times \begin{pmatrix} 3 \\ -3 \\ 4 \end{pmatrix}$ or the coordinates of A are $(5k, -3k, 4k)$	M1
	$k\sqrt{5^2 + (-3)^2 + 4^2} = \sqrt{2}$ or $k^2(5^2 + (-3)^2 + 4^2) = 2$ o.e.	A1
	Finds at least one value for k and substitutes into \overrightarrow{OA} to give position	dM1
	As $k = (\pm)\frac{1}{5}$, <i>A</i> has possible positions $\begin{pmatrix} 1\\ -\frac{3}{5}\\ \frac{4}{5} \end{pmatrix}$ or $\begin{pmatrix} -1\\ \frac{3}{5}\\ -\frac{4}{5} \end{pmatrix}$ (any notation)	A1
	As $k = (\pm)\frac{1}{5}$, <i>A</i> has possible positions $\begin{pmatrix} 1\\ -\frac{3}{5}\\ \frac{4}{5} \end{pmatrix}$ and $\begin{pmatrix} -1\\ \frac{3}{5}\\ -\frac{4}{5} \end{pmatrix}$ (any notation)	A1 (5)
	Way 2 $\overrightarrow{OA} = \begin{pmatrix} x \\ -\frac{3}{5}x \\ \frac{4}{5}x \end{pmatrix}$ or the coordinates of A are $(x, -\frac{3}{5}x, \frac{4}{5}x)$	M1
	$x\sqrt{1^2 + (-\frac{3}{5})^2 + \frac{4^2}{5}^2} = \sqrt{2}$ or equivalent (see notes for variations)	A1
	Finds x and substitutes into \overrightarrow{OA} to give position	dM1
	As $x = (\pm)1$ so <i>A</i> has coordinates $(1, -\frac{3}{5}, \frac{4}{5})$ or $(-1, \frac{3}{5}, -\frac{4}{5})$	A1
	As $x = (\pm)1$ so A has coordinates $\left(1, -\frac{3}{5}, \frac{4}{5}\right)$ and $\left(-1, \frac{3}{5}, -\frac{4}{5}\right)$	A1 (5)
	Way 3 (This is a common approach: see next page)	(3)

Question Number	Scheme	Marks
	Way 3 Writes $\frac{5}{x} = -\frac{3}{y} = \frac{4}{z}$ This is 2 equations, 3 unknowns	M1
	Writes $(x^2 + y^2 + z^2) = 2$ This is the third equation in 3 unknowns	A1
	Eliminates two of the variables to obtain either x , y , or z and uses it to find the other values	dM1
	A has coordinates $\left(1, -\frac{3}{5}, \frac{4}{5}\right)$ or $\left(-1, \frac{3}{5}, -\frac{4}{5}\right)$	A1
	A has coordinates $\left(1, -\frac{3}{5}, \frac{4}{5}\right)$ and $\left(-1, \frac{3}{5}, -\frac{4}{5}\right)$	A1 (5)
(ii)	Way 4 (minimal working): States $\frac{\sqrt{2}}{\sqrt{50}}$ or $\frac{\sqrt{50}}{\sqrt{2}}$	M1A1
	Deduces position vectors or coordinates $i - \frac{3}{5}j + \frac{4}{5}k$ or $-i + \frac{3}{5}j - \frac{4}{5}k$	M1A1
		A1
	$1 - \frac{2}{5}J + \frac{2}{5}K$ and $-1 + \frac{2}{5}J - \frac{2}{5}K$	(5) (10 marks)

(i) B1
$$\overrightarrow{OP} = \begin{pmatrix} -5+2\lambda \\ 1-3\lambda \\ 6+1\lambda \end{pmatrix}$$
 or coordinates of *P* are $(-5+2\lambda, 1-3\lambda, 6+\lambda)$. This may be implied by

its use in the scalar product.

n the scalar product. Attempts to use (may make slip copying) their $\overline{OP} \cdot \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix} = 0$ obtaining an equation in M1

 λ and solving to give $\lambda =$

They may use any multiple of $2\mathbf{i} - 3\mathbf{j} + \mathbf{k}$ (including $\lambda (2\mathbf{i} - 3\mathbf{j} + \mathbf{k})$ or $-(2\mathbf{i} - 3\mathbf{j} + \mathbf{k})$ etc...) $\lambda = \frac{1}{2}$

A1

Substitutes their λ into their \overrightarrow{OP} . Dependent upon the previous M dM1

A1 P has coordinates (-4, -0.5, 6.5). Accept it written in coordinate or in a vector form –

either as column vector or as - 4**i** - 0.5 **j** +6.5**k.** Accept $\overrightarrow{OP} = \begin{pmatrix} -4 \\ -\frac{1}{2} \\ \frac{13}{2} \end{pmatrix}$ for example.

(ii) Way 1

M1 Writing
$$\overrightarrow{OA} = k \times \begin{pmatrix} 5 \\ -3 \\ 4 \end{pmatrix}$$
 or the coordinates of A as $(5k, -3k, 4k)$.

This mark may be implied by correct coordinates later.

A1 Correct equation in k (one variable) using 3D Pythagoras' theorem dM1 (Dependent on first M mark) Solves their equation to give at least one value for k (their one variable, even allow use of μ here) and substitutes to find at least one possible position.

A1
$$\mathbf{a} = \begin{pmatrix} 1 \\ -\frac{3}{5} \\ \frac{4}{5} \end{pmatrix}$$
 OR $\begin{pmatrix} -1 \\ \frac{3}{5} \\ -\frac{4}{5} \end{pmatrix}$. Allow coordinates $\begin{pmatrix} 1, -\frac{3}{5}, \frac{4}{5} \end{pmatrix}$ or $\begin{pmatrix} -1, \frac{3}{5}, -\frac{4}{5} \end{pmatrix}$
A1 $\mathbf{a} = \begin{pmatrix} 1 \\ -\frac{3}{5} \\ \frac{4}{5} \end{pmatrix}$ AND $\begin{pmatrix} -1 \\ \frac{3}{5} \\ -\frac{4}{5} \end{pmatrix}$. Allow coordinates $\begin{pmatrix} 1, -\frac{3}{5}, \frac{4}{5} \end{pmatrix}$ and $\begin{pmatrix} -1, \frac{3}{5}, -\frac{4}{5} \end{pmatrix}$

If there are surds in the answer but it is otherwise correct then lose the final A1

Way 2:

This is described in the scheme where x is the single variable used but could be used with

$$\left(-\frac{5}{3}y, y, -\frac{4}{3}y\right)$$
 where y is $\left(\pm\right)\frac{3}{5}$, or with $\left(\frac{5}{4}z, -\frac{3}{4}z, z\right)$ where z is $\left(\pm\right)\frac{4}{5}$ in a similar way.

Way 3: The successful approach is described above in the scheme.

Solving the equations may be lengthy, they may eliminate their variables one at a time and they may make errors.

M1: This is for these correct equations in the scheme (Two equations – three unknowns) A1: A correct further equation (third equation – three unknowns)

- M1: Solves to obtain the three variables
- A1, A1 as before

(A common variation)

Some write $(x^2 + y^2 + z^2) = 2$ together with a scalar product giving $5x - 3y + 4z = \sqrt{2}\sqrt{50}$ This is two equations in three unknowns and usually stops there. This is marked M0A0M0A0A0 unless the candidate produces a third equation and makes progress towards the answer.

Way 4: (Special Case) M1A1: Writes down $\frac{\sqrt{2}}{\sqrt{50}}$ with little or no working (or $\frac{1}{5}$ or even 5) dM1A1A1: Uses their correct fraction to find the coordinates (position vectors) – as before



Figure 5 shows a sketch of part of the curve with equation $y = 2 - \ln x$, x > 0

The finite region *R*, shown shaded in Figure 5, is bounded by the curve, the *x*-axis and the line with equation x = e.

The table below shows corresponding values of x and y for $y = 2 - \ln x$

x	e	$\frac{e+e^2}{2}$	e ²
у	1		0

(a) Complete the table giving the value of *y* to 4 decimal places.

(1)

(b) Use the trapezium rule, with all the values of y in the completed table, to obtain an estimate for the area of R, giving your answer to 3 decimal places.

(3)

(c) Use integration by parts to show that
$$\int (\ln x)^2 dx = x (\ln x)^2 - 2x \ln x + 2x + c$$
 (4)

The area *R* is rotated through 360° about the *x*-axis.

(d) Use calculus to find the exact volume of the solid generated.

Write your answer in the form $\pi e(pe + q)$, where p and q are integers to be found.

(6)



Question Number	Scheme	Marks
13 (a)	awrt 0.3799 – may be seen in the table	B1 (1)
(b)	Area = $\frac{1}{2} \times \left(\frac{e^2 - e}{2}\right) (1 + 2 \times 0.3799' + 0)$ [The +0 is not required]	B1M1
	= awrt 2.055	A1 (2)
(c)	$\int (\ln x)^2 dx = \int 1 \times (\ln x)^2 dx = x (\ln x)^2 - \int x \times \frac{2 \ln x}{dx} dx$, <u>(3)</u> M1
Way 1	$\int (-1x)^{2} dx = r(\ln x)^{2} - \int 2\ln x dx$	Δ 1
	$-x(mx)^2 - \int 2mx dx$	
	$= x(\ln x) - 2x\ln x + \int 2dx$	dM1
	$= x(\ln x)^{2} - 2x\ln x + 2x(+c)$	A1* (4)
(c) Way 2	$\int (\ln x)^2 dx = \int (\ln x) \times (\ln x) dx = \ln x (x \ln x - x) - \int \frac{1}{x} \times (x \ln x - x) dx$	M1
	$= \ln x (x \ln x - x) - \int \ln x - 1 dx$	A1
	$= \ln x (x \ln x - x) - (x \ln x - x - x)$	dM1
	$=x(\ln x)^2-2x\ln x+2x(+c)$	A1*
(c)	Use $u = \ln x$ substitution to get to	,
Way3	$\int u^2 e^u du = u^2 e^u - \int 2u e^u du$	M1
	$\int u^2 e^{u} - u^2 e^{u} - \int 2u e^{u} du$	A 1
	$= u^{2}c^{u} \qquad \begin{bmatrix} 2ue & \int 2e^{u} du \end{bmatrix}$ $= u^{2}c^{u} \qquad 2uc^{u} + 2c^{u} + k$	AI
	$-u e^{-2u^{-2ue^{-1}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}}$	dM1A1
(1)	$-\chi(\mathrm{in}\chi) = 2\chi(\mathrm{in}\chi + 2\chi(\mathrm{ic}))$	(4)
(a)	$\int \pi y dx = \int \pi (2 - \ln x) dx$	BI
	$\int (2 - \ln x)^2 dx = \int 4 - 4 \ln x + (\ln x)^2 dx$	M1
	Correct integration of at least two of their three terms (see notes)	M1
	$= 4x - 4(x \ln x - x) + x(\ln x)^2 - 2x \ln x + 2x (+c)$	A1
	$V_{2} = \left[$	ddM1
	volume = $\pi [4x - 4(x \ln x - x) + x(\ln x)^{-} - 2x \ln x + 2x]_{e}$	
	$=2\pi e^2 - 5\pi e$ $=\pi e(2e-5)$	Δ 1
	-nc(2c-3)	AI (6)
		(14 marks)

- (a) B1 awrt 0.3799 should be in the table or given as answer not just appear in trapezium rule
- (b) B1 For the strip width of $\frac{e^2 e}{2}$ or correct equivalent e.g. $\frac{e^2 + e}{2} e$, or $e^2 \frac{e^2 + e}{2}$ Also accept awrt 2.34 o.e. This may be stated as h = the values above, or may be used correctly in the rule so $\frac{e^2 - e}{4}$, etc or 1.17 may be seen. M1 For **correct application** of the trapezium rule – requires correct ft bracket×(1+2× their answer to(a)+0)
 - A1 awrt 2.055

(c) Must use integration by parts or 0/4. Differentiating the answer is not acceptable.

Accept
$$\int (\ln x)^2 dx = x(\ln x)^2 - \int x \times \frac{A \ln x}{x} dx$$

A1
$$x(\ln x)^2 - \int 2\ln x dx$$

dM1 A second application of integration by parts the correct way around Accept = $x(\ln x)^2 - Ax \ln x \pm \int B dx$

Accept the answer to the $\int \ln x dx$ part just written down as $x \ln x - x$

- A1* Correct solution only = $x(\ln x)^2 2x \ln x + 2x(+c)$ with or without 'c'
- Way 2 M1 One correct application of integration by parts the correct way around. Accept $\int (\ln x)^2 dx = \int (\ln x) \times (\ln x) dx = \ln x (x \ln x - x) - \int \frac{1}{x} \times (x \ln x - x) dx$

A1
$$\ln x(x\ln x - x) - \int (\ln x - 1) dx$$

dM1 A second application of integration by parts the correct way around Accept = $\ln x (x \ln x - x) - (x \ln x - x - x)$ Accept the answer to the $\int \ln x dx$ part just written down as $x \ln x - x$ and substituted into their expression

A1* Correct solution only =
$$x(\ln x)^2 - 2x\ln x + 2x(+c)$$
 with or without 'c'

Way 3 M1 Uses substitution and performs one correct application of integration by parts the correct way around.

A1
$$= u^2 e^u - \left[2ue^u - \int 2e^u du\right]$$
 after second integration by parts
dM1 Final integration and returns to x

A1* Correct solution only $= x(\ln x)^2 - 2x\ln x + 2x(+c)$ with or without 'c'

(d) B1 Volume = $\int \pi y^2 dx = \int \pi (2 - \ln x)^2 dx$ (needs π and integral symbol) but not limits and can condone missing dx

M1 Multiplies out $(2 - \ln x)^2$ to $\lambda + \mu \ln x + \nu (\ln x)^2$ where λ , μ and ν are non zero positive or negative constants.

M1 Needs attempt to multiply out to at least $\lambda + \nu (\ln x)^2$ where λ and ν are non zero positive or negative constants and attempt to integrate. Look for $\lambda x \pm \mu (x \ln x - x) \pm \nu (x (\ln x)^2 - 2x \ln x + 2x)$ with **two of the three** terms integrated correctly (So if $\mu = 0$ could score M0M1 here)

- A1 Correct answer $4x 4(x \ln x x) + x(\ln x)^2 2x \ln x + 2x(+c)$ with or without c. Accept unsimplified and isw.
- ddM1 Attempts to substitute both correct limits into the result of their integral. Both previous M's must have been scored. If there has been small slips simplifying the result of the integral before use of limits then allow M1
- A1 Correct solution only $\pi e(2e-5)$

Special case: For those who misunderstand/misread and think that $y = (\ln x)^2$ so

Volume = $\int \pi y^2 dx = \int \pi (\ln x)^4 dx$ - the first B1 may be exceptionally awarded. If anyone appears to make progress with a method for this integration (not rubbish) please send to review.

This gains 1/6 marks