

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	5	/	0	1	Signature	

Paper Reference(s)

6665/01

Edexcel GCE

Core Mathematics C3

Advanced

Wednesday 20 January 2010 – Afternoon
Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink or Green)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 9 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with
Edexcel Limited copyright policy.
©2010 Edexcel Limited

Printer's Log. No. _____

Printer's Log. No.
N35381A

W850/R6665/57570 5/5/5/4/3/3



Turn over

edexcel 
advancing learning, changing lives

Leave
blank

$$\frac{x+1}{3x^2-3} - \frac{1}{3x+1}$$

as a single fraction in its simplest form.

(4)



January 2010
6665 Core Mathematics C3
Mark Scheme

Question Number	Scheme	Marks
Q1	$\frac{x+1}{3x^2-3} - \frac{1}{3x+1}$ $= \frac{x+1}{3(x^2-1)} - \frac{1}{3x+1}$ $= \frac{x+1}{3(x+1)(x-1)} - \frac{1}{3x+1}$ $= \frac{1}{3(x-1)} - \frac{1}{3x+1}$ $= \frac{3x+1-3(x-1)}{3(x-1)(3x+1)}$ <p>or $\frac{3x+1}{3(x-1)(3x+1)} - \frac{3(x-1)}{3(x-1)(3x+1)}$</p> $= \frac{4}{3(x-1)(3x+1)}$ <p> <i>Decide to award M1 here!!</i> Either $\frac{4}{3(x-1)(3x+1)}$ or $\frac{\frac{4}{3}}{(x-1)(3x+1)}$ or $\frac{4}{(3x-3)(3x+1)}$ or $\frac{4}{9x^2-6x-3}$ </p>	<p>Award below</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 aef</p> <p>[4]</p>

2.

$$f(x) = x^3 + 2x^2 - 3x - 11$$

(a) Show that $f(x) = 0$ can be rearranged as

$$x = \sqrt{\left(\frac{3x+11}{x+2}\right)}, \quad x \neq -2.$$

(2)

The equation $f(x) = 0$ has one positive root α .

The iterative formula $x_{n+1} = \sqrt{\left(\frac{3x_n + 11}{x_n + 2}\right)}$ is used to find an approximation to α .

(b) Taking $x_1 = 0$, find, to 3 decimal places, the values of x_2 , x_3 and x_4 .

(3)

(c) Show that $\alpha = 2.057$ correct to 3 decimal places.

(3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
Q2	$f(x) = x^3 + 2x^2 - 3x - 11$	
(a)	<p>Sets $f(x) = 0$ (can be implied) and takes out a factor of x^2 from $x^3 + 2x^2$, or x from $x^3 + 2x$ (slip).</p> $f(x) = 0 \Rightarrow x^3 + 2x^2 - 3x - 11 = 0$ $\Rightarrow x^2(x + 2) - 3x - 11 = 0$ $\Rightarrow x^2(x + 2) = 3x + 11$ $\Rightarrow x^2 = \frac{3x + 11}{x + 2}$ $\Rightarrow x = \sqrt{\left(\frac{3x + 11}{x + 2}\right)}$ <p>then rearranges to give the quoted result on the question paper.</p>	M1 A1 AG (2)
(b)	<p>Iterative formula: $x_{n+1} = \sqrt{\left(\frac{3x_n + 11}{x_n + 2}\right)}$, $x_1 = 0$</p> $x_2 = \sqrt{\left(\frac{3(0) + 11}{(0) + 2}\right)}$ <p>An attempt to substitute $x_1 = 0$ into the iterative formula. Can be implied by $x_2 = \sqrt{5.5}$ or 2.35 or awrt 2.345</p> <p>Both $x_2 =$ awrt 2.345 and $x_3 =$ awrt 2.037 $x_4 =$ awrt 2.059</p>	M1 A1 A1 (3)
(c)	<p>Let $f(x) = x^3 + 2x^2 - 3x - 11 = 0$</p> <p>$f(2.0565) = -0.013781637...$ $f(2.0575) = 0.0041401094...$ Sign change (and $f(x)$ is continuous) therefore a root α is such that $\alpha \in (2.0565, 2.0575) \Rightarrow \alpha = 2.057$ (3 dp)</p> <div style="border: 1px solid black; padding: 5px; margin: 5px;">Choose suitable interval for x, e.g. [2.0565, 2.0575] or tighter</div> <div style="border: 1px solid black; padding: 5px; margin: 5px;">any one value awrt 1 sf</div> <div style="border: 1px solid black; padding: 5px; margin: 5px;">both values correct awrt 1sf, sign change and conclusion</div> <div style="border: 1px solid black; padding: 5px; margin: 5px;">As a minimum, both values must be correct to 1 sf, candidate states "change of sign, hence root".</div>	M1 dM1 A1 (3)
		[8]

3. (a) Express $5 \cos x - 3 \sin x$ in the form $R \cos(x + \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$. (4)

for $0 \leq x < 2\pi$, giving your answers to 2 decimal places. (5)

[illegible]

Question Number	Scheme	Marks
Q3 (a)	$5\cos x - 3\sin x = R\cos(x + \alpha), \quad R > 0, \quad 0 < x < \frac{\pi}{2}$ $5\cos x - 3\sin x = R\cos x\cos\alpha - R\sin x\sin\alpha$ <p>Equate $\cos x$: $5 = R\cos\alpha$ Equate $\sin x$: $3 = R\sin\alpha$</p> $R = \sqrt{5^2 + 3^2}; = \sqrt{34} \quad \{= 5.83095..\}$ $\tan\alpha = \frac{3}{5} \Rightarrow \alpha = 0.5404195003...^{\circ}$ <p>Hence, $5\cos x - 3\sin x = \sqrt{34}\cos(x + 0.5404)$</p>	<p>M1; A1</p> <p>M1 A1</p>
(b)	$5\cos x - 3\sin x = 4$ $\sqrt{34}\cos(x + 0.5404) = 4$ $\cos(x + 0.5404) = \frac{4}{\sqrt{34}} \quad \{= 0.68599...\}$ $(x + 0.5404) = 0.814826916...^{\circ}$ $x = 0.2744...^{\circ}$ $(x + 0.5404) = 2\pi - 0.814826916...^{\circ} \quad \{= 5.468358...^{\circ}\}$ $x = 4.9279...^{\circ}$ <p>Hence, $x = \{0.27, 4.93\}$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>ddM1 A1</p> <p>(4)</p> <p>(5)</p> <p>[9]</p>

Part (b): If there are any EXTRA solutions inside the range $0 \leq x < 2\pi$, then withhold the final accuracy mark if the candidate would otherwise score all 5 marks. Also ignore EXTRA solutions outside the range $0 \leq x < 2\pi$.

Leave
blank

4. (i) Given that $y = \frac{\ln(x^2 + 1)}{x}$, find $\frac{dy}{dx}$.

(4)

(ii) Given that $x = \tan y$, show that $\frac{dy}{dx} = \frac{1}{1 + x^2}$.

(5)




Question Number	Scheme	Marks
Q4 (i)	$y = \frac{\ln(x^2 + 1)}{x}$ $u = \ln(x^2 + 1) \Rightarrow \frac{du}{dx} = \frac{2x}{x^2 + 1}$ <p>Apply quotient rule: $\left\{ \begin{array}{l} u = \ln(x^2 + 1) \quad v = x \\ \frac{du}{dx} = \frac{2x}{x^2 + 1} \quad \frac{dv}{dx} = 1 \end{array} \right\}$</p> $\frac{dy}{dx} = \frac{\left(\frac{2x}{x^2 + 1} \right)(x) - \ln(x^2 + 1)}{x^2}$ $\left\{ \frac{dy}{dx} = \frac{2}{(x^2 + 1)} - \frac{1}{x^2} \ln(x^2 + 1) \right\}$	<p>$\ln(x^2 + 1) \rightarrow \frac{\text{something}}{x^2 + 1}$ M1</p> <p>$\ln(x^2 + 1) \rightarrow \frac{2x}{x^2 + 1}$ A1</p> <p>Applying $\frac{xu' - \ln(x^2 + 1)v'}{x^2}$ correctly. M1</p> <p>Correct differentiation with correct bracketing but allow recovery. A1</p> <p>{Ignore subsequent working.}</p> <p>(4)</p>
(ii)	$x = \tan y$ $\frac{dx}{dy} = \sec^2 y$ $\frac{dy}{dx} = \frac{1}{\sec^2 y} \{ = \cos^2 y \}$ $\frac{dy}{dx} = \frac{1}{1 + \tan^2 y}$ <p>Hence, $\frac{dy}{dx} = \frac{1}{1 + x^2}$, (as required)</p>	<p>$\tan y \rightarrow \sec^2 y$ or an attempt to differentiate $\frac{\sin y}{\cos y}$ using either the quotient rule or product rule. M1*</p> <p>$\frac{dx}{dy} = \sec^2 y$ A1</p> <p>Finding $\frac{dy}{dx}$ by reciprocating $\frac{dx}{dy}$. dM1*</p> <p>For writing down or applying the identity $\sec^2 y = 1 + \tan^2 y$, which must be applied/stated completely in y. dM1*</p> <p>For the correct proof, leading on from the previous line of working. A1 AG</p> <p>(5)</p> <p>[9]</p>

Leave
blank

5. Sketch the graph of $y = \ln|x|$, stating the coordinates of any points of intersection with the axes.

(3)

Question Number	Scheme	Marks
Q5	<div><div><div>$y = \ln x$</div><div></div></div><div><div>Right-hand branch in quadrants 4 and 1. Correct shape.</div><div>Left-hand branch in quadrants 2 and 3. Correct shape.</div><div>Completely correct sketch and both $(-1, \{0\})$ and $(1, \{0\})$</div></div></div> <div><div>B1</div><div>B1</div><div>B1</div><div>(3)</div><div>[3]</div></div>	

6.


**Figure 1**

Figure 1 shows a sketch of the graph of $y = f(x)$.






The graph intersects the y -axis at the point $(0, 1)$ and the point $A(2, 3)$ is the maximum turning point.

Sketch, on separate axes, the graphs of

- (i) $y = f(-x) + 1$,
- (ii) $y = f(x + 2) + 3$,
- (iii) $y = 2f(2x)$.

On each sketch, show the coordinates of the point at which your graph intersects the y -axis and the coordinates of the point to which A is transformed.

(9)

Question Number	Scheme	Marks
Q6 (i)	<p>$y = f(-x) + 1$</p>  <p>Shape of </p> <p>and must have a maximum in quadrant 2 and a minimum in quadrant 1 or on the positive y-axis.</p> <p>Either $(\{0\}, 2)$ or $A'(-2, 4)$</p> <p>Both $(\{0\}, 2)$ and $A'(-2, 4)$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(3)</p>
(ii)	<p>$y = f(x + 2) + 3$</p>  <p>Any translation of the original curve.</p> <p>The translated maximum has either x-coordinate of 0 (can be implied) or y-coordinate of 6.</p> <p>The translated curve has maximum $(\{0\}, 6)$ and is in the correct position on the Cartesian axes.</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(3)</p>
(iii)	<p>$y = 2f(2x)$</p>  <p>Shape of </p> <p>with a minimum in quadrant 2 and a maximum in quadrant 1.</p> <p>Either $(\{0\}, 2)$ or $A'(1, 6)$</p> <p>Both $(\{0\}, 2)$ and $A'(1, 6)$</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(3)</p> <p>[9]</p>

7. (a) By writing $\sec x$ as $\frac{1}{\cos x}$, show that $\frac{d(\sec x)}{dx} = \sec x \tan x$.

Given that $y = e^{2x} \sec 3x$,

(b) find $\frac{dy}{dx}$.

The curve with equation $y = e^{2x} \sec 3x$, $-\frac{\pi}{6} < x < \frac{\pi}{6}$, has a minimum turning point at (a, b) .

(c) Find the values of the constants a and b , giving your answers to 3 significant figures.

This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire surface, typical of notebook or legal stationery. The paper is otherwise completely empty, with no margins, text, or other markings.



Question Number	Scheme	Marks
Q7 (a)	$y = \sec x = \frac{1}{\cos x} = (\cos x)^{-1}$	
	$\frac{dy}{dx} = -1(\cos x)^{-2}(-\sin x)$	M1
	$\frac{dy}{dx} = \pm((\cos x)^{-2}(\sin x))$ $-1(\cos x)^{-2}(-\sin x)$ or $(\cos x)^{-2}(\sin x)$	A1
	$\frac{dy}{dx} = \left\{ \frac{\sin x}{\cos^2 x} \right\} = \left(\frac{1}{\cos x} \right) \left(\frac{\sin x}{\cos x} \right) = \underline{\underline{\sec x \tan x}}$	Convincing proof. Must see both <u>underlined steps</u> . A1 AG
(b)	$y = e^{2x} \sec 3x$	
	$\left\{ \begin{array}{l} u = e^{2x} \quad v = \sec 3x \\ \frac{du}{dx} = 2e^{2x} \quad \frac{dv}{dx} = 3 \sec 3x \tan 3x \end{array} \right\}$	M1
	<div style="border: 1px solid black; padding: 5px; display: inline-block;">Seen or implied</div>	Either $e^{2x} \rightarrow 2e^{2x}$ or $\sec 3x \rightarrow 3 \sec 3x \tan 3x$ Both $e^{2x} \rightarrow 2e^{2x}$ and $\sec 3x \rightarrow 3 \sec 3x \tan 3x$ A1
	$\frac{dy}{dx} = 2e^{2x} \sec 3x + 3e^{2x} \sec 3x \tan 3x$	Applies $vu' + uv'$ correctly for their u, u', v, v' M1 $2e^{2x} \sec 3x + 3e^{2x} \sec 3x \tan 3x$ A1 isw
(c)	Turning point $\Rightarrow \frac{dy}{dx} = 0$	
	Hence, $e^{2x} \sec 3x (2 + 3 \tan 3x) = 0$	Sets their $\frac{dy}{dx} = 0$ and factorises (or cancels) out at least e^{2x} from at least two terms. M1
	{Note $e^{2x} \neq 0$, $\sec 3x \neq 0$, so $2 + 3 \tan 3x = 0$, }	
	giving $\tan 3x = -\frac{2}{3}$	$\tan 3x = \pm k$; $k \neq 0$ M1
	$\Rightarrow 3x = -0.58800 \Rightarrow x = \{a\} = -0.19600\dots$	Either awrt -0.196° or awrt -11.2° A1
	Hence, $y = \{b\} = e^{2(-0.196)} \sec(3 \times -0.196)$	
	$= 0.812093\dots = 0.812$ (3sf)	0.812 A1 cao
		(4)

[11]

Part (c): If there are any EXTRA solutions for x (or a) inside the range $-\frac{\pi}{6} < x < \frac{\pi}{6}$, ie. $-0.524 < x < 0.524$ or ANY EXTRA solutions for y (or b), (for these values of x) then withhold the final accuracy mark.
Also ignore EXTRA solutions outside the range $-\frac{\pi}{6} < x < \frac{\pi}{6}$, ie. $-0.524 < x < 0.524$.

Leave
blank

8. Solve

$$\operatorname{cosec}^2 2x - \cot 2x = 1$$

for $0 \leq x \leq 180^\circ$.

(7)



Question Number	Scheme	Marks
Q8	<p>$\operatorname{cosec}^2 2x - \cot 2x = 1, \text{ (eqn *) } 0 \leq x \leq 180^\circ$</p> <p>Using $\operatorname{cosec}^2 2x = 1 + \cot^2 2x$ gives</p> <p>$1 + \cot^2 2x - \cot 2x = 1$</p> <p>$\cot^2 2x - \cot 2x = 0 \quad \text{or} \quad \cot^2 2x = \cot 2x$</p> <p>$\cot 2x(\cot 2x - 1) = 0 \quad \text{or} \quad \cot 2x = 1$</p> <p>$\cot 2x = 0 \quad \text{or} \quad \cot 2x = 1$</p> <p>$\cot 2x = 0 \Rightarrow (\tan 2x \rightarrow \infty) \Rightarrow 2x = 90, 270$</p> <p>$\Rightarrow x = 45, 135$</p> <p>$\cot 2x = 1 \Rightarrow \tan 2x = 1 \Rightarrow 2x = 45, 225$</p> <p>$\Rightarrow x = 22.5, 112.5$</p> <p>Overall, $x = \{22.5, 45, 112.5, 135\}$</p>	<p>M1</p> <p>A1</p> <p>dM1</p> <p>A1</p> <p>ddM1</p> <p>A1</p> <p>B1</p> <p>[7]</p>

If there are any EXTRA solutions inside the range $0 \leq x \leq 180^\circ$ and the candidate would otherwise score FULL MARKS then withhold the final accuracy mark (the sixth mark in this question). Also ignore EXTRA solutions outside the range $0 \leq x \leq 180^\circ$.

Leave
blank

- (3)

(5)

- (4)

- (3)

[illegible]

Question Number	Scheme	Marks
Q9 (i)(a)	$\ln(3x - 7) = 5$ $e^{\ln(3x - 7)} = e^5$ $3x - 7 = e^5 \Rightarrow x = \frac{e^5 + 7}{3} \{= 51.804...\}$	<p>Takes e of both sides of the equation. This can be implied by $3x - 7 = e^5$. M1</p> <p>Then rearranges to make x the subject. dM1</p> <p>Exact answer of $\frac{e^5 + 7}{3}$. A1</p> <p>(3)</p>
(b)	$3^x e^{7x+2} = 15$ $\ln(3^x e^{7x+2}) = \ln 15$ $\ln 3^x + \ln e^{7x+2} = \ln 15$ $x \ln 3 + 7x + 2 = \ln 15$ $x(\ln 3 + 7) = -2 + \ln 15$ $x = \frac{-2 + \ln 15}{7 + \ln 3} \{= 0.0874...\}$	<p>Takes ln (or logs) of both sides of the equation. M1</p> <p>Applies the addition law of logarithms. M1</p> <p>$x \ln 3 + 7x + 2 = \ln 15$ A1 oe</p> <p>Factorising out at least two x terms on one side and collecting number terms on the other side. ddM1</p> <p>Exact answer of $\frac{-2 + \ln 15}{7 + \ln 3}$ A1 oe</p> <p>(5)</p>
(ii) (a)	$f(x) = e^{2x} + 3, x \in \mathbb{R}$ $y = e^{2x} + 3 \Rightarrow y - 3 = e^{2x}$ $\Rightarrow \ln(y - 3) = 2x$ $\Rightarrow \frac{1}{2} \ln(y - 3) = x$ Hence $f^{-1}(x) = \frac{1}{2} \ln(x - 3)$ $f^{-1}(x)$: Domain: $x > 3$ or $(3, \infty)$	<p>Attempt to make x (or swapped y) the subject M1</p> <p>Makes e^{2x} the subject and takes ln of both sides M1</p> <p>$\frac{1}{2} \ln(x - 3)$ or $\ln \sqrt{x - 3}$ A1 cao</p> <p>or $f^{-1}(y) = \frac{1}{2} \ln(y - 3)$ (see appendix)</p> <p>Either $x > 3$ or $(3, \infty)$ or Domain > 3. B1</p> <p>(4)</p>
(b)	$g(x) = \ln(x - 1), x \in \mathbb{R}, x > 1$ $fg(x) = e^{2 \ln(x-1)} + 3 \{= (x - 1)^2 + 3\}$ $fg(x)$: Range: $y > 3$ or $(3, \infty)$	<p>An attempt to put function g into function f. M1</p> <p>$e^{2 \ln(x-1)} + 3$ or $(x - 1)^2 + 3$ or $x^2 - 2x + 4$. A1 isw</p> <p>Either $y > 3$ or $(3, \infty)$ or Range > 3 or $fg(x) > 3$. B1</p> <p>(3)</p>
		[15]