

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	5	/	0	1	Signature	

Paper Reference(s)

6665/01

Edexcel GCE

Core Mathematics C3

Advanced

Monday 24 January 2011 – Morning
Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2011 Edexcel Limited.

Printer's Log. No.

Printer's Log. No.
H35404RA

W850/R6665/57570 5/5/5/3/4



Turn over

edexcel 
advancing learning, changing lives

Leave
blank

- (3)

- (1)**

- (5)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

January 2011
Core Mathematics C3 6665
Mark Scheme

Question Number	Scheme	Marks
1.		
(a)	$7\cos x - 24\sin x = R\cos(x + \alpha)$ $7\cos x - 24\sin x = R\cos x \cos \alpha - R\sin x \sin \alpha$ Equate $\cos x$: $7 = R\cos \alpha$ Equate $\sin x$: $24 = R\sin \alpha$ $R = \sqrt{7^2 + 24^2} = 25$ $\tan \alpha = \frac{24}{7} \Rightarrow \alpha = 1.287002218...^{\circ}$ Hence, $7\cos x - 24\sin x = 25\cos(x + 1.287)$	B1 M1 A1 (3)
(b)	Minimum value = <u>-25</u> -25 or $-R$	B1ft (1)
(c)	$7\cos x - 24\sin x = 10$ $25\cos(x + 1.287) = 10$ $\cos(x + 1.287) = \frac{10}{25}$ $PV = 1.159279481...^{\circ}$ or $66.42182152...^{\circ}$ So, $x + 1.287 = \{1.159279...^{\circ}, 5.123906...^{\circ}, 7.442465...^{\circ}\}$ gives, $x = \{3.836906..., 6.155465...\}$	$\cos(x \pm \text{their } \alpha) = \frac{10}{(\text{their } R)}$ M1 For applying $\cos^{-1}\left(\frac{10}{(\text{their } R)}\right)$ M1 either $2\pi +$ or $-$ their PV° or $360^{\circ} +$ or $-$ their PV° M1 awrt 3.84 OR 6.16 A1 awrt 3.84 AND 6.16 A1 (5) [9]

2. (a) Express

$$\frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)}$$

as a single fraction in its simplest form.

(4)

Given that

$$f(x) = \frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)} - 2, \quad x > 1,$$

(b) show that

$$f(x) = \frac{3}{2x-1}$$

(2)

(c) Hence differentiate $f(x)$ and find $f'(2)$.

(3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
2.		
(a)	$\frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)}$ $= \frac{(4x-1)(2x-1) - 3}{2(x-1)(2x-1)}$ $= \frac{8x^2 - 6x - 2}{\{2(x-1)(2x-1)\}}$ $= \frac{2(x-1)(4x+1)}{\{2(x-1)(2x-1)\}}$ $= \frac{4x+1}{2x-1}$	<p>An attempt to form a single fraction</p> <p>Simplifies to give a correct quadratic numerator over a correct quadratic denominator</p> <p>An attempt to factorise a 3 term quadratic numerator</p> <p>M1</p> <p>A1 aef</p> <p>M1</p> <p>A1</p> <p>(4)</p>
(b)	$f(x) = \frac{4x-1}{2(x-1)} - \frac{3}{2(x-1)(2x-1)} - 2, \quad x > 1$ $f(x) = \frac{(4x+1)}{(2x-1)} - 2$ $= \frac{(4x+1) - 2(2x-1)}{(2x-1)}$ $= \frac{4x+1-4x+2}{(2x-1)}$ $= \frac{3}{(2x-1)}$	<p>An attempt to form a single fraction</p> <p>Correct result</p> <p>M1</p> <p>A1 *</p> <p>(2)</p>
(c)	$f(x) = \frac{3}{(2x-1)} = 3(2x-1)^{-1}$ $f'(x) = 3(-1)(2x-1)^{-2}(2)$ $f'(2) = \frac{-6}{9} = -\frac{2}{3}$	<p>$\pm k(2x-1)^{-2}$</p> <p>Either $\frac{-6}{9}$ or $-\frac{2}{3}$</p> <p>M1</p> <p>A1 aef</p> <p>A1</p> <p>(3)</p> <p>[9]</p>

Leave
blank

3. Find all the solutions of

$$2 \cos 2\theta = 1 - 2 \sin \theta$$

in the interval $0 \leq \theta < 360^\circ$.

(6)



Question Number	Scheme	Marks
3.	$2 \cos 2\theta = 1 - 2 \sin \theta$ $2(1 - 2 \sin^2 \theta) = 1 - 2 \sin \theta$ $2 - 4 \sin^2 \theta = 1 - 2 \sin \theta$ $4 \sin^2 \theta - 2 \sin \theta - 1 = 0$ $\sin \theta = \frac{2 \pm \sqrt{4 - 4(4)(-1)}}{8}$ PVs: $\alpha_1 = 54^\circ$ or $\alpha_2 = -18^\circ$ $\theta = \{54, 126, 198, 342\}$	<p>Substitutes either $1 - 2 \sin^2 \theta$ or $2 \cos^2 \theta - 1$ or $\cos^2 \theta - \sin^2 \theta$ for $\cos 2\theta$. M1</p> <p>Forms a “quadratic in sine” = 0 M1(*)</p> <p>Applies the quadratic formula See notes for alternative methods. M1</p> <p>Any one correct answer A1 180-their pv dM1(*) All four solutions correct. A1</p> <p>[6]</p>

Leave
blank



Question Number	Scheme	Marks
4.		
(a)	$\theta = 20 + Ae^{-kt}$ (eqn *) $\{t = 0, \theta = 90 \Rightarrow\} \quad 90 = 20 + Ae^{-k(0)} \quad \text{Substitutes } t = 0 \text{ and } \theta = 90 \text{ into eqn } *$ $90 = 20 + A \Rightarrow \underline{A = 70} \quad \underline{A = 70}$	M1 A1 (2)
(b)	$\theta = 20 + 70e^{-kt}$ $\{t = 5, \theta = 55 \Rightarrow\} \quad 55 = 20 + 70e^{-k(5)} \quad \text{Substitutes } t = 5 \text{ and } \theta = 55 \text{ into eqn } *$ $\frac{35}{70} = e^{-5k} \quad \text{and rearranges eqn } * \text{ to make } e^{\pm 5k} \text{ the subject.}$ $\ln\left(\frac{35}{70}\right) = -5k \quad \text{Takes 'lns' and proceeds to make '}\pm 5k\text{' the subject.}$ $-5k = \ln\left(\frac{1}{2}\right)$ $-5k = \ln 1 - \ln 2 \Rightarrow -5k = -\ln 2 \Rightarrow \underline{k = \frac{1}{5}\ln 2} \quad \text{Convincing proof that } k = \frac{1}{5}\ln 2$	M1 dM1 A1 * (3)
(c)	$\theta = 20 + 70e^{-\frac{1}{5}t\ln 2}$ $\frac{d\theta}{dt} = -\frac{1}{5}\ln 2 \cdot (70)e^{-\frac{1}{5}t\ln 2} \quad \pm \alpha e^{-kt} \text{ where } k = \frac{1}{5}\ln 2$ $-14\ln 2 e^{-\frac{1}{5}t\ln 2}$ When $t = 10$, $\frac{d\theta}{dt} = -14\ln 2 e^{-2\ln 2}$ $\frac{d\theta}{dt} = -\frac{7}{2}\ln 2 = -2.426015132...$ Rate of decrease of $\theta = 2.426^\circ\text{C/min}$ (3 dp.) awrt ± 2.426	M1 A1 oe A1 (3) [8]

5.




Figure 1

Figure 1 shows a sketch of part of the curve with equation $y = f(x)$, where

$$f(x) = (8-x)\ln x, \quad x > 0$$

The curve cuts the x -axis at the points A and B and has a maximum turning point at Q , as shown in Figure 1.

(a) Write down the coordinates of A and the coordinates of B .

(2)

(b) Find $f'(x)$.

(3)

(c) Show that the x -coordinate of Q lies between 3.5 and 3.6

(2)

(d) Show that the x -coordinate of Q is the solution of

$$x = \frac{8}{1 + \ln x}$$

(3)

To find an approximation for the x -coordinate of Q , the iteration formula

$$x_{n+1} = \frac{8}{1 + \ln x_n}$$

is used.

(e) Taking $x_0 = 3.55$, find the values of x_1 , x_2 and x_3 .

Give your answers to 3 decimal places.

(3)



Question Number	Scheme	Marks
5.		
(a)	<p>Crosses x-axis $\Rightarrow f(x) = 0 \Rightarrow (8 - x)\ln x = 0$</p> <p>Either $(8 - x) = 0$ or $\ln x = 0 \Rightarrow x = 8, 1$</p> <p>Coordinates are $A(1, 0)$ and $B(8, 0)$.</p>	<p>Either one of $\{x\}=1$ OR $x=\{8\}$ B1</p> <p>Both $A(1, \{0\})$ and $B(8, \{0\})$ B1</p> <p>(2)</p>
(b)	<p>Apply product rule: $\left\{ \begin{array}{l} u = (8 - x) \quad v = \ln x \\ \frac{du}{dx} = -1 \quad \frac{dv}{dx} = \frac{1}{x} \end{array} \right\}$</p> <p>$f'(x) = -\ln x + \frac{8-x}{x}$</p>	<p>$vu' + uv'$ M1</p> <p>Any one term correct A1</p> <p>Both terms correct A1</p> <p>(3)</p>
(c)	<p>$f'(3.5) = 0.032951317...$</p> <p>$f'(3.6) = -0.058711623...$</p> <p>Sign change (and as $f'(x)$ is continuous) therefore the x-coordinate of Q lies between 3.5 and 3.6.</p>	<p>Attempts to evaluate both $f'(3.5)$ and $f'(3.6)$ M1</p> <p>both values correct to at least 1 sf, sign change and conclusion A1</p> <p>(2)</p>
(d)	<p>At Q, $f'(x) = 0 \Rightarrow -\ln x + \frac{8-x}{x} = 0$</p> <p>$\Rightarrow -\ln x + \frac{8}{x} - 1 = 0$</p> <p>$\Rightarrow \frac{8}{x} = \ln x + 1 \Rightarrow 8 = x(\ln x + 1)$</p> <p>$\Rightarrow x = \frac{8}{\ln x + 1}$ (as required)</p>	<p>Setting $f'(x) = 0$. M1</p> <p>Splitting up the numerator and proceeding to $x=$ M1</p> <p>For correct proof. No errors seen in working. A1</p> <p>(3)</p>

Question Number	Scheme	Marks
(e)	<p>Iterative formula: $x_{n+1} = \frac{8}{\ln x_n + 1}$</p> <p>$x_1 = \frac{8}{\ln(3.55) + 1}$</p> <p>$x_1 = 3.528974374...$</p> <p>$x_2 = 3.538246011...$</p> <p>$x_3 = 3.534144722...$</p> <p>$x_1 = 3.529, x_2 = 3.538, x_3 = 3.534, \text{ to 3 dp.}$</p>	<p>An attempt to substitute $x_0 = 3.55$ into the iterative formula. Can be implied by $x_1 = 3.528(97)...$ Both $x_1 = \text{awrt } 3.529$ and $x_2 = \text{awrt } 3.538$</p> <p>M1</p> <p>A1</p> <p>x_1, x_2, x_3 all stated correctly to 3 dp</p> <p>A1</p> <p>(3) [13]</p>

6. The function f is defined by

$$f: x \mapsto \frac{3 - 2x}{x - 5}, \quad x \in \mathbb{R}, \quad x \neq 5$$

(a) Find $f^{-1}(x)$.

(3)




Figure 2

The function g has domain $-1 \leq x \leq 8$, and is linear from $(-1, -9)$ to $(2, 0)$ and from $(2, 0)$ to $(8, 4)$. Figure 2 shows a sketch of the graph of $y = g(x)$.

(b) Write down the range of g .

(1)

(c) Find $gg(2)$.

(2)

(d) Find $fg(8)$.

(2)

(e) On separate diagrams, sketch the graph with equation

(i) $y = |g(x)|$,

(ii) $y = g^{-1}(x)$.


Show on each sketch the coordinates of each point at which the graph meets or cuts the axes.


(4)

(f) State the domain of the inverse function g^{-1} .

(1)



Question Number	Scheme	Marks
6.		
(a)	$y = \frac{3-2x}{x-5} \Rightarrow y(x-5) = 3-2x$ <p>Attempt to make x (or swapped y) the subject</p> $xy - 5y = 3 - 2x$ $\Rightarrow xy + 2x = 3 + 5y \Rightarrow x(y+2) = 3 + 5y$ <p>Collect x terms together and factorise.</p> $\Rightarrow x = \frac{3+5y}{y+2} \therefore f^{-1}(x) = \frac{3+5x}{x+2}$	<p>M1</p> <p>M1</p> <p>A1 oe (3)</p>
(b)	<p>Range of g is $-9 \leq g(x) \leq 4$ or $-9 \leq y \leq 4$</p> <p>Correct Range</p>	<p>B1 (1)</p>
(c)	<p>$g g(2) = g(0) = -6$, from sketch.</p>	<p>Deduces that $g(2)$ is 0. Seen or implied.</p> <p>-6</p> <p>M1</p> <p>A1 (2)</p>
(d)	<p>$fg(8) = f(4)$</p> $= \frac{3-4(2)}{4-5} = \frac{-5}{-1} = 5$	<p>Correct order g followed by f</p> <p>5</p> <p>M1</p> <p>A1 (2)</p>
(e)(i)		<p>Correct shape</p> <p>(2, {0}), ({0}, 6)</p> <p>B1</p> <p>B1</p>

Question Number	Scheme	Marks
(e)(ii)	 <p>Correct shape</p> <p>Graph goes through $(\{0\}, 2)$ and $(-6, \{0\})$ which are marked.</p>	<p>B1</p> <p>B1</p> <p>(4)</p>
(f)	<p>Domain of g^{-1} is $-9 \leq x \leq 4$</p> <p>Either correct answer or a follow through from part (b) answer</p>	<p>B1 $\sqrt{\quad}$</p> <p>(1)</p> <p>[13]</p>

7. The curve C has equation

$$y = \frac{3 + \sin 2x}{2 + \cos 2x}$$

(a) Show that

$$\frac{dy}{dx} = \frac{6 \sin 2x + 4 \cos 2x + 2}{(2 + \cos 2x)^2}$$

(4)

(b) Find an equation of the tangent to C at the point on C where $x = \frac{\pi}{2}$.

Write your answer in the form $y = ax + b$, where a and b are exact constants.

(4)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
7	<p>(a) $y = \frac{3 + \sin 2x}{2 + \cos 2x}$</p> <p>Apply quotient rule:</p> $\left\{ \begin{array}{l} u = 3 + \sin 2x \quad v = 2 + \cos 2x \\ \frac{du}{dx} = 2 \cos 2x \quad \frac{dv}{dx} = -2 \sin 2x \end{array} \right\}$ $\frac{dy}{dx} = \frac{2 \cos 2x(2 + \cos 2x) - (-2 \sin 2x)(3 + \sin 2x)}{(2 + \cos 2x)^2}$ $= \frac{4 \cos 2x + 2 \cos^2 2x + 6 \sin 2x + 2 \sin^2 2x}{(2 + \cos 2x)^2}$ $= \frac{4 \cos 2x + 6 \sin 2x + 2(\cos^2 2x + \sin^2 2x)}{(2 + \cos 2x)^2}$ $= \frac{4 \cos 2x + 6 \sin 2x + 2}{(2 + \cos 2x)^2} \quad (\text{as required})$	<p>Applying $\frac{u'v - uv'}{v^2}$ M1</p> <p>Any one term correct on the numerator A1</p> <p>Fully correct (unsimplified). A1</p> <p>For correct proof with an understanding that $\cos^2 2x + \sin^2 2x = 1$. A1*</p> <p>No errors seen in working. A1*</p> <p>(4)</p>
(b)	<p>When $x = \frac{\pi}{2}$, $y = \frac{3 + \sin \pi}{2 + \cos \pi} = \frac{3}{1} = 3$</p> <p>At $(\frac{\pi}{2}, 3)$, $m(\mathbf{T}) = \frac{6 \sin \pi + 4 \cos \pi + 2}{(2 + \cos \pi)^2} = \frac{-4 + 2}{1^2} = -2$</p> <p>Either \mathbf{T}: $y - 3 = -2(x - \frac{\pi}{2})$ or $y = -2x + c$ and $3 = -2(\frac{\pi}{2}) + c \Rightarrow c = 3 + \pi$; \mathbf{T}: $y = -2x + (\pi + 3)$</p>	<p>$y = 3$ B1</p> <p>$m(\mathbf{T}) = -2$ B1</p> <p>$y - y_1 = m(x - \frac{\pi}{2})$ with 'their TANGENT gradient' and their y_1; M1 or uses $y = mx + c$ with 'their TANGENT gradient';</p> <p>$y = -2x + \pi + 3$ A1</p> <p>(4) [8]</p>

Leave
blank

$$\frac{d}{dx}(\cos x) = -\sin x$$

show that $\frac{d}{dx}(\sec x) = \sec x \tan x$.

(3)

Given that

$$x = \sec 2y$$

(b) find $\frac{dx}{dy}$ in terms of y .

(2)

(c) Hence find $\frac{dy}{dx}$ in terms of x .

(4)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
8.		
(a)	$y = \sec x = \frac{1}{\cos x} = (\cos x)^{-1}$ $\frac{dy}{dx} = -1(\cos x)^{-2}(-\sin x)$ $\frac{dy}{dx} = \left\{ \frac{\sin x}{\cos^2 x} \right\} = \left(\frac{1}{\cos x} \right) \left(\frac{\sin x}{\cos x} \right) = \underline{\underline{\sec x \tan x}}$	<p>Writes $\sec x$ as $(\cos x)^{-1}$ and gives</p> $\frac{dy}{dx} = \pm((\cos x)^{-2}(\sin x))$ <p>$-1(\cos x)^{-2}(-\sin x)$ or $(\cos x)^{-2}(\sin x)$</p> <p>Convincing proof. Must see both <u>underlined steps.</u></p> <p>M1 A1 A1 AG (3)</p>
(b)	$x = \sec 2y, \quad y \neq (2n+1)\frac{\pi}{4}, n \in \mathbb{Z}.$ $\frac{dx}{dy} = 2 \sec 2y \tan 2y$	<p>$K \sec 2y \tan 2y$ $2 \sec 2y \tan 2y$</p> <p>M1 A1 (2)</p>
(c)	$\frac{dy}{dx} = \frac{1}{2 \sec 2y \tan 2y}$ $\frac{dy}{dx} = \frac{1}{2x \tan 2y}$ $1 + \tan^2 A = \sec^2 A \Rightarrow \tan^2 2y = \sec^2 2y - 1$ <p>So $\tan^2 2y = x^2 - 1$</p> $\frac{dy}{dx} = \frac{1}{2x\sqrt{(x^2-1)}}$	<p>Applies $\frac{dy}{dx} = \frac{1}{\left(\frac{dx}{dy}\right)}$</p> <p>Substitutes x for $\sec 2y$.</p> <p>Attempts to use the identity $1 + \tan^2 A = \sec^2 A$</p> <p>$\frac{dy}{dx} = \frac{1}{2x\sqrt{(x^2-1)}}$</p> <p>M1 M1 M1 A1 (4) [9]</p>