

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	5	/	0	1	Signature	

Paper Reference(s)

6665/01

Edexcel GCE

Core Mathematics C3

Advanced

Tuesday 15 June 2010 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2010 Edexcel Limited

Printer's Log. No.

Printer's Log. No.
H35385A

W850/R6665/57570 4/5/3/3/3



Turn over

edexcel 
advancing learning, changing lives

Leave
blank

$$\frac{\sin 2\theta}{1 + \cos 2\theta} = \tan \theta$$

(2)

$$\frac{2 \sin 2\theta}{1 + \cos 2\theta} = 1$$

(3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

June 2010
6665 Core Mathematics C3
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) $\frac{2 \sin \theta \cos \theta}{1 + 2 \cos^2 \theta - 1}$ $\frac{\cancel{2} \sin \theta \cancel{\cos \theta}}{\cancel{2} \cos \theta \cancel{\cos \theta}} = \tan \theta$ (as required) AG</p> <p>(b) $2 \tan \theta = 1 \Rightarrow \tan \theta = \frac{1}{2}$ $\theta_1 = \text{awrt } 26.6^\circ$ $\theta_2 = \text{awrt } -153.4^\circ$</p>	<p>M1</p> <p>A1 cso</p> <p>(2)</p> <p>M1</p> <p>A1</p> <p>A1 $\sqrt{}$</p> <p>(3)</p> <p>[5]</p>
	<p>(a) M1: Uses both a correct identity for $\sin 2\theta$ and a correct identity for $\cos 2\theta$. Also allow a candidate writing $1 + \cos 2\theta = 2 \cos^2 \theta$ on the denominator. Also note that angles must be consistent in when candidates apply these identities. A1: Correct proof. No errors seen.</p> <p>(b) 1st M1 for either $2 \tan \theta = 1$ or $\tan \theta = \frac{1}{2}$, seen or implied. A1: awrt 26.6 A1 $\sqrt{}$: awrt -153.4° or $\theta_2 = -180^\circ + \theta_1$</p> <p>Special Case: For candidate solving, $\tan \theta = k$, where $k \neq \frac{1}{2}$, to give θ_1 and $\theta_2 = -180^\circ + \theta_1$, then award M0A0B1 in part (b). Special Case: Note that those candidates who writes $\tan \theta = 1$, and gives ONLY two answers of 45° and -135° that are inside the range will be awarded SC M0A0B1.</p>	

Leave
blank

- $$y = \frac{3}{(5-3x)^2}, \quad x \neq \frac{5}{3}$$

(7)

[illegible]

Question Number	Scheme	Marks
2.	<p>At P, $y = 3$</p> $\frac{dy}{dx} = \frac{3(-2)(5-3x)^{-3}(-3)}{(5-3x)^3} \left\{ \text{or } \frac{18}{(5-3x)^3} \right\}$ $\frac{dy}{dx} = \frac{18}{(5-3(2))^3} \{ = -18 \}$ $m(N) = \frac{-1}{-18} \text{ or } \frac{1}{18}$ $N: y - 3 = \frac{1}{18}(x - 2)$ $N: \underline{x - 18y + 52 = 0}$	<p>B1</p> <p>M1A1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[7]</p>
	<p>1st M1: $\pm k(5-3x)^{-3}$ can be implied. See appendix for application of the quotient rule.</p> <p>2nd M1: Substituting $x = 2$ into an equation involving their $\frac{dy}{dx}$;</p> <p>3rd M1: Uses $m(N) = -\frac{1}{\text{their } m(T)}$.</p> <p>4th M1: $y - y_1 = m(x - 2)$ with 'their NORMAL gradient' or a "changed" tangent gradient and their y_1. Or uses a complete method to express the equation of the tangent in the form $y = mx + c$ with 'their NORMAL ("changed" numerical) gradient', their y_1 and $x = 2$.</p> <p>Note: To gain the final A1 mark all the previous 6 marks in this question need to be earned. Also there must be a completely correct solution given.</p>	

3. $f(x) = 4 \operatorname{cosec} x - 4x + 1$, where x is in radians.

- (b) Show that the equation $f(x) = 0$ can be written in the form

$$x = \frac{1}{\sin x} + \frac{1}{4} \quad (2)$$

- (c) Use the iterative formula

$$x_{n+1} = \frac{1}{\sin x_n} + \frac{1}{4}, \quad x_0 = 1.25,$$

to calculate the values of x_1 , x_2 and x_3 , giving your answers to 4 decimal places. **(3)**

- (d) By considering the change of sign of $f(x)$ in a suitable interval, verify that $\alpha=1.291$ correct to 3 decimal places.



Question Number	Scheme	Marks
3.	<p>(a) $f(1.2) = 0.49166551\dots$, $f(1.3) = -0.048719817\dots$ Sign change (and as $f(x)$ is continuous) therefore a root α is such that $\alpha \in [1.2, 1.3]$</p> <p>(b) $4\operatorname{cosec} x - 4x + 1 = 0 \Rightarrow 4x = 4\operatorname{cosec} x + 1$ $\Rightarrow x = \operatorname{cosec} x + \frac{1}{4} \Rightarrow x = \frac{1}{\sin x} + \frac{1}{4}$</p> <p>(c) $x_1 = \frac{1}{\sin(1.25)} + \frac{1}{4}$ $x_1 = 1.303757858\dots$, $x_2 = 1.286745793\dots$ $x_3 = 1.291744613\dots$</p> <p>(d) $f(1.2905) = 0.00044566695\dots$, $f(1.2915) = -0.00475017278\dots$ Sign change (and as $f(x)$ is continuous) therefore a root α is such that $\alpha \in (1.2905, 1.2915) \Rightarrow \alpha = 1.291$ (3 dp)</p>	<p>M1A1 (2)</p> <p>M1 A1 *</p> <p>(2)</p> <p>M1 A1 A1 (3)</p> <p>M1 A1 (2) [9]</p>
	<p>(a) M1: Attempts to evaluate both $f(1.2)$ and $f(1.3)$ and evaluates at least one of them correctly to awrt (or truncated) 1 sf. A1: both values correct to awrt (or truncated) 1 sf, sign change and conclusion.</p> <p>(b) M1: Attempt to make $4x$ or x the subject of the equation. A1: Candidate must then rearrange the equation to give the required result. It must be clear that candidate has made their initial $f(x) = 0$.</p> <p>(c) M1: An attempt to substitute $x_0 = 1.25$ into the iterative formula Eg $= \frac{1}{\sin(1.25)} + \frac{1}{4}$. Can be implied by $x_1 = \text{awrt } 1.3$ or $x_1 = \text{awrt } 46^\circ$. A1: Both $x_1 = \text{awrt } 1.3038$ and $x_2 = \text{awrt } 1.2867$ A1: $x_3 = \text{awrt } 1.2917$</p> <p>(d) M1: Choose suitable interval for x, e.g. $[1.2905, 1.2915]$ or tighter and at least one attempt to evaluate $f(x)$. A1: both values correct to awrt (or truncated) 1 sf, sign change and conclusion.</p>	

4. The function f is defined by

$$f : x \mapsto |2x - 5|, \quad x \in \mathbb{R}$$

- (a) Sketch the graph with equation $y = f(x)$, showing the coordinates of the points where the graph cuts or meets the axes.

(2)

- (b) Solve $f(x) = 15 + x$.

(3)

The function g is defined by

$$g : x \mapsto x^2 - 4x + 1, \quad x \in \mathbb{R}, \quad 0 \leq x \leq 5$$




- (c) Find $fg(2)$.

(2)

- (d) Find the range of g .

(3)



Question Number	Scheme	Marks
4.	<p>(a) </p> <p>(b) $x = 20$ $2x - 5 = -(15 + x) ; \Rightarrow x = -\frac{10}{3}$</p> <p>(c) $fg(2) = f(-3) = 2(-3) - 5 = -11 = 11$</p> <p>(d) $g(x) = x^2 - 4x + 1 = (x - 2)^2 - 4 + 1 = (x - 2)^2 - 3$. Hence $g_{\min} = -3$ Either $g_{\min} = -3$ or $g(x) \geq -3$ or $g(5) = 25 - 20 + 1 = 6$ $-3 \leq g(x) \leq 6$ or $-3 \leq y \leq 6$</p>	<p>M1A1</p> <p>(2)</p> <p>B1 M1;A1 oe.</p> <p>(3)</p> <p>M1;A1</p> <p>(2)</p> <p>M1</p> <p>B1</p> <p>A1</p> <p>(3)</p> <p>[10]</p>
	<p>(a) M1: V or  or  graph with vertex on the x-axis.</p> <p>A1: $(\frac{5}{2}, \{0\})$ and $(\{0\}, 5)$ seen and the graph appears in both the first and second quadrants.</p> <p>(b) M1: Either $2x - 5 = -(15 + x)$ or $-(2x - 5) = 15 + x$</p> <p>(c) M1: Full method of inserting $g(2)$ into $f(x) = 2x - 5$ or for inserting $x = 2$ into $2(x^2 - 4x + 1) - 5$. There must be evidence of the modulus being applied.</p> <p>(d) M1: Full method to establish the minimum of g. Eg: $(x \pm \alpha)^2 + \beta$ leading to $g_{\min} = \beta$. Or for candidate to differentiate the quadratic, set the result equal to zero, find x and insert this value of x back into $f(x)$ in order to find the minimum.</p> <p>B1: For either finding the correct minimum value of g (can be implied by $g(x) \geq -3$ or $g(x) > -3$) or for stating that $g(5) = 6$.</p> <p>A1: $-3 \leq g(x) \leq 6$ or $-3 \leq y \leq 6$ or $-3 \leq g \leq 6$. Note that: $-3 \leq x \leq 6$ is A0.</p> <p>Note that: $-3 \leq f(x) \leq 6$ is A0. Note that: $-3 \geq g(x) \geq 6$ is A0.</p> <p>Note that: $g(x) \geq -3$ or $g(x) > -3$ or $x \geq -3$ or $x > -3$ with no working gains M1B1A0.</p> <p>Note that for the final Accuracy Mark: If a candidate writes down $-3 < g(x) < 6$ or $-3 < y < 6$, then award M1B1A0. If, however, a candidate writes down $g(x) \geq -3$, $g(x) \leq 6$, then award A0. If a candidate writes down $g(x) \geq -3$ or $g(x) \leq 6$, then award A0.</p>	

5.




Figure 1

(a) Find the coordinates of the point where C crosses the y -axis. (1)

(b) Show that C crosses the x -axis at $x = 2$ and find the x -coordinate of the other point where C crosses the x -axis. (3)

(c) Find $\frac{dy}{dx}$. (3)

(d) Hence find the exact coordinates of the turning points of C . (5)

[illegible]

Question Number	Scheme	Marks
5.	(a) Either $y = 2$ or $(0, 2)$	B1 (1)
	(b) When $x = 2$, $y = (8 - 10 + 2)e^{-2} = 0e^{-2} = 0$ $(2x^2 - 5x + 2) = 0 \Rightarrow (x - 2)(2x - 1) = 0$ Either $x = 2$ (for possibly B1 above) or $x = \frac{1}{2}$.	B1 M1 A1 (3)
	(c) $\frac{dy}{dx} = (4x - 5)e^{-x} - (2x^2 - 5x + 2)e^{-x}$	M1A1A1 (3)
	(d) $(4x - 5)e^{-x} - (2x^2 - 5x + 2)e^{-x} = 0$ $2x^2 - 9x + 7 = 0 \Rightarrow (2x - 7)(x - 1) = 0$ $x = \frac{7}{2}, 1$ When $x = \frac{7}{2}$, $y = 9e^{-\frac{7}{2}}$, when $x = 1$, $y = -e^{-1}$	M1 M1 A1 ddM1A1 (5) [12]
	(b) If the candidate believes that $e^{-x} = 0$ solves to $x = 0$ or gives an extra solution of $x = 0$, then withhold the final accuracy mark. (c) M1: (their u') $e^{-x} + (2x^2 - 5x + 2)(\text{their } v')$ A1: Any one term correct. A1: Both terms correct. (d) 1 st M1: For setting their $\frac{dy}{dx}$ found in part (c) equal to 0. 2 nd M1: Factorise or eliminate out e^{-x} correctly and an attempt to factorise a 3-term quadratic or apply the formula to candidate's $ax^2 + bx + c$. See rules for solving a three term quadratic equation on page 1 of this Appendix. 3 rd ddM1: An attempt to use at least one x -coordinate on $y = (2x^2 - 5x + 2)e^{-x}$. Note that this method mark is dependent on the award of the two previous method marks in this part. Some candidates write down corresponding y -coordinates without any working. It may be necessary on some occasions to use your calculator to check that at least one of the two y -coordinates found is correct to awrt 2 sf. Final A1: Both $\{x = 1\}$, $y = -e^{-1}$ and $\{x = \frac{7}{2}\}$, $y = 9e^{-\frac{7}{2}}$. cao Note that both exact values of y are required.	

7. (a) Express $2 \sin \theta - 1.5 \cos \theta$ in the form $R \sin(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$.

(3)

- (ii) Find the value of θ , for $0 \leq \theta < \pi$, at which this maximum occurs.

(3)

$$H=6+2\sin\left(\frac{4\pi t}{25}\right)-1.5\cos\left(\frac{4\pi t}{25}\right), \quad 0\leq t<12,$$

(c) Calculate the maximum value of H predicted by this model and the value of t , to 2 decimal places, when this maximum occurs.

(3)

- (d) Calculate, to the nearest minute, the times when the height of sea water is predicted, by this model, to be 7 metres.

(6)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
7.	<p>(a) $R = \sqrt{6.25}$ or 2.5 $\tan \alpha = \frac{1.5}{2} = \frac{3}{4} \Rightarrow \alpha = \text{awrt } 0.6435$</p> <p>(b) (i) Max Value = 2.5 (ii) $\sin(\theta - 0.6435) = 1$ or $\theta - \text{their } \alpha = \frac{\pi}{2}; \Rightarrow \theta = \text{awrt } 2.21$</p> <p>(c) $H_{\text{Max}} = 8.5$ (m) $\sin\left(\frac{4\pi t}{25} - 0.6435\right) = 1$ or $\frac{4\pi t}{25} = \text{their (b) answer}; \Rightarrow t = \text{awrt } 4.41$</p> <p>(d) $\Rightarrow 6 + 2.5 \sin\left(\frac{4\pi t}{25} - 0.6435\right) = 7; \Rightarrow \sin\left(\frac{4\pi t}{25} - 0.6435\right) = \frac{1}{2.5} = 0.4$ $\left\{\frac{4\pi t}{25} - 0.6435\right\} = \sin^{-1}(0.4)$ or awrt 0.41 Either $t = \text{awrt } 2.1$ or awrt 6.7 So, $\left\{\frac{4\pi t}{25} - 0.6435\right\} = \{\pi - 0.411517... \text{ or } 2.730076...^c\}$ Times = $\{14:06, 18:43\}$</p>	<p>B1 M1A1 (3)</p> <p>B1 $\sqrt{\quad}$ M1;A1 $\sqrt{\quad}$ (3)</p> <p>B1 $\sqrt{\quad}$ M1;A1 (3)</p> <p>M1;M1 A1 A1 ddM1 A1 (6) [15]</p>
	<p>(a) B1: $R = 2.5$ or $R = \sqrt{6.25}$. For $R = \pm 2.5$, award B0. M1: $\tan \alpha = \pm \frac{1.5}{2}$ or $\tan \alpha = \pm \frac{2}{1.5}$ A1: $\alpha = \text{awrt } 0.6435$</p> <p>(b) B1 $\sqrt{\quad}$: 2.5 or follow through the value of R in part (a). M1: For $\sin(\theta - \text{their } \alpha) = 1$ A1 $\sqrt{\quad}$: awrt 2.21 or $\frac{\pi}{2} + \text{their } \alpha$ rounding correctly to 3 sf.</p> <p>(c) B1 $\sqrt{\quad}$: 8.5 or 6 + their R found in part (a) as long as the answer is greater than 6. M1: $\sin\left(\frac{4\pi t}{25} \pm \text{their } \alpha\right) = 1$ or $\frac{4\pi t}{25} = \text{their (b) answer}$ A1: For $\sin^{-1}(0.4)$ This can be implied by awrt 4.41 or awrt 4.40.</p> <p>(d) M1: $6 + (\text{their } R) \sin\left(\frac{4\pi t}{25} \pm \text{their } \alpha\right) = 7$, M1: $\sin\left(\frac{4\pi t}{25} \pm \text{their } \alpha\right) = \frac{1}{\text{their } R}$ A1: For $\sin^{-1}(0.4)$. This can be implied by awrt 0.41 or awrt 2.73 or other values for different α's. Note this mark can be implied by seeing 1.055. A1: Either $t = \text{awrt } 2.1$ or $t = \text{awrt } 6.7$ ddM1: either $\pi - \text{their PV}^c$. Note that this mark is dependent upon the two M marks. This mark will usually be awarded for seeing either 2.730... or 3.373... A1: Both $t = 14:06$ and $t = 18:43$ or both 126 (min) and 403 (min) or both 2 hr 6 min and 6 hr 43 min.</p>	

Leave
blank

$$\frac{2x^2 + 9x - 5}{x^2 + 2x - 15}$$

(3)

$$\ln(2x^2 + 9x - 5) = 1 + \ln(x^2 + 2x - 15), \quad x \neq -5,$$

(4)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
8.	<p>(a) $\frac{(x+5)(2x-1)}{(x+5)(x-3)} = \frac{(2x-1)}{(x-3)}$</p> <p>(b) $\ln\left(\frac{2x^2+9x-5}{x^2+2x-15}\right) = 1$</p> <p>$\frac{2x^2+9x-5}{x^2+2x-15} = e$</p> <p>$\frac{2x-1}{x-3} = e \Rightarrow 3e-1 = x(e-2)$</p> <p>$\Rightarrow x = \frac{3e-1}{e-2}$</p>	<p>M1 B1 A1 aef (3)</p> <p>M1</p> <p>dM1</p> <p>M1</p> <p>A1 aef cso (4) [7]</p>
	<p>(a) M1: An attempt to factorise the numerator. B1: Correct factorisation of denominator to give $(x+5)(x-3)$. Can be seen anywhere.</p> <p>(b) M1: Uses a correct law of logarithms to combine at least two terms. This usually is achieved by the subtraction law of logarithms to give $\ln\left(\frac{2x^2+9x-5}{x^2+2x-15}\right) = 1$.</p> <p>The product law of logarithms can be used to achieve $\ln(2x^2+9x-5) = \ln(e(x^2+2x-15))$.</p> <p>The product and quotient law could also be used to achieve $\ln\left(\frac{2x^2+9x-5}{e(x^2+2x-15)}\right) = 0$.</p> <p>dM1: Removing \ln's correctly by the realisation that the anti-\ln of 1 is e. Note that this mark is dependent on the previous method mark being awarded.</p> <p>M1: Collect x terms together and factorise. Note that this is not a dependent method mark.</p> <p>A1: $\frac{3e-1}{e-2}$ or $\frac{3e^1-1}{e^1-2}$ or $\frac{1-3e}{2-e}$. aef</p> <p>Note that the answer needs to be in terms of e. The decimal answer is 9.9610559... Note that the solution must be correct in order for you to award this final accuracy mark.</p> <p>Note: See Appendix for an alternative method of long division.</p>	