

Question Number	Scheme	Marks												
1.	<p>Differentiates</p> <p>to obtain : $6x + 8y \frac{dy}{dx} - 2,$ $\dots\dots\dots + (6x \frac{dy}{dx} + 6y) = 0$</p> $\left[\frac{dy}{dx} = \frac{2 - 6x - 6y}{6x + 8y} \right]$ <p>Substitutes $x = 1, y = -2$ into expression involving $\frac{dy}{dx}$, to give $\frac{dy}{dx} = -\frac{8}{10}$</p> <p>Uses line equation with numerical 'gradient' $y - (-2) = (\text{their gradient})(x - 1)$ or finds c and uses $y = (\text{their gradient})x + "c"$</p> <p>To give $5y + 4x + 6 = 0$ (or equivalent = 0)</p>	<p>M1</p> <p>A1,</p> <p>+(B1)</p> <p>M1, A1</p> <p>M1</p> <p>A1√ [7]</p>												
2. (a)	<table border="1" data-bbox="220 976 1294 1133"> <tr> <td>x</td> <td>0</td> <td>$\frac{\pi}{16}$</td> <td>$\frac{\pi}{8}$</td> <td>$\frac{3\pi}{16}$</td> <td>$\frac{\pi}{4}$</td> </tr> <tr> <td>y</td> <td>1</td> <td>1.01959</td> <td>1.08239</td> <td>1.20269</td> <td>1.41421</td> </tr> </table> <p>M1 for one correct, A1 for all correct</p>	x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$	y	1	1.01959	1.08239	1.20269	1.41421	<p>M1 A1</p> <p>(2)</p>
x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$									
y	1	1.01959	1.08239	1.20269	1.41421									
(b)	<p>Integral = $\frac{1}{2} \times \frac{\pi}{16} \times \{1 + 1.4142 + 2(1.01959 + \dots + 1.20269)\}$</p> $\left(= \frac{\pi}{32} \times 9.02355 \right) = 0.8859$	<p>M1 A1√</p> <p>A1 cao</p> <p>(3)</p>												
(c)	<p>Percentage error = $\frac{\text{approx} - 0.88137}{0.88137} \times 100 = 0.51\%$ (allow 0.5% to 0.54% for A1)</p> <p>M1 gained for $(\pm) \frac{\text{approx} - \ln(1 + \sqrt{2})}{\ln(1 + \sqrt{2})}$</p>	<p>M1 A1</p> <p>(2)</p> <p>[7]</p>												

Question Number	Scheme	Marks												
1.	<p>Differentiates</p> <p>to obtain : $6x + 8y \frac{dy}{dx} - 2,$ $\dots\dots\dots + (6x \frac{dy}{dx} + 6y) = 0$</p> $\left[\frac{dy}{dx} = \frac{2 - 6x - 6y}{6x + 8y} \right]$ <p>Substitutes $x = 1, y = -2$ into expression involving $\frac{dy}{dx}$, to give $\frac{dy}{dx} = -\frac{8}{10}$</p> <p>Uses line equation with numerical 'gradient' $y - (-2) = (\text{their gradient})(x - 1)$ or finds c and uses $y = (\text{their gradient})x + "c"$</p> <p>To give $5y + 4x + 6 = 0$ (or equivalent = 0)</p>	<p>M1</p> <p>A1,</p> <p>+(B1)</p> <p>M1, A1</p> <p>M1</p> <p>A1√ [7]</p>												
2.	<table border="1" data-bbox="220 974 1294 1133"> <tr> <td>x</td> <td>0</td> <td>$\frac{\pi}{16}$</td> <td>$\frac{\pi}{8}$</td> <td>$\frac{3\pi}{16}$</td> <td>$\frac{\pi}{4}$</td> </tr> <tr> <td>y</td> <td>1</td> <td>1.01959</td> <td>1.08239</td> <td>1.20269</td> <td>1.41421</td> </tr> </table> <p>M1 for one correct, A1 for all correct</p> <p>(b) Integral = $\frac{1}{2} \times \frac{\pi}{16} \times \{1 + 1.4142 + 2(1.01959 + \dots + 1.20269)\}$ $\left(= \frac{\pi}{32} \times 9.02355 \right) = 0.8859$</p> <p>(c) Percentage error = $\frac{\text{approx} - 0.88137}{0.88137} \times 100 = 0.51\%$ (allow 0.5% to 0.54% for A1)</p> <p>M1 gained for $(\pm) \frac{\text{approx} - \ln(1 + \sqrt{2})}{\ln(1 + \sqrt{2})}$</p>	x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$	y	1	1.01959	1.08239	1.20269	1.41421	<p>M1 A1</p> <p>(2)</p> <p>M1 A1√</p> <p>A1 cao</p> <p>(3)</p> <p>M1 A1 (2)</p> <p>[7]</p>
x	0	$\frac{\pi}{16}$	$\frac{\pi}{8}$	$\frac{3\pi}{16}$	$\frac{\pi}{4}$									
y	1	1.01959	1.08239	1.20269	1.41421									

Question Number	Scheme	Marks
3.	<p>Uses substitution to obtain $x = f(u) \left[\frac{u^2 + 1}{2} \right]$,</p> <p>and to obtain $u \frac{du}{dx} = \text{const. or equiv.}$</p> <p>Reaches $\int \frac{3(u^2 + 1)}{2u} u du$ or equivalent</p> <p>Simplifies integrand to $\int \left(3u^2 + \frac{3}{2} \right) du$ or equiv.</p> <p>Integrates to $\frac{1}{2}u^3 + \frac{3}{2}u$</p> <p>A1√ dependent on all previous Ms</p> <p>Uses new limits 3 and 1 substituting and subtracting (or returning to function of x with old limits)</p> <p>To give 16 cso</p> <hr/> <p>“By Parts” Attempt at “right direction” by parts M1</p> $\left[3x \left(2x - 1 \right)^{\frac{1}{2}} \right] - \left\{ \int 3 \left(2x - 1 \right)^{\frac{1}{2}} dx \right\} \quad \text{M1}\{\text{M1A1}\}$ $\dots\dots\dots - \left(2x - 1 \right)^{\frac{3}{2}} \quad \text{M1A1}\checkmark$ <p>Uses limits 5 and 1 correctly; [42 – 26] 16 M1A1</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>M1 A1√</p> <p>M1</p> <p>A1</p> <p>[8]</p>

<p>4.</p>	<p>Attempts $V = \pi \int x^2 e^{2x} dx$</p> $= \pi \left[\frac{x^2 e^{2x}}{2} - \int x e^{2x} dx \right]$ <p>(M1 needs parts in the correct direction)</p> $= \pi \left[\frac{x^2 e^{2x}}{2} - \left(\frac{x e^{2x}}{2} - \int \frac{e^{2x}}{2} dx \right) \right]$ <p>(M1 needs second application of parts)</p> <p>M1A1√ refers to candidates $\int x e^{2x} dx$, but dependent on prev. M1</p> $= \pi \left[\frac{x^2 e^{2x}}{2} - \left(\frac{x e^{2x}}{2} - \frac{e^{2x}}{4} \right) \right]$ <p>Substitutes limits 3 and 1 and subtracts to give... [dep. on second and third Ms]</p> $= \pi \left[\frac{13}{4} e^6 - \frac{1}{4} e^2 \right] \text{ or any correct exact equivalent.}$ <p>[Omission of π loses first and last marks only]</p>	<p>M1</p> <p>M1 A1</p> <p>M1 A1√</p> <p>A1 cao</p> <p>dM1</p> <p>A1</p> <p>[8]</p>
-----------	--	--

Question Number	Scheme	Marks
5. (a)	<p>Considers $3x^2 + 16 = A(2+x)^2 + B(1-3x)(2+x) + C(1-3x)$</p> <p>and substitutes $x = -2$, or $x = 1/3$,</p> <p>or compares coefficients and solves simultaneous equations</p> <p>To obtain $A = 3$, and $C = 4$</p> <p>Compares coefficients or uses simultaneous equation to show $B = 0$.</p>	<p>M1</p> <p>A1, A1</p> <p>B1</p> <p>(4)</p>
5. (b)	<p>Writes $3(1-3x)^{-1} + 4(2+x)^{-2}$</p> <p>$= 3(1+3x, +9x^2 + 27x^3 + \dots) +$</p> $\frac{4}{4} \left(1 + \frac{(-2)}{1} \left(\frac{x}{2} \right) + \frac{(-2)(-3)}{1.2} \left(\frac{x}{2} \right)^2 + \frac{(-2)(-3)(-4)}{1.2.3} \left(\frac{x}{2} \right)^3 + \dots \right)$ <p>$= 4 + 8x, + 27\frac{3}{4}x^2 + 80\frac{1}{2}x^3 + \dots$</p> <p>Or uses $(3x^2 + 16)(1-3x)^{-1}(2+x)^{-2}$</p> <p>$(3x^2 + 16) (1 + 3x, +9x^2 + 27x^3 +) \times$</p> $\frac{1}{4} \left(1 + \frac{(-2)}{1} \left(\frac{x}{2} \right) + \frac{(-2)(-3)}{1.2} \left(\frac{x}{2} \right)^2 + \frac{(-2)(-3)(-4)}{1.2.3} \left(\frac{x}{2} \right)^3 \right)$ <p>$= 4 + 8x, + 27\frac{3}{4}x^2 + 80\frac{1}{2}x^3 + \dots$</p>	<p>M1</p> <p>(M1, A1)</p> <p>(M1 A1)</p> <p>A1, A1</p> <p>(7)</p> <p>M1</p> <p>(M1A1)×</p> <p>(M1A1)</p> <p>A1, A1</p> <p>(7)</p> <p>[11]</p>

6. The line l_1 has vector equation

$$\mathbf{r} = 8\mathbf{i} + 12\mathbf{j} + 14\mathbf{k} + \lambda(\mathbf{i} + \mathbf{j} - \mathbf{k}),$$

where λ is a parameter.

The point A has coordinates $(4, 8, a)$, where a is a constant. The point B has coordinates $(b, 13, 13)$, where b is a constant. Points A and B lie on the line l_1 .

(a) Find the values of a and b .

(3)

Given that the point O is the origin, and that the point P lies on l_1 such that OP is perpendicular to l_1 ,

(b) find the coordinates of P .

(5)

(c) Hence find the distance OP , giving your answer as a simplified surd.

(2)



<p>6. (a)</p> <p>(b)</p> <p>(c)</p>	<p>$\lambda = -4 \rightarrow a = 18, \quad \mu = 1 \rightarrow b = 9$</p> $\begin{pmatrix} 8 + \lambda \\ 12 + \lambda \\ 14 - \lambda \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 0$ <p>$\therefore 8 + \lambda + 12 + \lambda - 14 + \lambda = 0$</p> <p>Solves to obtain $\lambda \quad (\lambda = -2)$</p> <p>Then substitutes value for λ to give P at the point (6, 10, 16) (any form)</p> <p>$OP = \sqrt{36 + 100 + 256}$</p> <p>$(= \sqrt{392}) = 14\sqrt{2}$</p>	<p>M1 A1, A1 (3)</p> <p>M1</p> <p>A1</p> <p>dM1</p> <p>M1, A1 (5)</p> <p>M1</p> <p>A1 cao (2)</p> <p>[10]</p>
<p>7. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(i)</p> <p>(ii)</p>	<p>$\frac{dV}{dr} = 4\pi r^2$</p> <p>Uses $\frac{dr}{dt} = \frac{dV}{dt} \cdot \frac{dr}{dV}$ in any form, $= \frac{1000}{4\pi r^2 (2t+1)^2}$</p> <p>$V = \int 1000(2t+1)^{-2} dt$ and integrate to $p (2t+1)^{-1}, \quad = -500(2t+1)^{-1} (+c)$</p> <p>Using $V=0$ when $t=0$ to find c, ($c = 500$, or equivalent)</p> <p>$\therefore V = 500(1 - \frac{1}{2t+1})$ (any form)</p> <p>then use $r = \sqrt[3]{\left(\frac{3V}{4\pi}\right)}$ to give $r, = 4.77$</p> <p>Substitutes $t = 5$ and $r =$ 'their value' into 'their' part (b)</p> <p>$\frac{dr}{dt} = 0.0289 \quad (\approx 2.90 \times 10^{-2}) \text{ (cm/s) } * \quad \text{AG}$</p>	<p>B1 (1)</p> <p>M1, A1 (2)</p> <p>M1, A1</p> <p>M1</p> <p>A1 (4)</p> <p>M1,</p> <p>M1, A1 (3)</p> <p>M1</p> <p>A1 (2)</p> <p>[12]</p>

Leave blank

7. The volume of a spherical balloon of radius r cm is V cm³, where $V = \frac{4}{3}\pi r^3$.

(a) Find $\frac{dV}{dr}$. (1)

The volume of the balloon increases with time t seconds according to the formula

$$\frac{dV}{dt} = \frac{1000}{(2t+1)^2}, \quad t \geq 0.$$

(b) Using the chain rule, or otherwise, find an expression in terms of r and t for $\frac{dr}{dt}$. (2)

(c) Given that $V = 0$ when $t = 0$, solve the differential equation $\frac{dV}{dt} = \frac{1000}{(2t+1)^2}$, to obtain V in terms of t . (4)

(d) Hence, at time $t = 5$,

(i) find the radius of the balloon, giving your answer to 3 significant figures, (3)

(ii) show that the rate of increase of the radius of the balloon is approximately 2.90×10^{-2} cm s⁻¹. (2)




<p>6. (a)</p> <p>(b)</p> <p>(c)</p>	<p>$\lambda = -4 \rightarrow a = 18, \quad \mu = 1 \rightarrow b = 9$</p> $\begin{pmatrix} 8 + \lambda \\ 12 + \lambda \\ 14 - \lambda \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 0$ <p>$\therefore 8 + \lambda + 12 + \lambda - 14 + \lambda = 0$</p> <p>Solves to obtain $\lambda \quad (\lambda = -2)$</p> <p>Then substitutes value for λ to give P at the point (6, 10, 16) (any form)</p> <p>$OP = \sqrt{36 + 100 + 256}$</p> <p>$(= \sqrt{392}) = 14\sqrt{2}$</p>	<p>M1 A1, A1 (3)</p> <p>M1</p> <p>A1</p> <p>dM1</p> <p>M1, A1 (5)</p> <p>M1</p> <p>A1 cao (2)</p> <p>[10]</p>
<p>7. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(i)</p> <p>(ii)</p>	<p>$\frac{dV}{dr} = 4\pi r^2$</p> <p>Uses $\frac{dr}{dt} = \frac{dV}{dt} \cdot \frac{dr}{dV}$ in any form, $= \frac{1000}{4\pi r^2 (2t+1)^2}$</p> <p>$V = \int 1000(2t+1)^{-2} dt$ and integrate to $p (2t+1)^{-1}, \quad = -500(2t+1)^{-1} (+c)$</p> <p>Using $V=0$ when $t=0$ to find c, ($c = 500$, or equivalent)</p> <p>$\therefore V = 500(1 - \frac{1}{2t+1})$ (any form)</p> <p>(i) Substitute $t = 5$ to give V, then use $r = \sqrt[3]{\left(\frac{3V}{4\pi}\right)}$ to give $r, = 4.77$</p> <p>(ii) Substitutes $t = 5$ and $r =$ 'their value' into 'their' part (b)</p> <p>$\frac{dr}{dt} = 0.0289 \quad (\approx 2.90 \times 10^{-2}) \text{ (cm/s) } * \quad \text{AG}$</p>	<p>B1 (1)</p> <p>M1, A1 (2)</p> <p>M1, A1</p> <p>M1</p> <p>A1 (4)</p> <p>M1,</p> <p>M1, A1 (3)</p> <p>M1</p> <p>A1 (2)</p> <p>[12]</p>

Leave blank

8.

Figure 2



The curve shown in Figure 2 has parametric equations

$$x = t - 2 \sin t, \quad y = 1 - 2 \cos t, \quad 0 \leq t \leq 2\pi.$$

- (a) Show that the curve crosses the x -axis where $t = \frac{\pi}{3}$ and $t = \frac{5\pi}{3}$. (2)

The finite region R is enclosed by the curve and the x -axis, as shown shaded in Figure 2.

- (b) Show that the area of R is given by the integral

$$\int_{\frac{\pi}{3}}^{\frac{5\pi}{3}} (1 - 2 \cos t)^2 dt. \quad (3)$$

- (c) Use this integral to find the exact value of the shaded area. (7)



8.	<p>(a) Solves $y = 0 \Rightarrow \cos t = \frac{1}{2}$ to obtain $t = \frac{\pi}{3}$ or $\frac{5\pi}{3}$ (need both for A1)</p> <p>Or substitutes both values of t and shows that $y = 0$</p> <p>(b) $\frac{dx}{dt} = 1 - 2 \cos t$</p> <p>Area = $\int y dx = \int_{\frac{\pi}{3}}^{\frac{5\pi}{3}} (1 - 2 \cos t)(1 - 2 \cos t) dt = \int_{\frac{\pi}{3}}^{\frac{5\pi}{3}} (1 - 2 \cos t)^2 dt$ * AG</p> <p>(c) Area = $\int 1 - 4 \cos t + 4 \cos^2 t dt$ 3 terms</p> <p>= $\int 1 - 4 \cos t + 2(\cos 2t + 1) dt$ (use of correct double angle formula)</p> <p>= $\int 3 - 4 \cos t + 2 \cos 2t dt$</p> <p>= $[3t - 4 \sin t + \sin 2t]$</p> <p>Substitutes the two correct limits $t = \frac{5\pi}{3}$ and $\frac{\pi}{3}$ and subtracts.</p> <p>= $4\pi + 3\sqrt{3}$</p>	
----	---	--