

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	6	/	0	1	Signature	

Paper Reference(s)

6666/01

Edexcel GCE

Core Mathematics C4

Advanced

Friday 18 June 2010 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 32 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2010 Edexcel Limited

Printer's Log. No.

Printer's Log. No.
H35386A

W850/R6666/57570 4/5/3/3



Turn over

edexcel 
advancing learning, changing lives

1.




Figure 1

Figure 1 shows part of the curve with equation $y = \sqrt{0.75 + \cos^2 x}$. The finite region R , shown shaded in Figure 1, is bounded by the curve, the y -axis, the x -axis and the line with equation $x = \frac{\pi}{3}$.

(a) Complete the table with values of y corresponding to $x = \frac{\pi}{6}$ and $x = \frac{\pi}{4}$.

x	0	$\frac{\pi}{12}$	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$
y	1.3229	1.2973			1

(2)

(b) Use the trapezium rule

(i) with the values of y at $x = 0$, $x = \frac{\pi}{6}$ and $x = \frac{\pi}{3}$ to find an estimate of the area of R .

Give your answer to 3 decimal places.

(ii) with the values of y at $x = 0$, $x = \frac{\pi}{12}$, $x = \frac{\pi}{6}$, $x = \frac{\pi}{4}$ and $x = \frac{\pi}{3}$ to find a further estimate of the area of R . Give your answer to 3 decimal places.

(6)



June 2010
6666 Core Mathematics C4
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) $y\left(\frac{\pi}{6}\right) \approx 1.2247, y\left(\frac{\pi}{4}\right) = 1.1180$ accept awrt 4 d.p.</p> <p>(b)(i) $I \approx \left(\frac{\pi}{12}\right)(1.3229 + 2 \times 1.2247 + 1)$ B1 for $\frac{\pi}{12}$ ≈ 1.249 cao</p> <p>(ii) $I \approx \left(\frac{\pi}{24}\right)(1.3229 + 2 \times (1.2973 + 1.2247 + 1.1180) + 1)$ B1 for $\frac{\pi}{24}$ ≈ 1.257 cao</p>	<p>B1 B1 (2)</p> <p>B1 M1 A1</p> <p>B1 M1 A1 (6) [8]</p>

2. Using the substitution $u = \cos x + 1$, or otherwise, show that

$$\int_0^{\frac{\pi}{2}} e^{\cos x + 1} \sin x \, dx = e(e - 1)$$

(6)



Question Number	Scheme	Marks
2.	$\frac{du}{dx} = -\sin x$ $\int \sin x e^{\cos x + 1} dx = -\int e^u du$ $= -e^u$ $= -e^{\cos x + 1}$ $\left[-e^{\cos x + 1} \right]_0^{\frac{\pi}{2}} = -e^1 - (-e^2)$ $= e(e - 1) *$	<p>B1</p> <p>M1 A1</p> <p>A1ft</p> <p>ft sign error</p> <p>or equivalent with u</p> <p>M1</p> <p>A1</p> <p>cso</p> <p>(6)</p> <p>[6]</p>

3. A curve C has equation

$$2^x + y^2 = 2xy$$

Find the exact value of $\frac{dy}{dx}$ at the point on C with coordinates $(3, 2)$.

(7)



Question Number	Scheme	Marks
3.	$\frac{d}{dx}(2^x) = \ln 2 \cdot 2^x$ $\ln 2 \cdot 2^x + 2y \frac{dy}{dx} = 2y + 2x \frac{dy}{dx}$ <p>Substituting (3, 2)</p> $8 \ln 2 + 4 \frac{dy}{dx} = 4 + 6 \frac{dy}{dx}$ $\frac{dy}{dx} = 4 \ln 2 - 2$ <p>Accept exact equivalents</p>	<p>B1</p> <p>M1 A1= A1</p> <p>M1</p> <p>M1 A1 (7)</p> <p>[7]</p>

4. A curve C has parametric equations

$$x = \sin^2 t, \quad y = 2 \tan t, \quad 0 \leq t < \frac{\pi}{2}$$

(a) Find $\frac{dy}{dx}$ in terms of t .

(4)

The tangent to C at the point where $t = \frac{\pi}{3}$ cuts the x -axis at the point P .

(b) Find the x -coordinate of P .

(6)

[illegible]

Question Number	Scheme	Marks
4.	<p>(a) $\frac{dx}{dt} = 2 \sin t \cos t, \frac{dy}{dt} = 2 \sec^2 t$</p> <p>$\frac{dy}{dx} = \frac{\sec^2 t}{\sin t \cos t} \left(= \frac{1}{\sin t \cos^3 t} \right)$ or equivalent</p> <p>(b) At $t = \frac{\pi}{3}, x = \frac{3}{4}, y = 2\sqrt{3}$</p> <p>$\frac{dy}{dx} = \frac{\sec^2 \frac{\pi}{3}}{\sin \frac{\pi}{3} \cos \frac{\pi}{3}} = \frac{16}{\sqrt{3}}$</p> <p>$y - 2\sqrt{3} = \frac{16}{\sqrt{3}} \left(x - \frac{3}{4} \right)$</p> <p>$y = 0 \Rightarrow x = \frac{3}{8}$</p>	<p>B1 B1</p> <p>M1 A1 (4)</p> <p>B1</p> <p>M1 A1</p> <p>M1</p> <p>M1 A1 (6)</p> <p>[10]</p>

5.

$$\frac{2x^2+5x-10}{(x-1)(x+2)} \equiv A + \frac{B}{x-1} + \frac{C}{x+2}$$

- (a) Find the values of the constants A , B and C .

(4)

- (b) Hence, or otherwise, expand $\frac{2x^2+5x-10}{(x-1)(x+2)}$ in ascending powers of x , as far as the term in x^2 . Give each coefficient as a simplified fraction.

(7)

[illegible]

Question Number	Scheme	Marks
5.	<p>(a) $A = 2$</p> $2x^2 + 5x - 10 = A(x-1)(x+2) + B(x+2) + C(x-1)$ <p>$x \rightarrow 1 \quad -3 = 3B \Rightarrow B = -1$</p> <p>$x \rightarrow -2 \quad -12 = -3C \Rightarrow C = 4$</p> <p>(b) $\frac{2x^2 + 5x - 10}{(x-1)(x+2)} = 2 + (1-x)^{-1} + 2\left(1 + \frac{x}{2}\right)^{-1}$</p> $(1-x)^{-1} = 1 + x + x^2 + \dots$ $\left(1 + \frac{x}{2}\right)^{-1} = 1 - \frac{x}{2} + \frac{x^2}{4} + \dots$ $\frac{2x^2 + 5x - 10}{(x-1)(x+2)} = (2+1+2) + (1-1)x + \left(1 + \frac{1}{2}\right)x^2 + \dots$ $= 5 + \dots \quad \text{ft their } A - B + \frac{1}{2}C$ $= \dots + \frac{3}{2}x^2 + \dots \quad 0x \text{ stated or implied}$	<p>B1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>M1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1 ft</p> <p>A1 A1 (7)</p> <p>[11]</p>

Leave
blank

$$f(\theta) = 4 \cos^2 \theta - 3 \sin^2 \theta$$

(3)

(7)



Question Number	Scheme	Marks
6.	<p>(a)</p> $f(\theta) = 4\cos^2\theta - 3\sin^2\theta$ $= 4\left(\frac{1}{2} + \frac{1}{2}\cos 2\theta\right) - 3\left(\frac{1}{2} - \frac{1}{2}\cos 2\theta\right)$ $= \frac{1}{2} + \frac{7}{2}\cos 2\theta \quad *$	<p>M1 M1</p> <p>A1 (3)</p> <p>cs0</p>
	<p>(b)</p> $\int \theta \cos 2\theta d\theta = \frac{1}{2}\theta \sin 2\theta - \frac{1}{2} \int \sin 2\theta d\theta$ $= \frac{1}{2}\theta \sin 2\theta + \frac{1}{4}\cos 2\theta$ $\int \theta f(\theta) d\theta = \frac{1}{4}\theta^2 + \frac{7}{4}\theta \sin 2\theta + \frac{7}{8}\cos 2\theta$ $\left[\dots \right]_0^{\frac{\pi}{2}} = \left[\frac{\pi^2}{16} + 0 - \frac{7}{8} \right] - \left[0 + 0 + \frac{7}{8} \right]$ $= \frac{\pi^2}{16} - \frac{7}{4}$	<p>M1 A1</p> <p>A1</p> <p>M1 A1</p> <p>M1</p> <p>A1 (7)</p> <p>[10]</p>

The line l_2 has equation $\mathbf{r} = \begin{pmatrix} 0 \\ 9 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix}$, where μ is a scalar parameter.

(a) the coordinates of C .

(3)

The point A is the point on l_1 where $\lambda = 0$ and the point B is the point on l_2 where $\mu = -1$.

(b) Find the size of the angle ACB . Give your answer in degrees to 2 decimal places.

(4)

(c) Hence, or otherwise, find the area of the triangle ABC .

(5)



Question Number	Scheme	Marks
7.	<p>(a) j components $3 + 2\lambda = 9 \Rightarrow \lambda = 3$ ($\mu = 1$) Leading to $C : (5, 9, -1)$ accept vector forms</p> <p>(b) Choosing correct directions or finding \overrightarrow{AC} and \overrightarrow{BC} $\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix} = 5 + 2 = \sqrt{6}\sqrt{29} \cos \angle ACB$ use of scalar product $\angle ACB = 57.95^\circ$ awrt 57.95°</p> <p>(c) $A : (2, 3, -4)$ $B : (-5, 9, -5)$ $\overrightarrow{AC} = \begin{pmatrix} 3 \\ 6 \\ 3 \end{pmatrix}$, $\overrightarrow{BC} = \begin{pmatrix} 10 \\ 0 \\ 4 \end{pmatrix}$ $AC^2 = 3^2 + 6^2 + 3^2 \Rightarrow AC = 3\sqrt{6}$ $BC^2 = 10^2 + 4^2 \Rightarrow BC = 2\sqrt{29}$ $\Delta ABC = \frac{1}{2} AC \times BC \sin \angle ACB$ $= \frac{1}{2} 3\sqrt{6} \times 2\sqrt{29} \sin \angle ACB \approx 33.5$ $15\sqrt{5}$, awrt 34</p>	<p>M1 A1 A1 (3)</p> <p>M1 M1 A1 A1 (4)</p> <p>M1 A1 A1 M1 A1 (5) [12]</p>
	<p><i>Alternative method for (b) and (c)</i></p> <p>(b) $A : (2, 3, -4)$ $B : (-5, 9, -5)$ $C : (5, 9, -1)$ $AB^2 = 7^2 + 6^2 + 1^2 = 86$ $AC^2 = 3^2 + 6^2 + 3^2 = 54$ $BC^2 = 10^2 + 0^2 + 4^2 = 116$ Finding all three sides $\cos \angle ACB = \frac{116 + 54 - 86}{2\sqrt{116}\sqrt{54}} (= 0.53066 \dots)$ $\angle ACB = 57.95^\circ$ awrt 57.95°</p> <p>If this method is used some of the working may gain credit in part (c) and appropriate marks may be awarded if there is an attempt at part (c).</p>	<p>M1 M1 A1 A1 (4)</p>




Figure 2

Figure 2 shows a cylindrical water tank. The diameter of a circular cross-section of the tank is 6 m. Water is flowing into the tank at a constant rate of $0.48\pi \text{ m}^3 \text{ min}^{-1}$. At time t minutes, the depth of the water in the tank is h metres. There is a tap at a point T at the bottom of the tank. When the tap is open, water leaves the tank at a rate of $0.6\pi h \text{ m}^3 \text{ min}^{-1}$.

- (a) Show that t minutes after the tap has been opened

$$75 \frac{dh}{dt} = (4 - 5h) \quad (5)$$

When $t = 0$, $h = 0.2$

- (b) Find the value of t when $h = 0.5$ (6)



Question Number	Scheme	Marks
8.	<p>(a)</p> $\frac{dV}{dt} = 0.48\pi - 0.6\pi h$ $V = 9\pi h \Rightarrow \frac{dV}{dt} = 9\pi \frac{dh}{dt}$ $9\pi \frac{dh}{dt} = 0.48\pi - 0.6\pi h$ <p>Leading to $75 \frac{dh}{dt} = 4 - 5h$ *</p> <p>(b)</p> $\int \frac{75}{4-5h} dh = \int 1 dt$ <p>separating variables</p> $-15 \ln(4-5h) = t (+C)$ $-15 \ln(4-5h) = t + C$ <p>When $t = 0$, $h = 0.2$</p> $-15 \ln 3 = C$ $t = 15 \ln 3 - 15 \ln(4-5h)$ <p>When $h = 0.5$</p> $t = 15 \ln 3 - 15 \ln 1.5 = 15 \ln \left(\frac{3}{1.5} \right) = 15 \ln 2$ <p>awrt 10.4</p> <p>Alternative for last 3 marks</p> $t = \left[-15 \ln(4-5h) \right]_{0.2}^{0.5}$ $= -15 \ln 1.5 + 15 \ln 3$ $= 15 \ln \left(\frac{3}{1.5} \right) = 15 \ln 2$ <p>awrt 10.4</p>	<p>M1 A1</p> <p>B1</p> <p>M1</p> <p>cs0 A1 (5)</p> <p>M1</p> <p>M1 A1</p> <p>M1</p> <p>M1 A1</p> <p>M1 M1</p> <p>A1 (6)</p>