

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	6	/	0	1	Signature	

Paper Reference(s)

**6666/01**

# Edexcel GCE

# Core Mathematics C4

## Advanced

Wednesday 18 June 2014 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

### Materials required for examination

---

Mathematical Formulae (Pink)

### Items included with question papers

Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.**

### Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer for each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

## Advice to Candidates

---

---

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.  
©2014 Pearson Education Ltd

Printer's Log. No.

Printer's Log. No.  
**P43165A**

W850/R6666/57570 5/5/5/1/1/1/



*Turn over*

PEARSON

1. A curve  $C$  has the equation

$$x^3 + 2xy - x - y^3 - 20 = 0$$

- (a) Find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .

(5)

- (b) Find an equation of the tangent to  $C$  at the point  $(3, -2)$ , giving your answer in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers.

(2)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme		Marks
1.	$x^3 + 2xy - x - y^3 - 20 = 0$		
(a)	<div><math display="block">\left\{ \begin{array}{c} \cancel{\frac{dx}{dy}} \\ \cancel{\frac{dy}{dx}} \end{array} \right\} \times \left\{ \begin{array}{c} \cancel{\frac{dy}{dx}} \\ \cancel{\frac{dx}{dy}} \end{array} \right\} \quad \underline{3x^2 + \left( 2y + 2x \frac{dy}{dx} \right) - 1 - 3y^2 \frac{dy}{dx} = 0}</math><math display="block">3x^2 + 2y - 1 + \left( 2x - 3y^2 \right) \frac{dy}{dx} = 0</math><math display="block">\frac{dy}{dx} = \frac{3x^2 + 2y - 1}{3y^2 - 2x} \quad \text{or} \quad \frac{1 - 3x^2 - 2y}{2x - 3y^2}</math></div>		M1 <u>A1</u> <u>B1</u>  dM1  A1 <b>cso</b>  <b>[5]</b>
(b)	<div>At <math>P(3, -2)</math>, <math>m(\mathbf{T}) = \frac{dy}{dx} = \frac{3(3)^2 + 2(-2) - 1}{3(-2)^2 - 2(3)}; = \frac{22}{6}</math> or <math>\frac{11}{3}</math>  <b>and either</b> <math>\mathbf{T}: y - -2 = \frac{11}{3} (x - 3)</math>  <b>or</b> <math>(-2) = \left( \frac{11}{3} \right) (3) + c \Rightarrow c = \dots,</math></div> <div><b>T:</b> <math>11x - 3y - 39 = 0</math> or <math>K(11x - 3y - 39) = 0</math></div>		<div>see notes</div> M1          A1 <b>cso</b>
			<b>[2]</b> 7
(a)	<div><b><u>Alternative method for part (a)</u></b>  <math display="block">\left\{ \begin{array}{c} \cancel{\frac{dx}{dy}} \\ \cancel{\frac{dy}{dx}} \end{array} \right\} \times \left\{ \begin{array}{c} \cancel{\frac{dy}{dx}} \\ \cancel{\frac{dx}{dy}} \end{array} \right\} \quad \underline{3x^2 \frac{dx}{dy} + \left( 2y \frac{dx}{dy} + 2x \right) - \frac{dx}{dy} - 3y^2 = 0}</math><math display="block">2x - 3y^2 + \left( 3x^2 + 2y - 1 \right) \frac{dx}{dy} = 0</math><math display="block">\frac{dy}{dx} = \frac{3x^2 + 2y - 1}{3y^2 - 2x} \quad \text{or} \quad \frac{1 - 3x^2 - 2y}{2x - 3y^2}</math></div>		M1 <u>A1</u> <u>B1</u>  dM1  A1 <b>cso</b>  <b>[5]</b>
<b>Question 1 Notes</b>			
(a) <b>General</b>	<b>Note</b>	Writing down $\frac{dy}{dx} = \frac{3x^2 + 2y - 1}{3y^2 - 2x}$ or $\frac{1 - 3x^2 - 2y}{2x - 3y^2}$ from <b>no working</b> is full marks.	
	<b>Note</b>	Writing down $\frac{dy}{dx} = \frac{3x^2 + 2y - 1}{2x - 3y^2}$ or $\frac{1 - 3x^2 - 2y}{3y^2 - 2x}$ from <b>no working</b> is M1A0B0M1A0	
	<b>Note</b>	Few candidates will write $3x^2 + 2y + 2x dy - 1 - 3y^2 dy = 0$ leading to $\frac{dy}{dx} = \frac{3x^2 + 2y - 1}{3y^2 - 2x}$ , o.e.  This should get full marks.	
1. (a)	<b>M1</b>	Differentiates implicitly to include either $2x \frac{dy}{dx}$ or $-y^3 \rightarrow \pm k y^2 \frac{dy}{dx}$ . (Ignore $\left( \frac{dy}{dx} = \right)$ ).	
	<b>A1</b>	$x^3 \rightarrow 3x^2$ <b>and</b> $-x - y^3 - 20 = 0 \rightarrow -1 - 3y^2 \frac{dy}{dx} = 0$	
	<b>B1</b>	$2xy \rightarrow 2y + 2x \frac{dy}{dx}$	
	<b>Note</b>	If an extra term appears then award 1 <sup>st</sup> A0.	

1. (a) ctd	<p><b>Note</b></p> <p><b>dM1</b></p> <p><b>Note</b></p> <p><b>A1</b></p>	$3x^2 + 2y + 2x \frac{dy}{dx} - 1 - 3y^2 \frac{dy}{dx} \rightarrow 3x^2 + 2y - 1 = 3y^2 \frac{dy}{dx} - 2x \frac{dy}{dx}$ <p>will get 1<sup>st</sup> A1 (implied) as the "= 0" can be implied by rearrangement of their equation.</p> <p><b>dependent on the first method mark being awarded.</b></p> <p>An attempt to factorise out <b>all the terms in</b> <math>\frac{dy}{dx}</math> as long as there are <b>at least two terms</b> in <math>\frac{dy}{dx}</math>.</p> <p>ie. <math>\dots + (2x - 3y^2) \frac{dy}{dx} = \dots</math></p> <p>Placing an extra <math>\frac{dy}{dx}</math> at the beginning and then including it in their factorisation is fine for dM1.</p> <p>For <math>\frac{1 - 2y - 3x^2}{2x - 3y^2}</math> or equivalent. Eg: <math>\frac{3x^2 + 2y - 1}{3y^2 - 2x}</math></p> <p><b>cso:</b> If the candidate's solution is not completely correct, then do not give this mark.</p> <p><b>isw:</b> You can, however, ignore subsequent working following on from correct solution.</p>
1. (b)	<p><b>M1</b></p> <p><b>Note</b></p> <p><b>A1</b></p> <p><b>cso</b></p> <p><b>isw</b></p>	<p><b>Some</b> attempt to substitute <b>both</b> <math>x = 3</math> <b>and</b> <math>y = -2</math> into their <math>\frac{dy}{dx}</math> which contains both <math>x</math> and <math>y</math> to find <math>m_T</math> <b>and</b></p> <ul style="list-style-type: none"> <li><b>either</b> applies <math>y - -2 = (\text{their } m_T)(x - 3)</math>, where <math>m_T</math> is a numerical value.</li> <li><b>or</b> finds <math>c</math> by solving <math>(-2) = (\text{their } m_T)(3) + c</math>, where <math>m_T</math> is a numerical value.</li> </ul> <p>Using a changed gradient (i.e. applying <math>\frac{-1}{\text{their } \frac{dy}{dx}}</math> or <math>\frac{1}{\text{their } \frac{dy}{dx}}</math> is M0).</p> <p>Accept any integer multiple of <math>11x - 3y - 39 = 0</math> or <math>11x - 39 - 3y = 0</math> or <math>-11x + 3y + 39 = 0</math>, where their tangent equation is equal to 0.</p> <p>A correct solution is required from a correct <math>\frac{dy}{dx}</math>.</p> <p>You can ignore subsequent working following a correct solution.</p>
1. (a)	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>B1</b></p> <p><b>dM1</b></p> <p><b>A1</b></p>	<p><b><u>Alternative method for part (a): Differentiating with respect to y</u></b></p> <p>Differentiates implicitly to include either <math>2y \frac{dx}{dy}</math> or <math>x^3 \rightarrow \pm kx^2 \frac{dx}{dy}</math> or <math>-x \rightarrow -\frac{dx}{dy}</math> (Ignore <math>\left(\frac{dx}{dy} = \right)</math>).</p> <p><math>x^3 \rightarrow 3x^2 \frac{dx}{dy}</math> <b>and</b> <math>-x - y^3 - 20 = 0 \rightarrow -\frac{dx}{dy} - 3y^2 = 0</math></p> <p><math>2xy \rightarrow 2y \frac{dx}{dy} + 2x</math></p> <p><b>dependent on the first method mark being awarded.</b></p> <p>An attempt to factorise out <b>all the terms in</b> <math>\frac{dx}{dy}</math> as long as there are <b>at least two terms</b> in <math>\frac{dx}{dy}</math>.</p> <p>For <math>\frac{1 - 2y - 3x^2}{2x - 3y^2}</math> or equivalent. Eg: <math>\frac{3x^2 + 2y - 1}{3y^2 - 2x}</math></p> <p><b>cso:</b> If the candidate's solution is not completely correct, then do not give this mark.</p>

Leave  
blank

- $$1 - 6x + Ax^2 + \dots$$

- (2)

- (3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
2.	$\left\{ (1 + kx)^{-4} = 1 + (-4)(kx) + \frac{(-4)(-4-1)}{2!}(kx)^2 + \dots \right\}$	
(a)	<p><b>Either</b> <math>(-4)k = -6</math> <b>or</b> <math>(1 + kx)^{-4} = 1 + (-4)(kx)</math> <b>see notes</b></p> <p>leading to <math>k = \frac{3}{2}</math> <span style="float: right;"><math>k = \frac{3}{2}</math> or 1.5 or <math>\frac{6}{4}</math></span></p>	M1 A1 [2]
(b)	<p><math>\frac{(-4)(-5)}{2}(k)^2</math> <b>Either</b> <math>\frac{(-4)(-5)}{2!}</math> <b>or</b> <math>(k)^2</math> <b>or</b> <math>(kx)^2</math></p> <p><b>Either</b> <math>\frac{(-4)(-5)}{2!}(k)^2</math> <b>or</b> <math>\frac{(-4)(-5)}{2!}(kx)^2</math></p> <p><math>\left\{ A = \frac{(-4)(-5)}{2!} \left( \frac{3}{2} \right)^2 \right\} \Rightarrow A = \frac{45}{2}</math> <span style="float: right;"><math>\frac{45}{2}</math> or 22.5</span></p>	M1 M1 A1 [3] 5
<b>Question 2 Notes</b>		
<b>Note</b>	<b>In this question ignore part labelling and mark part (a) and part (b) together.</b>	
<b>Note</b>	Writing down $\left\{ (1 + kx)^{-4} \right\} = 1 + (-4)(kx) + \frac{(-4)(-4-1)}{2!}(kx)^2 + \dots$ gets all the method marks in Q2. i.e. (a) M1 and (b) M1M1	
(a)	<p><b>M1</b> Award M1 for</p> <ul style="list-style-type: none"> <li><b>either</b> writing down <math>(-4)k = -6</math> or <math>4k = 6</math></li> <li><b>or</b> expanding <math>(1 + kx)^{-4}</math> to give <math>1 + (-4)(kx)</math></li> <li><b>or</b> writing down <math>(-4)kx = -6</math> or <math>(-4k) = -6x</math> or <math>-4kx = -6x</math></li> </ul> <p><b>A1</b> <math>k = \frac{3}{2}</math> or 1.5 or <math>\frac{6}{4}</math> <b>from no incorrect sign errors.</b></p>	
<b>Note</b>	The M1 mark can be implied by a candidate writing down the correct value of $k$ .	
<b>Note</b>	Award M1 for writing down $4k = 6$ and then A1 for $k = 1.5$ (or equivalent).	
<b>Note</b>	Award M0 for $4k = -6$ (if there is no evidence that $(1 + kx)^{-4}$ expands to give $1 + (-4)(kx) + \dots$ )	
<b>Note</b>	$1 + (-4)(kx)$ leading to $(-4)k = 6$ leading to $k = \frac{3}{2}$ is M1A0.	
(b)	<p><b>M1</b> For <b>either</b> <math>\frac{(-4)(-4-1)}{2!}</math> <b>or</b> <math>\frac{(-4)(-5)}{2!}</math> <b>or</b> 10 <b>or</b> <math>(k)^2</math> <b>or</b> <math>(kx)^2</math></p> <p><b>M1</b> <b>Either</b> <math>\frac{(-4)(-4-1)}{2!}(k)^2</math> <b>or</b> <math>\frac{(-4)(-5)}{2!}(k)^2</math> <b>or</b> <math>\frac{(-4)(-5)}{2!}(kx)^2</math> <b>or</b> <math>\frac{(-4)(-5)}{2!}(\text{their } k)^2</math> <b>or</b> <math>10k^2</math></p>	
<b>Note</b>	Candidates are allowed to use 2 instead of 2!	
<b>A1</b>	Uses $k = 1.5$ to give $A = \frac{45}{2}$ or 22.5	
<b>Note</b>	$A = \frac{90}{4}$ which has not been simplified is A0.	
<b>Note</b>	Award A0 for $A = \frac{45}{2}x^2$ .	
<b>Note</b>	Allow A1 for $A = \frac{45}{2}x^2$ followed by $A = \frac{45}{2}$	
<b>Note</b>	$k = -1.5$ leading to $A = \frac{45}{2}$ or 22.5 is A0.	

3.




Figure 1

Figure 1 shows a sketch of part of the curve with equation  $y = \frac{10}{2x + 5\sqrt{x}}$ ,  $x > 0$

The finite region  $R$ , shown shaded in Figure 1, is bounded by the curve, the  $x$ -axis, and the lines with equations  $x = 1$  and  $x = 4$

The table below shows corresponding values of  $x$  and  $y$  for  $y = \frac{10}{2x + 5\sqrt{x}}$

$x$	1	2	3	4
$y$	1.42857	0.90326		0.55556

- Complete the table above by giving the missing value of  $y$  to 5 decimal places. (1)
- Use the trapezium rule, with all the values of  $y$  in the completed table, to find an estimate for the area of  $R$ , giving your answer to 4 decimal places. (3)
- By reference to the curve in Figure 1, state, giving a reason, whether your estimate in part (b) is an overestimate or an underestimate for the area of  $R$ . (1)
- Use the substitution  $u = \sqrt{x}$ , or otherwise, to find the exact value of


$$\int_1^4 \frac{10}{2x + 5\sqrt{x}} dx$$

(6)




Question Number	Scheme					Marks
3.	$\frac{x}{y}$	1	2	3	4	$y = \frac{10}{2x + 5\sqrt{x}}$
(a)	{ At $x = 3,$ } $y = 0.68212$ (5 dp)					0.68212 B1 cao [1]
(b)	$\frac{1}{2} \times 1 \times [1.42857 + 0.55556 + 2(0.90326 + \text{their } 0.68212)]$					Outside brackets $\frac{1}{2} \times 1$ or $\frac{1}{2}$ B1 aef For structure of [.....] M1
(c)	{ = $\frac{1}{2}(5.15489)$ } = 2.577445 = 2.5774 (4 dp)					anything that rounds to 2.5774 A1 [3]
(d)	<ul style="list-style-type: none"><li>Overestimate</li><li>and a reason such as<ul style="list-style-type: none"><li>{ top of } <u>trapezia lie above the curve</u></li><li>a diagram which gives reference to the extra area</li><li>concave or convex</li><li><math>\frac{d^2y}{dx^2} &gt; 0</math> (can be implied)</li><li>bends inwards</li><li>curves downwards</li></ul></li></ul>					B1 [1]
	$\{u = \sqrt{x} \Rightarrow \} \frac{du}{dx} = \frac{1}{2}x^{-\frac{1}{2}} \text{ or } \frac{dx}{du} = 2u$ $\int \frac{10}{2u^2 + 5u} \cdot 2u \, du$ Either $\left\{ \int \right\} \frac{\pm k u}{\alpha u^2 \pm \beta u} \{du\}$ or $\left\{ \int \right\} \frac{\pm k}{u(\alpha u^2 \pm \beta u)} \{du\}$					M1
	$\left\{ = \int \frac{20}{2u + 5} \, du \right\} = \frac{20}{2} \ln(2u + 5)$ $\pm \lambda \ln(2u + 5)$ or $\pm \lambda \ln\left(u + \frac{5}{2}\right), \lambda \neq 0$ <b>with no other terms.</b> $\frac{20}{2u + 5} \rightarrow \frac{20}{2} \ln(2u + 5)$ or $10 \ln\left(u + \frac{5}{2}\right)$					M1 A1 cso
	$\left\{ \left[ \frac{20}{2} \ln(2u + 5) \right]_1^2 \right\} = 10 \ln(2(2) + 5) - 10 \ln(2(1) + 5)$ $10 \ln 9 - 10 \ln 7$ or $10 \ln\left(\frac{9}{7}\right)$ or $20 \ln 3 - 10 \ln 7$					Substitutes limits of 2 and 1 in $u$ (or 4 and 1 in $x$ ) and subtracts the correct way round. M1 A1 oe cso [6] 11
Question 3 Notes						
3. (a)	B1	0.68212 correct answer only. Look for this on the table or in the candidate's working.				
(b)	B1	Outside brackets $\frac{1}{2} \times 1$ or $\frac{1}{2}$ or equivalent.				
	M1	For structure of trapezium rule [ ..... ]				
	Note A1	No errors are allowed [eg. an omission of a y-ordinate or an extra y-ordinate or a repeated y ordinate]. anything that rounds to 2.5774				
	Note	Working must be seen to demonstrate the use of the trapezium rule. (Actual area is 2.51314428...)				



3. (b) contd	<p><b>Note</b> Award B1M1A1 for <math>\frac{1}{2}(1.42857 + 0.55556) + (0.90326 + \text{their } 0.68212) = 2.577445</math></p> <p><b>Bracketing mistake:</b> Unless the final answer implies that the calculation has been done correctly award B1M0A0 for <math>\frac{1}{2} \times 1 + 1.42857 + 2(0.90326 + \text{their } 0.68212) + 0.55556</math> (nb: answer of 5.65489).</p> <p>award B1M0A0 for <math>\frac{1}{2} \times 1 (1.42857 + 0.55556) + 2(0.90326 + \text{their } 0.68212)</math> (nb: answer of 4.162825).</p> <p><b>Alternative method: Adding individual trapezia</b></p> $\text{Area} \approx 1 \times \left[ \frac{1.42857 + 0.90326}{2} + \frac{0.90326 + "0.68212"}{2} + \frac{"0.68212" + 0.55556}{2} \right] = 2.577445$ <p><b>B1</b> B1: 1 and a divisor of 2 on all terms inside brackets.</p> <p><b>M1</b> M1: First and last ordinates once and two of the middle ordinates twice inside brackets ignoring the 2.</p> <p><b>A1</b> A1: anything that rounds to 2.5774</p>
(c)	<p><b>B1</b> Overestimate <b>and</b> either trapezia lie above curve <b>or</b> a diagram that gives reference to the extra area</p> <p><b>eg.</b> This diagram is sufficient. It must show the top of a trapezium lying above the curve.</p>  <p><b>or</b> concave or convex <b>or</b> <math>\frac{d^2y}{dx^2} &gt; 0</math> (can be implied) <b>or</b> bends inwards <b>or</b> curves downwards.</p> <p><b>Note</b> Reason of "gradient is negative" by itself is B0.</p>
(d)	<p><b>B1</b> <math>\frac{du}{dx} = \frac{1}{2}x^{-\frac{1}{2}}</math> or <math>du = \frac{1}{2\sqrt{x}} dx</math> or <math>2\sqrt{x} du = dx</math> or <math>dx = 2u du</math> or <math>\frac{dx}{du} = 2u</math> o.e.</p> <p><b>M1</b> Applying the substitution and achieving <math>\left\{ \int \right\} \frac{\pm ku}{\alpha u^2 \pm \beta u} \{du\}</math> or <math>\left\{ \int \right\} \frac{\pm k}{u(\alpha u^2 \pm \beta u)} \{du\}</math>, <math>k, \alpha, \beta \neq 0</math>. Integral sign and <math>du</math> not required for this mark.</p> <p><b>M1</b> Cancelling <math>u</math> and integrates to achieve <math>\pm \lambda \ln(2u + 5)</math> or <math>\pm \lambda \ln\left(u + \frac{5}{2}\right)</math>, <math>\lambda \neq 0</math> <b>with no other terms</b>.</p> <p><b>A1</b> <b>cso.</b> Integrates <math>\frac{20}{2u + 5}</math> to give <math>\frac{20}{2} \ln(2u + 5)</math> or <math>10 \ln\left(u + \frac{5}{2}\right)</math>, un-simplified or simplified.</p> <p><b>Note</b> BE CAREFUL! Candidates must be integrating <math>\frac{20}{2u + 5}</math> or equivalent.</p> <p>So <math>\int \frac{10}{2u + 5} du = 10 \ln(2u + 5)</math> WOULD BE A0 and final A0.</p> <p><b>M1</b> Applies limits of 2 and 1 in <math>u</math> or 4 and 1 in <math>x</math> in their (i.e. any) changed function and subtracts the correct way round.</p> <p><b>A1</b> Exact answers of either <math>10 \ln 9 - 10 \ln 7</math> or <math>10 \ln\left(\frac{9}{7}\right)</math> or <math>20 \ln 3 - 10 \ln 7</math> or <math>20 \ln\left(\frac{3}{\sqrt{7}}\right)</math> or <math>\ln\left(\frac{9^{10}}{7^{10}}\right)</math> or equivalent. <b>Correct solution only.</b></p> <p><b>Note</b> You can ignore subsequent working which follows from a correct answer.</p> <p><b>Note</b> A decimal answer of 2.513144283... (without a correct <b>exact</b> answer) is A0.</p>

4.



### Figure 2

A vase with a circular cross-section is shown in Figure 2. Water is flowing into the vase.

When the depth of the water is  $h$  cm, the volume of water  $V$  cm<sup>3</sup> is given by

$$V = 4\pi h(h + 4), \quad 0 \leq h \leq 25$$

Water flows into the vase at a constant rate of  $80\pi \text{ cm}^3 \text{ s}^{-1}$

Find the rate of change of the depth of the water, in  $\text{cm s}^{-1}$ , when  $h = 6$


(5)



Question Number	Scheme	Marks
4.	$\frac{dV}{dt} = 80\pi$ , $V = 4\pi h(h + 4) = 4\pi h^2 + 16\pi h$ , $\frac{dV}{dh} = 8\pi h + 16\pi$	$\pm \alpha h \pm \beta$ , $\alpha \neq 0$ , $\beta \neq 0$ $8\pi h + 16\pi$ M1 A1
	$\left\{ \frac{dV}{dh} \times \frac{dh}{dt} = \frac{dV}{dt} \Rightarrow \right\} (8\pi h + 16\pi) \frac{dh}{dt} = 80\pi$ $\left( \text{Candidate's } \frac{dV}{dh} \right) \times \frac{dh}{dt} = 80\pi$ $\left\{ \frac{dh}{dt} = \frac{dV}{dt} \div \frac{dV}{dh} \Rightarrow \right\} \frac{dh}{dt} = 80\pi \times \frac{1}{8\pi h + 16\pi}$ or $80\pi \div \text{Candidate's } \frac{dV}{dh}$	M1 oe A1 oe
	When $h = 6$ , $\left\{ \frac{dh}{dt} = \right\} \frac{1}{8\pi(6) + 16\pi} \times 80\pi \left\{ = \frac{80\pi}{64\pi} \right\}$ $\frac{dh}{dt} = \underline{1.25} \text{ (cms}^{-1}\text{)}$	<b>dependent on the previous M1 see notes</b> $1.25$ or $\frac{5}{4}$ or $\frac{10}{8}$ or $\frac{80}{64}$ dM1 A1 oe <b>[5]</b> <b>5</b>
	<b><u>Alternative Method for the first M1A1</u></b> Product rule: $\begin{cases} u = 4\pi h & v = h + 4 \\ \frac{du}{dh} = 4\pi & \frac{dv}{dh} = 1 \end{cases}$ $\frac{dV}{dh} = 4\pi(h + 4) + 4\pi h$	$\pm \alpha h \pm \beta$ , $\alpha \neq 0$ , $\beta \neq 0$ $4\pi(h + 4) + 4\pi h$ M1 A1
<b>Question 4 Notes</b>		
	<b>M1</b>	An expression of the form $\pm \alpha h \pm \beta$ , $\alpha \neq 0$ , $\beta \neq 0$ . <b>Can be simplified or un-simplified.</b>
	<b>A1</b>	Correct simplified or un-simplified differentiation of $V$ . eg. $8\pi h + 16\pi$ or $4\pi(h + 4) + 4\pi h$ or $8\pi(h + 2)$ or equivalent.
	<b>Note</b>	Some candidates will use the product rule to differentiate $V$ with respect to $h$ . <b>(See Alt Method 1).</b>
	<b>Note</b>	$\frac{dV}{dh}$ does not have to be explicitly stated, but it should be clear that they are differentiating their $V$ .
	<b>M1</b>	$\left( \text{Candidate's } \frac{dV}{dh} \right) \times \frac{dh}{dt} = 80\pi$ or $80\pi \div \text{Candidate's } \frac{dV}{dh}$
	<b>Note</b>	Also allow 2 <sup>nd</sup> M1 for $\left( \text{Candidate's } \frac{dV}{dh} \right) \times \frac{dh}{dt} = \mathbf{80}$ or $\mathbf{80} \div \text{Candidate's } \frac{dV}{dh}$
	<b>Note</b>	Give 2 <sup>nd</sup> M0 for $\left( \text{Candidate's } \frac{dV}{dh} \right) \times \frac{dh}{dt} = \mathbf{80\pi t}$ or $\mathbf{80k}$ or $\mathbf{80\pi t}$ or $\mathbf{80k} \div \text{Candidate's } \frac{dV}{dh}$
	<b>dM1</b>	<b>which is dependent on the previous M1 mark.</b> Substitutes $h = 6$ into an expression which is a result of a quotient of their $\frac{dV}{dh}$ and $80\pi$ (or 80)
	<b>A1</b>	$1.25$ or $\frac{5}{4}$ or $\frac{10}{8}$ or $\frac{80}{64}$ (units are not required).
	<b>Note</b>	$\frac{80\pi}{64\pi}$ as a final answer is A0.
	<b>Note</b>	Substituting $h = 6$ into a correct $\frac{dV}{dh}$ gives $64\pi$ but the final M1 mark can only be awarded if this is used as a quotient with $80\pi$ (or 80)

Leave  
blank

5.



### Figure 3

Figure 3 shows a sketch of the curve  $C$  with parametric equations

$$x = 4 \cos\left(t + \frac{\pi}{6}\right), \quad y = 2 \sin t, \quad 0 \leq t < 2\pi$$

(a) Show that

$$x + y = 2\sqrt{3} \cos t \quad (3)$$

(b) Show that a cartesian equation of  $C$  is

$$(x + y)^2 + ay^2 = b$$

where  $a$  and  $b$  are integers to be determined.

(2)



Question Number	Scheme	Marks
5.	$x = 4 \cos\left(t + \frac{\pi}{6}\right), \quad y = 2 \sin t$	
(a)	<p><b>Main Scheme</b></p> $x = 4 \left( \cos t \cos\left(\frac{\pi}{6}\right) - \sin t \sin\left(\frac{\pi}{6}\right) \right) \qquad \cos\left(t + \frac{\pi}{6}\right) \rightarrow \cos t \cos\left(\frac{\pi}{6}\right) \pm \sin t \sin\left(\frac{\pi}{6}\right)$ <p>So, <math>\{x + y\} = 4 \left( \cos t \cos\left(\frac{\pi}{6}\right) - \sin t \sin\left(\frac{\pi}{6}\right) \right) + 2 \sin t</math> <span style="float: right;">Adds their expanded <math>x</math> (which is in terms of <math>t</math>) to <math>2 \sin t</math></span></p> $= 4 \left( \left( \frac{\sqrt{3}}{2} \right) \cos t - \left( \frac{1}{2} \right) \sin t \right) + 2 \sin t$ $= 2\sqrt{3} \cos t \quad *$ <p style="text-align: right;">Correct proof</p>	<p>M1 oe</p> <p>dM1</p> <p>A1 * <b>[3]</b></p>
(a)	<p><b>Alternative Method 1</b></p> $x = 4 \left( \cos t \cos\left(\frac{\pi}{6}\right) - \sin t \sin\left(\frac{\pi}{6}\right) \right) \qquad \cos\left(t + \frac{\pi}{6}\right) \rightarrow \cos t \cos\left(\frac{\pi}{6}\right) \pm \sin t \sin\left(\frac{\pi}{6}\right)$ $= 4 \left( \left( \frac{\sqrt{3}}{2} \right) \cos t - \left( \frac{1}{2} \right) \sin t \right) = 2\sqrt{3} \cos t - 2 \sin t$ <p>So, <math>x = 2\sqrt{3} \cos t - y</math> <span style="float: right;">Forms an equation in <math>x, y</math> and <math>t</math>.</span></p> $x + y = 2\sqrt{3} \cos t \quad *$ <p style="text-align: right;">Correct proof</p>	<p>M1 oe</p> <p>dM1</p> <p>A1 * <b>[3]</b></p>
(b)	<p><b>Main Scheme</b></p> $\left( \frac{x+y}{2\sqrt{3}} \right)^2 + \left( \frac{y}{2} \right)^2 = 1$ $\Rightarrow \frac{(x+y)^2}{12} + \frac{y^2}{4} = 1$ $\Rightarrow (x+y)^2 + 3y^2 = 12$ <p style="text-align: right;">Applies <math>\cos^2 t + \sin^2 t = 1</math> to achieve an equation containing <b>only</b> <math>x</math>'s and <math>y</math>'s.</p> <p style="text-align: right;"><math>(x+y)^2 + 3y^2 = 12</math> <math>\{a=3, b=12\}</math></p>	<p>M1</p> <p>A1 <b>[2]</b></p>
(b)	<p><b>Alternative Method 1</b></p> $(x+y)^2 = 12 \cos^2 t = 12(1 - \sin^2 t) = 12 - 12 \sin^2 t$ <p>So, <math>(x+y)^2 = 12 - 3y^2</math> <span style="float: right;">Applies <math>\cos^2 t + \sin^2 t = 1</math> to achieve an equation containing <b>only</b> <math>x</math>'s and <math>y</math>'s.</span></p> $\Rightarrow (x+y)^2 + 3y^2 = 12$ <p style="text-align: right;"><math>(x+y)^2 + 3y^2 = 12</math></p>	<p>M1</p> <p>A1 <b>[2]</b></p>
(b)	<p><b>Alternative Method 2</b></p> $(x+y)^2 = 12 \cos^2 t$ <p>As <math>12 \cos^2 t + 12 \sin^2 t = 12</math></p> <p>then <math>(x+y)^2 + 3y^2 = 12</math></p>	<p>M1, A1 <b>[2]</b></p>
		<b>5</b>

Question 5 Notes		
5. (a)	M1	$\cos\left(t + \frac{\pi}{6}\right) \rightarrow \cos t \cos\left(\frac{\pi}{6}\right) \pm \sin t \sin\left(\frac{\pi}{6}\right)$ or $\cos\left(t + \frac{\pi}{6}\right) \rightarrow \left(\frac{\sqrt{3}}{2}\right)\cos t \pm \left(\frac{1}{2}\right)\sin t$
	Note	If a candidate states $\cos(A + B) = \cos A \cos B \pm \sin A \sin B$ , but there is an error <i>in its application</i> then give M1.  <u>Awarding the dM1 mark which is dependent on the first method mark</u>
Main	dM1	Adds their expanded $x$ (which is in terms of $t$ ) to $2\sin t$
	Note	Writing $x + y = \dots$ is not needed in the <b>Main Scheme</b> method.
Alt 1	dM1	Forms an equation in $x$ , $y$ and $t$ .
(b)	A1*	Evidence of $\cos\left(\frac{\pi}{6}\right)$ and $\sin\left(\frac{\pi}{6}\right)$ evaluated and the proof is correct with no errors.
	Note	$\{x + y\} = 4\cos\left(t + \frac{\pi}{6}\right) + 2\sin t$ , by itself is M0M0A0.
	M1	Applies $\cos^2 t + \sin^2 t = 1$ to achieve an equation containing <b>only</b> $x$ 's and $y$ 's.
	A1	leading $(x + y)^2 + 3y^2 = 12$
	SC	Award <b>Special Case B1B0</b> for a candidate who writes down <b>either</b> <ul style="list-style-type: none"> <li><math>(x + y)^2 + 3y^2 = 12</math> from no working</li> <li><math>a = 3, b = 12</math>, but <b><u>does not provide a correct proof.</u></b></li> </ul>
	Note	Alternative method 2 is fine for M1 A1
	Note	Writing $(x + y)^2 = 12\cos^2 t$ followed by $12\cos^2 t + a(4\sin^2 t) = b \Rightarrow a = 3, b = 12$ is SC: B1B0
	Note	Writing $(x + y)^2 = 12\cos^2 t$ followed by $12\cos^2 t + a(4\sin^2 t) = b$ <ul style="list-style-type: none"> <li>states <math>a = 3, b = 12</math></li> <li>and refers to either <math>\cos^2 t + \sin^2 t = 1</math> or <math>12\cos^2 t + 12\sin^2 t = 12</math></li> <li>and there is no incorrect working</li> </ul> would get M1A1

**6. (i) Find**

(3)

(ii) Find

(2)

(7)

Question Number	Scheme	Marks
6. (i)	$\int x e^{4x} dx = \frac{1}{4} x e^{4x} - \int \frac{1}{4} e^{4x} \{dx\}$ $= \frac{1}{4} x e^{4x} - \frac{1}{16} e^{4x} \{+ c\}$	<div> <math>\pm \alpha x e^{4x} - \int \beta e^{4x} \{dx\}, \alpha \neq 0, \beta &gt; 0</math> M1         </div> <div> <math>\frac{1}{4} x e^{4x} - \int \frac{1}{4} e^{4x} \{dx\}</math> A1         </div> <div> <math>\frac{1}{4} x e^{4x} - \frac{1}{16} e^{4x}</math> A1         </div>
(ii)	$\int \frac{8}{(2x-1)^3} dx = \frac{8(2x-1)^{-2}}{(2)(-2)} \{+ c\}$ $\{-2(2x-1)^{-2} \{+ c\}\}$	<div> <math>\pm \lambda (2x-1)^{-2}</math> M1         </div> <div> <math>\frac{8(2x-1)^{-2}}{(2)(-2)}</math> or equivalent. A1         </div> <div> <i>{Ignore subsequent working}.</i> [2]         </div>
(iii)	$\frac{dy}{dx} = e^x \operatorname{cosec} 2y \operatorname{cosec} y \quad y = \frac{\pi}{6} \text{ at } x = 0$	
	<p><b>Main Scheme</b></p> $\int \frac{1}{\operatorname{cosec} 2y \operatorname{cosec} y} dy = \int e^x dx \quad \text{or} \quad \int \sin 2y \sin y dy = \int e^x dx$ $\int 2 \sin y \cos y \sin y dy = \int e^x dx$ $\frac{2}{3} \sin^3 y = e^x \{+ c\}$ $\frac{2}{3} \sin^3 \left( \frac{\pi}{6} \right) = e^0 + c \quad \text{or} \quad \frac{2}{3} \left( \frac{1}{8} \right) - 1 = c$ $\left\{ \Rightarrow c = -\frac{11}{12} \right\} \quad \text{giving} \quad \frac{2}{3} \sin^3 y = e^x - \frac{11}{12}$	<div>B1 oe</div> <div>M1</div> <div>M1</div> <div>A1</div> <div>B1</div> <div>M1</div> <div>A1</div> <div>[7]</div>
	<p><b>Alternative Method 1</b></p> $\int \frac{1}{\operatorname{cosec} 2y \operatorname{cosec} y} dy = \int e^x dx \quad \text{or} \quad \int \sin 2y \sin y dy = \int e^x dx$ $\int -\frac{1}{2} (\cos 3y - \cos y) dy = \int e^x dx$ $-\frac{1}{2} \left( \frac{1}{3} \sin 3y - \sin y \right) = e^x \{+ c\}$ $-\frac{1}{2} \left( \frac{1}{3} \sin \left( \frac{3\pi}{6} \right) - \sin \left( \frac{\pi}{6} \right) \right) = e^0 + c \quad \text{or} \quad -\frac{1}{2} \left( \frac{1}{3} - \frac{1}{2} \right) - 1 = c$ $\left\{ \Rightarrow c = -\frac{11}{12} \right\} \quad \text{giving} \quad -\frac{1}{6} \sin 3y + \frac{1}{2} \sin y = e^x - \frac{11}{12}$	<div>B1 oe</div> <div>M1</div> <div>M1</div> <div>A1</div> <div>B1</div> <div>M1</div> <div>A1</div> <div>[7]</div>
		12



Question 6 Notes		
6. (i)	<b>M1</b>	Integration by parts is applied in the form $\pm \alpha x e^{4x} - \int \beta e^{4x} \{dx\}$ , where $\alpha \neq 0$ , $\beta > 0$ . (must be in this form).
	<b>A1</b>	$\frac{1}{4} x e^{4x} - \int \frac{1}{4} e^{4x} \{dx\}$ or equivalent.
	<b>A1</b>	$\frac{1}{4} x e^{4x} - \frac{1}{16} e^{4x}$ with/without $+ c$ . Can be un-simplified.
	<b>isw</b>	You can ignore subsequent working following on from a correct solution.
	<b>SC</b>	<b>SPECIAL CASE:</b> A candidate who uses $u = x$ , $\frac{dv}{dx} = e^{4x}$ , writes down the correct “by parts” formula, but makes only one error when applying it can be awarded Special Case M1.
(ii)	<b>M1</b>	$\pm \lambda (2x - 1)^{-2}$ , $\lambda \neq 0$ . <b>Note</b> that $\lambda$ can be 1.
	<b>A1</b>	$\frac{8(2x - 1)^{-2}}{(2)(-2)}$ or $-2(2x - 1)^{-2}$ or $\frac{-2}{(2x - 1)^2}$ with/without $+ c$ . Can be un-simplified.
	<b>Note</b>	You can ignore subsequent working which follows from a correct answer.
(iii)	<b>B1</b>	Separates variables as shown. $dy$ and $dx$ should be in the correct positions, though this mark can be implied by later working. Ignore the integral signs.
	<b>Note</b>	<b>Allow B1 for</b> $\int \frac{1}{\csc 2y \csc y} = \int e^x$ <b>or</b> $\int \sin 2y \sin y = \int e^x$
	<b>M1</b>	$\frac{1}{\csc 2y} \rightarrow 2 \sin y \cos y$ <b>or</b> $\sin 2y \rightarrow 2 \sin y \cos y$ <b>or</b> $\sin 2y \sin y \rightarrow \pm \lambda \cos 3y \pm \lambda \cos y$ seen anywhere in the candidate’s working to (iii).
	<b>M1</b>	Integrates to give $\pm \mu \sin^3 y$ , $\mu \neq 0$ <b>or</b> $\pm \alpha \sin 3y \pm \beta \sin y$ , $\alpha \neq 0$ , $\beta \neq 0$
	<b>A1</b>	$2 \sin^2 y \cos y \rightarrow \frac{2}{3} \sin^3 y$ (with no extra terms) <b>or</b> integrates to give $-\frac{1}{2} \left( \frac{1}{3} \sin 3y - \sin y \right)$
	<b>B1</b>	Evidence that $e^x$ has been integrated to give $e^x$ <b>as part of solving their DE</b> .
	<b>M1</b>	Some evidence of using both $y = \frac{\pi}{6}$ and $x = 0$ in an integrated or changed equation containing $c$ .
	<b>Note</b>	that is mark can be implied by the correct value of $c$ .
	<b>A1</b>	$\frac{2}{3} \sin^3 y = e^x - \frac{11}{12}$ <b>or</b> $-\frac{1}{6} \sin 3y + \frac{1}{2} \sin y = e^x - \frac{11}{12}$ <b>or any equivalent correct answer.</b>
	<b>Note</b>	You can ignore subsequent working which follows from a correct answer.
<b>Alternative Method 2 (Using integration by parts twice)</b>		
$\int \sin 2y \sin y dy = \int e^x dx$		B1 oe
		Applies integration by parts <b>twice</b> to give $\pm \alpha \cos y \sin 2y \pm \beta \sin y \cos 2y$ <b>M2</b>
$\frac{1}{3} \cos y \sin 2y - \frac{2}{3} \sin y \cos 2y = e^x \{ + c \}$		$\frac{1}{3} \cos y \sin 2y - \frac{2}{3} \sin y \cos 2y$ (simplified or un-simplified) <b>A1</b>
		$e^x \rightarrow e^x$ <b>as part of solving their DE.</b> <b>B1</b>
		as in the main scheme <b>M1</b>
$\frac{1}{3} \cos y \sin 2y - \frac{2}{3} \sin y \cos 2y = e^x - \frac{11}{12}$		$-\frac{1}{6} \sin 3y + \frac{1}{2} \sin y = e^x - \frac{11}{12}$ <b>A1</b>
<b>[7]</b>		

7.




Figure 4

Figure 4 shows a sketch of part of the curve  $C$  with parametric equations

$$x = 3 \tan \theta, \quad y = 4 \cos^2 \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The point  $P$  lies on  $C$  and has coordinates  $(3, 2)$ .

The line  $l$  is the normal to  $C$  at  $P$ . The normal cuts the  $x$ -axis at the point  $Q$ .

(a) Find the  $x$  coordinate of the point  $Q$ .

(6)

The finite region  $S$ , shown shaded in Figure 4, is bounded by the curve  $C$ , the  $x$ -axis, the  $y$ -axis and the line  $l$ . This shaded region is rotated  $2\pi$  radians about the  $x$ -axis to form a solid of revolution.

(b) Find the exact value of the volume of the solid of revolution, giving your answer in the form  $p\pi + q\pi^2$ , where  $p$  and  $q$  are rational numbers to be determined.

[You may use the formula  $V = \frac{1}{3}\pi r^2 h$  for the volume of a cone.]

(9)

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
7.	$x = 3 \tan \theta$ , $y = 4 \cos^2 \theta$ or $y = 2 + 2 \cos 2\theta$ , $0 \leq \theta < \frac{\pi}{2}$ .	
(a)	$\frac{dx}{d\theta} = 3 \sec^2 \theta$ , $\frac{dy}{d\theta} = -8 \cos \theta \sin \theta$ or $\frac{dy}{d\theta} = -4 \sin 2\theta$	
	$\frac{dy}{dx} = \frac{-8 \cos \theta \sin \theta}{3 \sec^2 \theta} \left\{ = -\frac{8}{3} \cos^3 \theta \sin \theta = -\frac{4}{3} \sin 2\theta \cos^2 \theta \right\}$ their $\frac{dy}{d\theta}$ divided by their $\frac{dx}{d\theta}$ Correct $\frac{dy}{dx}$	M1 A1 oe
	At $P(3, 2)$ , $\theta = \frac{\pi}{4}$ , $\frac{dy}{dx} = -\frac{8}{3} \cos^3 \left( \frac{\pi}{4} \right) \sin \left( \frac{\pi}{4} \right) \left\{ = -\frac{2}{3} \right\}$ Some evidence of substituting $\theta = \frac{\pi}{4}$ into their $\frac{dy}{dx}$ So, $m(N) = \frac{3}{2}$ applies $m(N) = \frac{-1}{m(T)}$	M1 M1
	Either N: $y - 2 = \frac{3}{2} (x - 3)$ or $2 = \left( \frac{3}{2} \right) (3) + c$ see notes	M1
	{ At Q, $y = 0$ , so, $-2 = \frac{3}{2} (x - 3)$ } giving $x = \frac{5}{3}$ $x = \frac{5}{3}$ or $1 \frac{2}{3}$ or awrt 1.67	A1 cso
(b)	$\left\{ \int y^2 dx = \int y^2 \frac{dx}{d\theta} d\theta \right\} = \left\{ \int \right\} (4 \cos^2 \theta)^2 3 \sec^2 \theta \{d\theta\}$ see notes So, $\pi \int y^2 dx = \pi \int (4 \cos^2 \theta)^2 3 \sec^2 \theta \{d\theta\}$ see notes $\int y^2 dx = \int 48 \cos^2 \theta d\theta$ $\int 48 \cos^2 \theta \{d\theta\}$ $= \{48\} \int \left( \frac{1 + \cos 2\theta}{2} \right) d\theta \left\{ = \int (24 + 24 \cos 2\theta) d\theta \right\}$ Applies $\cos 2\theta = 2 \cos^2 \theta - 1$	M1 A1 A1 M1
	$= \{48\} \left( \frac{1}{2} \theta + \frac{1}{4} \sin 2\theta \right) \left\{ = 24\theta + 12 \sin 2\theta \right\}$ Dependent on the first method mark. For $\pm \alpha \theta \pm \beta \sin 2\theta$ $\cos^2 \theta \rightarrow \left( \frac{1}{2} \theta + \frac{1}{4} \sin 2\theta \right)$	dM1 A1
	$\int_0^{\frac{\pi}{4}} y^2 dx \left\{ = 48 \left[ \frac{1}{2} \theta + \frac{1}{4} \sin 2\theta \right]_0^{\frac{\pi}{4}} \right\} = \{48\} \left( \left( \frac{\pi}{8} + \frac{1}{4} \right) - (0 + 0) \right) \{ = 6\pi + 12 \}$ Dependent on the third method mark. { So $V = \pi \int_0^{\frac{\pi}{4}} y^2 dx = 6\pi^2 + 12\pi$ } $V_{\text{cone}} = \frac{1}{3} \pi (2)^2 \left( 3 - \frac{5}{3} \right) \left\{ = \frac{16\pi}{9} \right\}$ $V_{\text{cone}} = \frac{1}{3} \pi (2)^2 (3 - \text{their } (a))$ $\left\{ \text{Vol}(S) = 6\pi^2 + 12\pi - \frac{16\pi}{9} \right\} \Rightarrow \text{Vol}(S) = \frac{92}{9} \pi + 6\pi^2$ $\frac{92}{9} \pi + 6\pi^2$ $\left\{ p = \frac{92}{9}, q = 6 \right\}$	dM1 M1 A1
		[9]

[6]

[9]

15

Question 7 Notes		
7. (a)	<b>1<sup>st</sup> M1</b>	Applies their $\frac{dy}{d\theta}$ divided by their $\frac{dx}{d\theta}$ <b>or</b> applies $\frac{dy}{d\theta}$ multiplied by their $\frac{d\theta}{dx}$
	<b>SC</b>	Award <b>Special Case 1<sup>st</sup> M1</b> if <b>both</b> $\frac{dx}{d\theta}$ <b>and</b> $\frac{dy}{d\theta}$ are both correct.
	<b>1<sup>st</sup> A1</b>	Correct $\frac{dy}{dx}$ i.e. $\frac{-8\cos\theta\sin\theta}{3\sec^2\theta}$ or $-\frac{8}{3}\cos^3\theta\sin\theta$ or $-\frac{4}{3}\sin 2\theta\cos^2\theta$ or any equivalent form.
	<b>2<sup>nd</sup> M1</b>	<b>Some evidence</b> of substituting $\theta = \frac{\pi}{4}$ or $\theta = 45^\circ$ into their $\frac{dy}{dx}$
	<b>Note</b>	For 3 <sup>rd</sup> M1 and 4 <sup>th</sup> M1, $m(\mathbf{T})$ must be found by using $\frac{dy}{dx}$ .
	<b>3<sup>rd</sup> M1</b>	applies $m(\mathbf{N}) = \frac{-1}{m(\mathbf{T})}$ . Numerical value for $m(\mathbf{N})$ is required here.
	<b>4<sup>th</sup> M1</b>	<ul style="list-style-type: none"> <li>Applies <math>y - 2 = (\text{their } m_N)(x - 3)</math>, where <math>m(\mathbf{N})</math> is a numerical value,</li> <li>or <b>finds c</b> by solving <math>2 = (\text{their } m_N)3 + c</math>, where <math>m(\mathbf{N})</math> is a numerical value,</li> </ul> and $m_N = -\frac{1}{\text{their } m(\mathbf{T})}$ or $m_N = \frac{1}{\text{their } m(\mathbf{T})}$ or $m_N = -\text{their } m(\mathbf{T})$ .
	<b>Note</b>	This mark can be implied by subsequent working.
	<b>2<sup>nd</sup> A1</b>	$x = \frac{5}{3}$ or $1\frac{2}{3}$ or awrt 1.67 <b>from a correct solution only</b> .
	<b>1<sup>st</sup> M1</b>	Applying $\int y^2 dx$ as $y^2 \frac{dx}{d\theta}$ with their $\frac{dx}{d\theta}$ . Ignore $\pi$ or $\frac{1}{3}\pi$ outside integral.
(b)	<b>Note</b>	You can ignore the omission of an integral sign and/or $d\theta$ for the 1 <sup>st</sup> M1.
	<b>Note</b>	Allow 1 <sup>st</sup> M1 for $\int (\cos^2\theta)^2 \times \text{"their } 3\sec^2\theta"$ $d\theta$ or $\int 4(\cos^2\theta)^2 \times \text{"their } 3\sec^2\theta"$ $d\theta$
	<b>1<sup>st</sup> A1</b>	Correct expression $\left\{ \pi \int y^2 dx \right\} = \pi \int (4\cos^2\theta)^2 3\sec^2\theta \{d\theta\}$ (Allow the omission of $d\theta$ )
	<b>Note</b>	<b>IMPORTANT:</b> The $\pi$ can be recovered later, <b>but as a correct statement only</b> .
	<b>2<sup>nd</sup> A1</b>	$\left\{ \int y^2 dx \right\} = \int 48\cos^2\theta \{d\theta\}$ . (Ignore $d\theta$ ). <b>Note:</b> 48 can be written as 24(2) for example.
	<b>2<sup>nd</sup> M1</b>	Applies $\cos 2\theta = 2\cos^2\theta - 1$ to their integral. (Seen or <b>implied</b> .)
	<b>3<sup>rd</sup> dM1*</b>	<b>which is dependent on the 1<sup>st</sup> M1 mark.</b> Integrating $\cos^2\theta$ to give $\pm\alpha\theta \pm \beta\sin 2\theta$ , $\alpha \neq 0$ , $\beta \neq 0$ , un-simplified or simplified.
	<b>3<sup>rd</sup> A1</b>	<b>which is dependent on the 3<sup>rd</sup> M1 mark and the 1<sup>st</sup> M1 mark.</b> Integrating $\cos^2\theta$ to give $\frac{1}{2}\theta + \frac{1}{4}\sin 2\theta$ , un-simplified or simplified.  This can be implied by $k\cos^2\theta$ giving $\frac{k}{2}\theta + \frac{k}{4}\sin 2\theta$ , un-simplified or simplified.
	<b>4<sup>th</sup> dM1</b>	<b>which is dependent on the 3<sup>rd</sup> M1 mark and the 1<sup>st</sup> M1 mark.</b> Some evidence of applying limits of $\frac{\pi}{4}$ and 0 (0 can be implied) to an integrated function in $\theta$
	<b>5<sup>th</sup> M1</b>	Applies $V_{\text{cone}} = \frac{1}{3}\pi(2)^2(3 - \text{their part (a) answer})$ .
	<b>Note</b>	Also allow the 5 <sup>th</sup> M1 for $V_{\text{cone}} = \pi \int_{\text{their } \frac{5}{3}}^3 \left( \frac{3}{2}x - \frac{5}{2} \right)^2 \{dx\}$ , which includes the correct limits.
	<b>4<sup>th</sup> A1</b>	$\frac{92}{9}\pi + 6\pi^2$ or $10\frac{2}{9}\pi + 6\pi^2$
	<b>Note</b> <b>Note</b>	A decimal answer of 91.33168464... (without a correct <b>exact</b> answer) is A0. The $\pi$ in the volume formula is only needed for the 1 <sup>st</sup> A1 mark and the final accuracy mark.

7.		<p><b><u>Working with a Cartesian Equation</u></b></p> <p>A cartesian equation for <math>C</math> is <math>y = \frac{36}{x^2 + 9}</math></p>
(a)	<p><b>1<sup>st</sup> M1</b></p> <p><b>1<sup>st</sup> A1</b></p> <p><b>2<sup>nd</sup> dM1</b></p>	<p><math>\frac{dy}{dx} = \pm \lambda x (\pm \alpha x^2 \pm \beta)^{-2}</math> <b>or</b> <math>\frac{dy}{dx} = \frac{\pm \lambda x}{(\pm \alpha x^2 \pm \beta)^2}</math></p> <p><math>\frac{dy}{dx} = -36(x^2 + 9)^{-2}(2x)</math> <b>or</b> <math>\frac{dy}{dx} = \frac{-72x}{(x^2 + 9)^2}</math> un-simplified or simplified.</p> <p><b>Dependent on the 1<sup>st</sup> M1 mark if a candidate uses this method</b></p> <p>For substituting <math>x = 3</math> into their <math>\frac{dy}{dx}</math></p> <p>i.e. at <math>P(3, 2)</math>, <math>\frac{dy}{dx} = \frac{-72(3)}{(3^2 + 9)^2} \left\{ = -\frac{2}{3} \right\}</math></p> <p>From this point onwards the original scheme can be applied.</p>
(b)	<p><b>1<sup>st</sup> M1</b></p> <p><b>A1</b></p>	<p>For <math>\int \left( \frac{\pm \lambda}{\pm \alpha x^2 \pm \beta} \right)^2 \{dx\}</math> (<math>\pi</math> not required for this mark)</p> <p>For <math>\pi \int \left( \frac{36}{x^2 + 9} \right)^2 \{dx\}</math> (<math>\pi</math> required for this mark)</p> <p>To integrate, a substitution of <math>x = 3 \tan \theta</math> is required which will lead to <math>\int 48 \cos^2 \theta d\theta</math> and so from this point onwards the original scheme can be applied.</p>
(a)	<p><b>1<sup>st</sup> M1</b></p> <p><b>1<sup>st</sup> A1</b></p> <p><b>2<sup>nd</sup> dM1</b></p>	<p>Another cartesian equation for <math>C</math> is <math>x^2 = \frac{36}{y} - 9</math></p> <p><math>\pm \alpha x = \pm \frac{\beta}{y^2} \frac{dy}{dx}</math> <b>or</b> <math>\pm \alpha x \frac{dx}{dy} = \pm \frac{\beta}{y^2}</math></p> <p><math>2x = -\frac{36}{y^2} \frac{dy}{dx}</math> <b>or</b> <math>2x \frac{dx}{dy} = -\frac{36}{y^2}</math></p> <p><b>Dependent on the 1<sup>st</sup> M1 mark if a candidate uses this method</b></p> <p>For substituting <math>x = 3</math> to find <math>\frac{dy}{dx}</math></p> <p>i.e. at <math>P(3, 2)</math>, <math>2(3) = -\frac{36}{4} \frac{dy}{dx} \Rightarrow \frac{dy}{dx} = \dots</math></p> <p>From this point onwards the original scheme can be applied.</p>

8. Relative to a fixed origin  $O$ , the point  $A$  has position vector  $\begin{pmatrix} -2 \\ 4 \\ 7 \end{pmatrix}$

and the point  $B$  has position vector  $\begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix}$

The line  $l_1$  passes through the points  $A$  and  $B$ .

(a) Find the vector  $\overrightarrow{AB}$ . (2)

(b) Hence find a vector equation for the line  $l_1$  (1)

The point  $P$  has position vector  $\begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix}$

Given that angle  $PBA$  is  $\theta$ ,

(c) show that  $\cos \theta = \frac{1}{3}$  (3)

The line  $l_2$  passes through the point  $P$  and is parallel to the line  $l_1$

(d) Find a vector equation for the line  $l_2$  (2)

The points  $C$  and  $D$  both lie on the line  $l_2$

Given that  $AB = PC = DP$  and the  $x$  coordinate of  $C$  is positive,

(e) find the coordinates of  $C$  and the coordinates of  $D$ . (3)

(f) find the exact area of the trapezium  $ABCD$ , giving your answer as a simplified surd. (4)

---

---

---

---

---

---

---


---

---

---



Question Number	Scheme	Marks
8.	$\overrightarrow{OA} = -2\mathbf{i} + 4\mathbf{j} + 7\mathbf{k}$ , $\overrightarrow{OB} = -\mathbf{i} + 3\mathbf{j} + 8\mathbf{k}$ & $\overrightarrow{OP} = 0\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$	
(a)	$\overrightarrow{AB} = \pm((- \mathbf{i} + 3\mathbf{j} + 8\mathbf{k}) - (-2\mathbf{i} + 4\mathbf{j} + 7\mathbf{k})); = \mathbf{i} - \mathbf{j} + \mathbf{k}$	M1; A1 [2]
(b)	$\{l_1 : \mathbf{r}\} = \begin{pmatrix} -2 \\ 4 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ or $\{\mathbf{r}\} = \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$	B1ft [1]
(c)	$\overrightarrow{PB} = \overrightarrow{OB} - \overrightarrow{OP} = \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} - \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix}$ or $\overrightarrow{BP} = \begin{pmatrix} 1 \\ -1 \\ -5 \end{pmatrix}$	M1
	$\{\cos \theta = \frac{\overrightarrow{AB} \bullet \overrightarrow{PB}}{ \overrightarrow{AB}   \overrightarrow{PB} } = \frac{\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \bullet \begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix}}{\sqrt{(1)^2 + (-1)^2 + (1)^2} \cdot \sqrt{(-1)^2 + (1)^2 + (5)^2}}$ Applies dot product formula between their $(\overrightarrow{AB}$ or $\overrightarrow{BA})$ and their $(\overrightarrow{PB}$ or $\overrightarrow{BP})$ .	M1
	$\{\cos \theta\} = \frac{-1-1+5}{\sqrt{3} \cdot \sqrt{27}} = \frac{3}{9} = \frac{1}{3}$ Correct proof	A1 cso [3]
(d)	$\{l_2 : \mathbf{r} = \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ $\mathbf{p} + \lambda \mathbf{d}$ or $\mathbf{p} + \mu \mathbf{d}$ , $\mathbf{p} \neq 0$ , $\mathbf{d} \neq 0$ with either $\mathbf{p} = 0\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ or $\mathbf{d} =$ their $\overrightarrow{AB}$ , or a multiple of their $\overrightarrow{AB}$ . Correct vector equation.	M1 A1 ft [2]
(e)	$\overrightarrow{OC} = \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 4 \end{pmatrix}$ or $\overrightarrow{OD} = \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} = \begin{pmatrix} -1 \\ 3 \\ 2 \end{pmatrix}$ $\{C(1, 1, 4), D(-1, 3, 2)\}$ Either $\overrightarrow{OP} +$ their $\overrightarrow{AB}$ or $\overrightarrow{OP} -$ their $\overrightarrow{AB}$ At least one set of coordinates are correct. Both sets of coordinates are correct.	M1 A1 ft A1 ft [3]
(f) Way 1	$\frac{h}{\sqrt{(-1)^2 + (1)^2 + (5)^2}} = \sin \theta$ $h = \sqrt{27} \sin(70.5\dots) \left\{ = \sqrt{27} \frac{\sqrt{8}}{3} = 2\sqrt{6} = \text{awrt } 4.9 \right\}$ Area $ABCD = \frac{1}{2} 2\sqrt{6}(\sqrt{3} + 2\sqrt{3})$ $\left\{ = \frac{1}{2} 2\sqrt{6}(3\sqrt{3}) = 3\sqrt{18} \right\} = 9\sqrt{2}$	$\frac{h}{\text{their }  \overrightarrow{PB} } = \sin \theta$ $\sqrt{27} \sin(70.5\dots)$ or $\sqrt{27} \cdot \frac{\sqrt{8}}{3}$ or $2\sqrt{6}$ or awrt 4.9 or equivalent $\frac{1}{2}(\text{their } h)(\text{their } AB + \text{their } CD)$ $9\sqrt{2}$ M1 A1 oe dM1 A1 cao [4] 15

<p>8. (f)</p>	<p><b>Helpful Diagram!</b></p>  <p>Area <math>\triangle APB = 4.2426\dots</math></p> <p><math>\vec{DA} = \vec{PB} = \begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix}</math></p> <p><math>\vec{PA} = \vec{CB} = \begin{pmatrix} -2 \\ 2 \\ 4 \end{pmatrix}</math></p>
	<p><math>\vec{PA} = \vec{CB} = \begin{pmatrix} -2 \\ 2 \\ 4 \end{pmatrix}</math> and <math>\vec{AB} = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math>, so <math>BC \perp AB</math></p> <p>Candidates do not need to prove this result for part (f)</p>
<p>8. (f) Way 2</p>	<p><math>h =  \vec{CB}  = \sqrt{(-2)^2 + (2)^2 + (4)^2} = \sqrt{24} = 2\sqrt{6} = 4.8989\dots</math></p> <p>Area <math>ABCD = \frac{1}{2} \sqrt{24} (\sqrt{3} + 2\sqrt{3})</math> or <math>\frac{1}{2} \sqrt{24} \sqrt{3} + \sqrt{24} \sqrt{3}</math></p> <p><math>= 9\sqrt{2}</math></p> <p>Attempts <math> \vec{PA} </math> or <math> \vec{CB} </math>  <math> \vec{PA}  =  \vec{CB}  = \sqrt{24}</math>  <math>\frac{1}{2} h(\text{their } AB + \text{their } CD)</math>  <math>9\sqrt{2}</math></p> <p>M1 A1 oe dM1 oe A1 cso</p>
<p>Way3 8. (f)</p>	<p><b>Finds the area of either triangle <math>APB</math> or <math>APD</math> or <math>BCP</math> and triples the result.</b></p> <p>Area <math>\triangle APB = \frac{1}{2} \sqrt{3} (3\sqrt{3}) \sin \theta</math></p> <p><math>= \frac{1}{2} \sqrt{3} (3\sqrt{3}) \sin(70.5\dots)</math></p> <p>Area <math>ABCD = 3 (3\sqrt{2})</math></p> <p><math>= 9\sqrt{2}</math></p> <p>Attempts <math>\frac{1}{2} (\text{their } AB)(\text{their } PB) \sin \theta</math>  <math>\frac{1}{2} \sqrt{3} (3\sqrt{3}) \sin(70.5\dots)</math> or <math>3\sqrt{2}</math>  or awrt 4.24 or equivalent  <math>3 \times \text{Area of } \triangle APB</math>  <math>9\sqrt{2}</math></p> <p>M1 A1 dM1 A1 cso</p>



Question 8 Notes	
8. (a)	<p><b>M1</b> Finding the difference (either way) between <math>\overrightarrow{OB}</math> and <math>\overrightarrow{OA}</math>. If no “subtraction” seen, you can award M1 for 2 out of 3 correct components of the difference.</p> <p><b>A1</b> <math>\mathbf{i} - \mathbf{j} + \mathbf{k}</math> or <math>\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math> or <math>(1, -1, 1)</math> or benefit of the doubt <math>\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math></p>
(b)	<p><b>B1ft</b> <math>\{\mathbf{r}\} = \begin{pmatrix} -2 \\ 4 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math> or <math>\{\mathbf{r}\} = \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math>, with <math>\overrightarrow{AB}</math> or <math>\overrightarrow{BA}</math> correctly followed through from (a).</p> <p><b>Note</b> <math>\mathbf{r} =</math> is not needed.</p>
(c)	<p><b>M1</b> An attempt to find either the vector <math>\overrightarrow{PB}</math> or <math>\overrightarrow{BP}</math>. If no “subtraction” seen, you can award M1 for 2 out of 3 correct components of the difference.</p> <p><b>M1</b> Applies dot product formula between their <math>(\overrightarrow{AB}</math> or <math>\overrightarrow{BA})</math> and their <math>(\overrightarrow{PB}</math> or <math>\overrightarrow{BP})</math>.</p> <p><b>A1</b> Obtains <math>\{\cos \theta\} = \frac{1}{3}</math> <b>by correct solution only.</b></p> <p><b>Note</b> If candidate starts by applying <math>\frac{\overrightarrow{AB} \cdot \overrightarrow{PB}}{ \overrightarrow{AB}   \overrightarrow{PB} }</math> correctly (without reference to <math>\cos \theta = \dots</math>) they can gain both 2<sup>nd</sup> M1 and A1 mark.</p> <p><b>Note</b> Award the final A1 mark if candidate achieves <math>\{\cos \theta\} = \frac{1}{3}</math> by either taking the dot product between</p> <p>(i) <math>\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math> and <math>\begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix}</math> or (ii) <math>\begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix}</math> and <math>\begin{pmatrix} 1 \\ -1 \\ -5 \end{pmatrix}</math>. Ignore if any of these vectors are labelled incorrectly.</p> <p><b>Note</b> Award final A0, cso for those candidates who take the dot product between</p> <p>(iii) <math>\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}</math> and <math>\begin{pmatrix} 1 \\ -1 \\ -5 \end{pmatrix}</math> or (iv) <math>\begin{pmatrix} -1 \\ 1 \\ -1 \end{pmatrix}</math> and <math>\begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix}</math></p> <p>They will usually find <math>\{\cos \theta\} = -\frac{1}{3}</math> or may fudge <math>\{\cos \theta\} = \frac{1}{3}</math>.</p> <p>If these candidates give a convincing detailed explanation which must include reference to the direction of their vectors then this can be given A1 cso</p>
(c)	<p><b>Alternative Method 1: The Cosine Rule</b></p> <p><math>\overrightarrow{PB} = \overrightarrow{OB} - \overrightarrow{OP} = \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} - \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix}</math> or <math>\overrightarrow{BP} = \begin{pmatrix} 1 \\ -1 \\ -5 \end{pmatrix}</math></p> <p>Note <math> \overrightarrow{PB}  = \sqrt{27}</math>, <math> \overrightarrow{AB}  = \sqrt{3}</math> and <math> \overrightarrow{PA}  = \sqrt{24}</math></p> <p><math>(\sqrt{24})^2 = (\sqrt{27})^2 + (\sqrt{3})^2 - 2(\sqrt{27})(\sqrt{3})\cos \theta</math></p> <p><math>\cos \theta = \frac{27 + 3 - 24}{18} = \frac{1}{3}</math></p> <p>Mark in the same way as the main scheme. M1</p> <p>Applies the cosine rule the correct way round M1 oe</p> <p>Correct proof A1 cso</p> <p>[3]</p>

8. (c)	<b>Alternative Method 2: Right-Angled Trigonometry</b>		
	$\overrightarrow{PB} = \overrightarrow{OB} - \overrightarrow{OP} = \begin{pmatrix} -1 \\ 3 \\ 8 \end{pmatrix} - \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 5 \end{pmatrix} \text{ or } \overrightarrow{BP} = \begin{pmatrix} 1 \\ -1 \\ -5 \end{pmatrix}$	Mark in the same way as the main scheme.	M1
	Either $(\sqrt{24})^2 + (\sqrt{3})^2 = (\sqrt{27})^2$		
	or $\overrightarrow{AB} \bullet \overrightarrow{PA} = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \bullet \begin{pmatrix} -2 \\ 2 \\ 4 \end{pmatrix} = -2 - 2 + 4 = 0$	Confirms $\Delta PAB$ is right-angled	M1
	So, $\left\{ \cos \theta = \frac{AB}{PB} \Rightarrow \right\} \cos \theta = \frac{\sqrt{3}}{\sqrt{27}} = \frac{1}{3}$	Correct proof	A1 cso
<b>[3]</b>			
(d)	<b>M1</b>	Writing down a line in the form $\mathbf{p} + \lambda \mathbf{d}$ or $\mathbf{p} + \mu \mathbf{d}$ with either $\mathbf{a} = \begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix}$ or $\mathbf{d} =$ their $\overrightarrow{AB}$ $\mathbf{d} =$ their $\overrightarrow{AB}$ , or a multiple of their $\overrightarrow{AB}$ found in part (a).	
	<b>A1ft</b>	Writing $\begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} + \mu \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$ or $\begin{pmatrix} 0 \\ 2 \\ 3 \end{pmatrix} + \mu \mathbf{d}$ , where $\mathbf{d} =$ their $\overrightarrow{AB}$ or a multiple of their $\overrightarrow{AB}$ found in part (a).	
	<b>Note</b>	$\mathbf{r} =$ is not needed.	
	<b>Note</b>	Using the same scalar parameter as in part (b) is fine for A1.	
(e)	<b>M1</b>	Either $\overrightarrow{OP} +$ their $\overrightarrow{AB}$ or $\overrightarrow{OP} -$ their $\overrightarrow{AB}$ .	
	<b>Note</b>	This can be implied at least two out of three correct components for either their $C$ or their $D$ .	
	<b>A1ft</b>	At least one set of coordinates are correct. Ignore labelling of $C, D$	
	<b>A1ft</b>	Both sets of coordinates are correct. Ignore labelling of $C, D$	
	<b>Note</b>	You can follow through either or both accuracy marks in this part using their $\overrightarrow{AB}$ from part (a).	
(f)	<b>M1</b>	Way 1: $\frac{h}{\text{their }  \overrightarrow{PB} } = \sin \theta$ Way 2: Attempts $ \overrightarrow{PA} $ or $ \overrightarrow{CB} $ Way 3: Attempts $\frac{1}{2} (\text{their } PB)(\text{their } AB) \sin \theta$	
	<b>Note</b>	Finding $AD$ by itself is M0.	
	<b>A1</b>	Either <ul style="list-style-type: none"><li><math>h = \sqrt{27} \sin(70.5\dots)</math> or <math> \overrightarrow{PA}  =  \overrightarrow{CB}  = \sqrt{24}</math> or equivalent. (See Way 1 and Way 2)</li></ul> or <ul style="list-style-type: none"><li>the area of either triangle <math>APB</math> or <math>APD</math> or <math>BDP = \frac{1}{2} \sqrt{3} (3\sqrt{3}) \sin(70.5\dots)</math> o.e. (See Way 3).</li></ul>	
	<b>dM1</b>	<b>which is dependent on the 1<sup>st</sup> M1 mark.</b>	
		A full method to find the area of trapezium $ABCD$ . (See Way 1, Way 2 and Way 3).	
	<b>A1</b>	$9\sqrt{2}$ from a correct solution only.	
	<b>Note</b>	A decimal answer of 12.7279... (without a correct <b>exact</b> answer) is A0.	