

Centre No.						Paper Reference				Surname	Initial(s)			
Candidate No.						6	6	6	3	/	0	1	Signature	

Paper Reference(s)  
**6663/01**

**Edexcel GCE  
Core Mathematics C1  
Advanced Subsidiary**

Tuesday 10 January 2006 – Afternoon  
Time: 1 hour 30 minutes



**Materials required for examination**  
Mathematical Formulae (Green)

**Items included with question papers**  
Nil

Calculators may NOT be used in this examination.

Examiner's use only		
Team Leader's use only		

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
Total	

**Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature.  
Check that you have the correct question paper.  
You must write your answer for each question in the space following the question.

**Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided.  
Full marks may be obtained for answers to ALL questions.  
The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).  
There are 10 questions in this question paper. The total mark for this paper is 75.  
There are 20 pages in this question paper. Any blank pages are indicated.

**Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled.  
You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.





Question number	Scheme	Marks
1.	$x(x^2 - 4x + 3)$ $= x(x - 3)(x - 1)$	Factor of $x$ . (Allow $(x - 0)$ ) Factorise 3 term quadratic M1 M1 A1 (3) <b>Total 3 marks</b>
2.	(a) $u_2 = (-2)^2 = 4$ $u_3 = 1, u_4 = 4$ (b) $u_{20} = 4$	For $u_3$ , ft $(u_2 - 3)^2$ B1 B1ft, B1 (3) B1ft (1) <b>Total 4 marks</b>
3.	(a) $y = 5 - (2 \times 3) = -1$ (b) Gradient of $L$ is $\frac{1}{2}$ $y - (-1) = \frac{1}{2}(x - 3)$ $x - 2y - 5 = 0$	(or equivalent verification) (*) B1 (1) B1 (ft from a <u>changed</u> gradient) M1 A1ft A1 (or equiv. with integer coefficients) (4) <b>Total 5 marks</b>



Question number	Scheme	Marks
1.	$x(x^2 - 4x + 3)$ $= x(x - 3)(x - 1)$	Factor of $x$ . (Allow $(x - 0)$ ) Factorise 3 term quadratic M1 M1 A1 (3) <b>Total 3 marks</b>
2.	(a) $u_2 = (-2)^2 = 4$ $u_3 = 1, u_4 = 4$ (b) $u_{20} = 4$	For $u_3$ , ft $(u_2 - 3)^2$ B1 B1ft, B1 (3) B1ft (1) <b>Total 4 marks</b>
3.	(a) $y = 5 - (2 \times 3) = -1$ (b) Gradient of $L$ is $\frac{1}{2}$ $y - (-1) = \frac{1}{2}(x - 3)$ $x - 2y - 5 = 0$	(or equivalent verification) (*) B1 (1) B1 (ft from a <u>changed</u> gradient) M1 A1ft A1 (or equiv. with integer coefficients) (4) <b>Total 5 marks</b>

Leave  
blank

3. The line  $L$  has equation  $y = 5 - 2x$ .

(a) Show that the point  $P(3, -1)$  lies on  $L$ .

(1)

(b) Find an equation of the line perpendicular to  $L$ , which passes through  $P$ . Give your answer in the form  $ax + by + c = 0$ , where  $a, b$  and  $c$  are integers.

(4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

(Total 5 marks)

Q3



Question number	Scheme	Marks
1.	$x(x^2 - 4x + 3)$ $= x(x - 3)(x - 1)$	Factor of $x$ . (Allow $(x - 0)$ ) Factorise 3 term quadratic M1 M1 A1 (3) <b>Total 3 marks</b>
2.	(a) $u_2 = (-2)^2 = 4$ $u_3 = 1, u_4 = 4$ (b) $u_{20} = 4$	For $u_3$ , ft $(u_2 - 3)^2$ B1 B1ft, B1 (3) B1ft (1) <b>Total 4 marks</b>
3.	(a) $y = 5 - (2 \times 3) = -1$ (b) Gradient of $L$ is $\frac{1}{2}$ $y - (-1) = \frac{1}{2}(x - 3)$ $x - 2y - 5 = 0$	(or equivalent verification) (*) B1 (1) B1 (ft from a <u>changed</u> gradient) M1 A1ft A1 (or equiv. with integer coefficients) (4) <b>Total 5 marks</b>

Leave blank

4. Given that  $y = 2x^2 - \frac{6}{x^3}$ ,  $x \neq 0$ ,

(a) find  $\frac{dy}{dx}$ ,

(2)

(b) find  $\int y \, dx$ .

(3)

Horizontal lines for student answers.

(Total 5 marks)

Q4

Small empty box for marking.





Question number	Scheme	Marks
4.	<p>(a) <math>\frac{dy}{dx} = 4x + 18x^{-4}</math> M1: <math>x^2 \rightarrow x</math> or <math>x^{-3} \rightarrow x^{-4}</math></p> <p>(b) <math>\frac{2x^3}{3} - \frac{6x^{-2}}{-2} + C</math> M1: <math>x^2 \rightarrow x^3</math> or <math>x^{-3} \rightarrow x^{-2}</math> or <math>+C</math></p> <p><math>\left( = \frac{2x^3}{3} + 3x^{-2} + C \right)</math> First A1: <math>\frac{2x^3}{3} + C</math></p> <p>Second A1: <math>-\frac{6x^{-2}}{-2}</math></p>	<p>M1 A1 (2)</p> <p>M1 A1 A1 (3)</p> <p><b>Total 5 marks</b></p>

5.	<p>(a) <math>3\sqrt{5}</math> (or <math>a = 3</math>)</p> <p>(b) <math>\frac{2(3+\sqrt{5})}{(3-\sqrt{5})} \times \frac{(3+\sqrt{5})}{(3+\sqrt{5})}</math></p> <p><math>(3-\sqrt{5})(3+\sqrt{5}) = 9-5</math> (= 4) (Used as or intended as denominator)</p> <p><math>(3+\sqrt{5})(p \pm q\sqrt{5}) = \dots</math> 4 terms (<math>p \neq 0, q \neq 0</math>) (Independent)</p> <p>or <math>(6+2\sqrt{5})(p \pm q\sqrt{5}) = \dots</math> 4 terms (<math>p \neq 0, q \neq 0</math>)</p> <p>[Correct version: <math>(3+\sqrt{5})(3+\sqrt{5}) = 9+3\sqrt{5}+3\sqrt{5}+5</math>, or double this.]</p> <p><math>\frac{2(14+6\sqrt{5})}{4} = 7+3\sqrt{5}</math> 1<sup>st</sup> A1: <math>b = 7</math>, 2<sup>nd</sup> A1: <math>c = 3</math></p>	<p>B1 (1)</p> <p>M1</p> <p>B1</p> <p>M1</p> <p>A1 A1 (5)</p> <p><b>Total 6 marks</b></p>
----	---	--



Question number	Scheme	Marks
4.	<p>(a) <math>\frac{dy}{dx} = 4x + 18x^{-4}</math> M1: <math>x^2 \rightarrow x</math> or <math>x^{-3} \rightarrow x^{-4}</math></p> <p>(b) <math>\frac{2x^3}{3} - \frac{6x^{-2}}{-2} + C</math> M1: <math>x^2 \rightarrow x^3</math> or <math>x^{-3} \rightarrow x^{-2}</math> or <math>+C</math></p> <p><math>\left( = \frac{2x^3}{3} + 3x^{-2} + C \right)</math> First A1: <math>\frac{2x^3}{3} + C</math></p> <p>Second A1: <math>-\frac{6x^{-2}}{-2}</math></p>	<p>M1 A1 (2)</p> <p>M1 A1 A1 (3)</p> <p><b>Total 5 marks</b></p>

5.	<p>(a) <math>3\sqrt{5}</math> (or <math>a = 3</math>)</p> <p>(b) <math>\frac{2(3+\sqrt{5})}{(3-\sqrt{5})} \times \frac{(3+\sqrt{5})}{(3+\sqrt{5})}</math></p> <p><math>(3-\sqrt{5})(3+\sqrt{5}) = 9-5</math> (= 4) (Used as or intended as denominator)</p> <p><math>(3+\sqrt{5})(p \pm q\sqrt{5}) = \dots</math> 4 terms (<math>p \neq 0, q \neq 0</math>) (Independent)</p> <p>or <math>(6+2\sqrt{5})(p \pm q\sqrt{5}) = \dots</math> 4 terms (<math>p \neq 0, q \neq 0</math>)</p> <p>[Correct version: <math>(3+\sqrt{5})(3+\sqrt{5}) = 9+3\sqrt{5}+3\sqrt{5}+5</math>, or double this.]</p> <p><math>\frac{2(14+6\sqrt{5})}{4} = 7+3\sqrt{5}</math> 1<sup>st</sup> A1: <math>b = 7</math>, 2<sup>nd</sup> A1: <math>c = 3</math></p>	<p>B1 (1)</p> <p>M1</p> <p>B1</p> <p>M1</p> <p>A1 A1 (5)</p> <p><b>Total 6 marks</b></p>
----	---	--

6.

Figure 1




Figure 1 shows a sketch of the curve with equation  $y = f(x)$ . The curve passes through the points  $(0, 3)$  and  $(4, 0)$  and touches the  $x$ -axis at the point  $(1, 0)$ .

On separate diagrams sketch the curve with equation




(a)  $y = f(x + 1)$ , (3)

(b)  $y = 2f(x)$ , (3)

(c)  $y = f\left(\frac{1}{2}x\right)$ . (3)

On each diagram show clearly the coordinates of all the points where the curve meets the axes.



Question number	Scheme	Marks
6.	<p>(a) </p> <p>(See below) Clearly through origin (or (0, 0) seen) 3 labelled (or (3, 0) seen)</p> <p>(b) </p> <p>Stretch parallel to y-axis 1 and 4 labelled (or (1, 0) and (4, 0) seen) 6 labelled (or (0, 6) seen)</p> <p>(c) </p> <p>Stretch parallel to x-axis 2 and 8 labelled (or (2, 0) and (8, 0) seen) 3 labelled (or (0, 3) seen)</p>	<p>M1 A1 A1 (3)</p> <p>M1 A1 A1 (3)</p> <p>M1 A1 A1 (3)</p> <p><b>Total 9 marks</b></p>

7.	<p>(a) <math>500 + (500 + 200) = 1200</math> or <math>S_2 = \frac{1}{2}2\{1000 + 200\} = 1200</math> (*)</p> <p>(b) Using <math>a = 500, d = 200</math> with <math>n = 7, 8</math> or <math>9</math> <math>a + (n - 1)d</math> or “listing” <math>500 + (7 \times 200) = (\pounds)1900</math></p> <p>(c) Using <math>\frac{1}{2}n\{2a + (n - 1)d\}</math> or <math>\frac{1}{2}n\{a + l\}</math>, or listing and “summing” terms <math>S_8 = \frac{1}{2}8\{2 \times 500 + 7 \times 200\}</math> or <math>S_8 = \frac{1}{2}8\{500 + 1900\}</math>, or all terms in list correct <math>= (\pounds) 9600</math></p> <p>(d) <math>\frac{1}{2}n\{2 \times 500 + (n - 1) \times 200\} = 32000</math> M1: General <math>S_n</math>, equated to 32000 <math>n^2 + 4n - 320 = 0</math> (or equiv.) M1: Simplify to 3 term quadratic <math>(n + 20)(n - 16) = 0</math> <math>n = \dots</math> M1: Attempt to solve 3 t.q. <math>n = 16,</math> Age is 26 A1cso,A1cso</p>	<p>B1 (1)</p> <p>M1 A1 (2)</p> <p>M1 A1 A1 (3)</p> <p>M1 A1 M1 A1 M1 A1cso,A1cso (7)</p> <p><b>Total 13 marks</b></p>
----	---	---

Leave  
blank

7. On Alice's 11th birthday she started to receive an annual allowance. The first annual allowance was £500 and on each following birthday the allowance was increased by £200.

(a) Show that, immediately after her 12th birthday, the total of the allowances that Alice had received was £1200. (1)

(b) Find the amount of Alice's annual allowance on her 18th birthday. (2)

(c) Find the total of the allowances that Alice had received up to and including her 18th birthday. (3)

When the total of the allowances that Alice had received reached £32 000 the allowance stopped.

(d) Find how old Alice was when she received her last allowance. (7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---




---

---

---

---



Question number	Scheme	Marks
6.	<p>(a)  (See below) Clearly through origin (or (0, 0) seen) 3 labelled (or (3, 0) seen)</p> <p>(b)  Stretch parallel to y-axis 1 and 4 labelled (or (1, 0) and (4, 0) seen) 6 labelled (or (0, 6) seen)</p> <p>(c)  Stretch parallel to x-axis 2 and 8 labelled (or (2, 0) and (8, 0) seen) 3 labelled (or (0, 3) seen)</p>	<p>M1 A1 A1 (3)</p> <p>M1 A1 A1 (3)</p> <p>M1 A1 A1 (3)</p> <p><b>Total 9 marks</b></p>

7.	<p>(a) <math>500 + (500 + 200) = 1200</math> or <math>S_2 = \frac{1}{2}2\{1000 + 200\} = 1200</math> (*)</p> <p>(b) Using <math>a = 500, d = 200</math> with <math>n = 7, 8</math> or <math>9</math> <math>a + (n - 1)d</math> or “listing” <math>500 + (7 \times 200) = (\pounds)1900</math></p> <p>(c) Using <math>\frac{1}{2}n\{2a + (n - 1)d\}</math> or <math>\frac{1}{2}n\{a + l\}</math>, or listing and “summing” terms <math>S_8 = \frac{1}{2}8\{2 \times 500 + 7 \times 200\}</math> or <math>S_8 = \frac{1}{2}8\{500 + 1900\}</math>, or all terms in list correct <math>= (\pounds) 9600</math></p> <p>(d) <math>\frac{1}{2}n\{2 \times 500 + (n - 1) \times 200\} = 32000</math> M1: General <math>S_n</math>, equated to 32000 <math>n^2 + 4n - 320 = 0</math> (or equiv.) M1: Simplify to 3 term quadratic <math>(n + 20)(n - 16) = 0</math> <math>n = \dots</math> M1: Attempt to solve 3 t.q. <math>n = 16,</math> Age is 26 A1cso,A1cso</p>	<p>B1 (1)</p> <p>M1 A1 (2)</p> <p>M1 A1 A1 (3)</p> <p>M1 A1 M1 A1 M1 A1cso,A1cso (7)</p> <p><b>Total 13 marks</b></p>
----	---	---







9.

Figure 2




Figure 2 shows part of the curve  $C$  with equation

$$y = (x - 1)(x^2 - 4).$$

The curve cuts the  $x$ -axis at the points  $P$ ,  $(1, 0)$  and  $Q$ , as shown in Figure 2.

(a) Write down the  $x$ -coordinate of  $P$ , and the  $x$ -coordinate of  $Q$ . (2)

(b) Show that  $\frac{dy}{dx} = 3x^2 - 2x - 4$ . (3)

(c) Show that  $y = x + 7$  is an equation of the tangent to  $C$  at the point  $(-1, 6)$ . (2)

The tangent to  $C$  at the point  $R$  is parallel to the tangent at the point  $(-1, 6)$ .

(d) Find the exact coordinates of  $R$ . (5)

---

---

---

---

---

---

---

---


---

---







Question number	Scheme	Marks
10.	(a) $x^2 + 2x + 3 = (x + 1)^2 + 2$ $(a = 1, b = 2)$	B1, B1      (2)
	(b) 	“U”-shaped parabola Vertex in correct quadrant (ft from $(-a, b)$ $(0, 3)$ (or 3 on y-axis)
	(c) $b^2 - 4ac = 4 - 12 = -8$ Negative, so curve does not cross $x$ -axis	B1 B1      (2)
	(d) $b^2 - 4ac = k^2 - 12$ (May be within the quadratic formula)	M1
	$k^2 - 12 < 0$ (Correct inequality expression in any form)	A1
	$-\sqrt{12} < k < \sqrt{12}$ (or $-2\sqrt{3} < k < 2\sqrt{3}$ )	M1 A1      (4)
<b>Total 11 marks</b>		