Winter 2009

www.mystudybro.com

Mathematics C1

This resource was created and owned by Pearson Edexcel Past Paper 6663 Surname Initial(s) Centre Paper Reference No. Signature Candidate 3 6 6 () 6 No. Paper Reference(s) 6663/01 Examiner's use only **Edexcel GCE** Team Leader's use only **Core Mathematics C1 Advanced Subsidiary** Question Leave Number Blank Friday 9 January 2009 – Morning 1 Time: 1 hour 30 minutes 2 3 4 Materials required for examination Items included with question papers 5 Mathematical Formulae (Green) Nil 6 Calculators may NOT be used in this examination. 7 8 9 10 **Instructions to Candidates** In the boxes above, write your centre number, candidate number, your surname, initials and signature. 11 Check that you have the correct question paper. Answer ALL the questions. You must write your answer for each question in the space following the question. **Information for Candidates** A booklet 'Mathematical Formulae and Statistical Tables' is provided. Full marks may be obtained for answers to ALL questions. The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 11 questions in this question paper. The total mark for this paper is 75. There are 28 pages in this question paper. Any blank pages are indicated. **Advice to Candidates** You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2009 Edexcel Limited.







Total

Turn over

Winter	2009
Past Pape	er

t Paper	This resource was created and owned by Pearson Edex	cel	666
			Leave
	$\frac{1}{2}$		
1. (a)	Write down the value of 125^3 .	(1)	
	2	(1)	
(b)	Find the value of $125^{-\frac{2}{3}}$.		
		(2)	
		(Total 3 marks)	
		. /	
			3
	\$\$\$ \$ \$\$\$ \$\$ \$\$ \$\$ 1		Turn ove

January 2009 6663 Core Mathematics C1 Mark Scheme

Que Num	stion nber	Scheme	Mark	٢S
1	(a) (b)	5 (±5 is B0)	B1	(1)
	(6)	$\frac{1}{(\text{their 5})^2}$ or $\left(\frac{1}{\text{their 5}}\right)^2$	M1	
		$=\frac{1}{25}$ or 0.04 $(\pm\frac{1}{25} \text{ is A0})$	A1	(2) [3]
	(b)	M1 follow through their value of 5. Must have reciprocal and square.		
		5^{-2} is <u>not</u> sufficient to score this mark, unless $\frac{1}{5^2}$ follows this.		
		A negative introduced at any stage can score the M1 but not the A1, e.g. $125^{-\frac{2}{3}} = \left(-\frac{1}{5}\right)^2 = \frac{1}{25}$ scores M1 A0		
		$125^{-\frac{2}{3}} = -\left(\frac{1}{5}\right)^2 = -\frac{1}{25}$ scores M1 A0.		
		Correct answer with no working scores both marks.		
		<u>Alternative</u> : $\frac{1}{\sqrt[3]{125^2}}$ or $\frac{1}{(125^2)^{\frac{1}{3}}}$ M1 (reciprocal and the correct number squared) $\left(=\frac{1}{\sqrt[3]{15625}}\right)$		
		$=\frac{1}{25}$ A1		

er 2009 aper	www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathema	tics C 666
			Leave blank
. Find	$\int (12x^5 - 8x^3 + 3) dx$, giving each term in its simplest form.		
	J ` ´	(4)	
			02
	(Ta	tal 1 marke)	
	(10	ται τ mai κ5j	

Question Number	Scheme	Marks
2	$(I =)\frac{12}{6}x^{6} - \frac{8}{4}x^{4} + 3x + c$ = 2x ⁶ - 2x ⁴ + 3x + c	M1 A1A1A1 [4]
	M1 for an attempt to integrate $x^n \to x^{n+1}$ (i.e. ax^6 or ax^4 or ax , where a is any non-zero constant). Also, this M mark can be scored for just the $+ c$ (seen at some stage), even if no other terms are correct. 1 st A1 for $2x^6$ 2 nd A1 for $-2x^4$ 3 rd A1 for $3x + c$ (or $3x + k$, etc., any appropriate letter can be used as the constant) Allow $3x^1 + c$, but not $\frac{3x^1}{1} + c$. Note that the A marks can be awarded at separate stages, e.g. $\frac{12}{6}x^6 - 2x^4 + 3x$ scores 2^{nd} A1 $\frac{12}{6}x^6 - 2x^4 + 3x + c$ scores 3^{rd} A1 $2x^6 - 2x^4 + 3x$ scores 1^{st} A1 (even though the c has now been lost). Remember that all the A marks are dependent on the M mark. If applicable, isw (ignore subsequent working) after a correct answer is seen.	
	Ignore wrong notation if the intention is clear, e.g. Answer $\int 2x^6 - 2x^4 + 3x + c dx$.	

2 Expand and simplify $(a/7 + 2)(a/7 - 2)$		Leave blank
5. Expand and simplify $(\sqrt{7} + 2)(\sqrt{7} - 2)$.	(2)	
		03
	(Total 2 marks)	
	(5

Question Number	Scheme	Marks
3	$\sqrt{7}^2 + 2\sqrt{7} - 2\sqrt{7} - 2^2$, or 7 - 4 or an exact equivalent such as $\sqrt{49} - 2^2 = 3$	M1 A1 [2]
	M1 for an expanded expression. At worst, there can be <u>one wrong term</u> and <u>one wrong sign</u> , or <u>two wrong signs</u> . e.g. $7+2\sqrt{7}-2\sqrt{7}-2$ is M1 (one wrong term -2) $7+2\sqrt{7}+2\sqrt{7}+4$ is M1 (two wrong signs $+2\sqrt{7}$ and $+4$) $7+2\sqrt{7}+2\sqrt{7}+2$ is M1 (one wrong term $+2$, one wrong sign $+2\sqrt{7}$) $\sqrt{7}+2\sqrt{7}-2\sqrt{7}+4$ is M1 (one wrong term $\sqrt{7}$, one wrong sign $+4$) $\sqrt{7}+2\sqrt{7}-2\sqrt{7}-2$ is M0 (two wrong terms $\sqrt{7}$ and -2) $7+\sqrt{14}-\sqrt{14}-4$ is M0 (two wrong terms $\sqrt{14}$ and $-\sqrt{14}$) If only 2 terms are given, they must be correct, i.e. $(7-4)$ or an equivalent unsimplified version to score M1. The terms can be seen <u>separately</u> for the M1. Correct answer with <u>no working</u> scores both marks.	

Ainter 2 ast Pape	2009 www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematic
		I
		t
4.	A curve has equation $y = I(x)$ and passes through the point (4, 22).	
	Given that	
	$f'(x) = 3x^2 - 3x^{\frac{1}{2}} - 7,$	
	use integration to find $f(x)$, giving each term in its simplest form.	
		(5)

N 3 0 0 8 1 A 0 6 2 8

Question Number	Scheme	Mark	S
4	$(f(x) =) \frac{3x^3}{3} - \frac{3x^2}{\frac{3}{2}} - 7x(+c)$ = $x^3 - 2x^{\frac{3}{2}} - 7x(+c)$ f(4) = 22 \Rightarrow 22 = 64 - 16 - 28 + c c = 2	M1 A1A1 M1 A1cso	(5)
			[5]
	1 st M1 for an attempt to integrate $(x^3 \text{ or } x^{\frac{3}{2}} \text{ seen})$. The <i>x</i> term is insufficient for this mark and similarly the + <i>c</i> is insufficient. 1 st A1 for $\frac{3}{3}x^3$ or $-\frac{3x^{\frac{3}{2}}}{\frac{3}{2}}$ (An unsimplified or simplified correct form) 2 nd A1 for all three <i>x</i> terms correct and simplified (the simplification may be seen later). The + <i>c</i> is not required for this mark. Allow $-7x^1$, but <u>not</u> $-\frac{7x^1}{1}$. 2 nd M1 for an attempt to use $x = 4$ <u>and</u> $y = 22$ in a changed function (even if differentiated) to form an equation in <i>c</i> . 3 rd A1 for <i>c</i> = 2 with no earlier incorrect work (a final expression for f(<i>x</i>) is not required).		

(3)

6663



Figure 1

Figure 1 shows a sketch of the curve C with equation y = f(x). There is a maximum at (0, 0), a minimum at (2, -1) and C passes through (3, 0).

On separate diagrams sketch the curve with equation

(a)
$$y = f(x+3)$$
,
(3)

(b)
$$y = f(-x)$$
.

On each diagram show clearly the coordinates of the maximum point, the minimum point and any points of intersection with the *x*-axis.



Question Number		Scheme	Marks
5	(a)	Shape \bigwedge , touching the x-axis maximum. Through (0,0) & -3 marked on x or (-3,0) seen. Allow (0,-3) if marked on the x Marked in the correct place, but 3 Min at (-1,-1)	at its M1 x-axis, A1 -axis. 3, is A0. A1 (3)
	(b)	Correct shape \bigvee (top left - bottom right) Through - 3 and max at (0, 0). Marked in the correct place, but 3 (-2,-1) Min at (-2,-1)	B1 B1 B1 B1 B1 (3) [6]
	(a)	M1 as described above. Be generous, even when the curve seems to be constraight line segments, but there must be a discernible 'curve' at the max 1 st A1 for curve passing through -3 and the origin. Max at (-3,0) 2 nd A1 for minimum at (-1,-1). Can simply be indicated on sketch.	nposed of and min.
	(b)	 1st B1 for the correct shape. A negative cubic passing from top left to bottom right. Shape: Be generous, even when the curve seems to be composed of straight line segments, but there must be a discernible 'curve' at the max. and min. 2nd B1 for curve passing through (-3,0) having a max at (0,0) and no other max. 3rd B1 for minimum at (-2,-1) and no other minimum. If in correct quadrant but labelled, e.g. (-2,1), this is B0. In each part the (0, 0) does <u>not</u> need to be written to score the second mark having the curve pass through the origin is sufficient. The last mark (for the minimum) in each part is dependent on a sketch being attempted, and the sketch must show the minimum in approximately the correct place (not, for example, (-2, -1) marked in the wrong quadrant). The mark for the minimum is <u>not</u> given for the coordinates just marked on the axes <u>unless</u> these are clearly linked to the minimum by vertical and horizontal lines. 	

Question Number		Scheme	Marks	
6	(a)	$2x^{\frac{3}{2}}$ or $p = \frac{3}{2}$ (<u>Not</u> $2x\sqrt{x}$)	B1	
	(b)	$ \begin{array}{ccc} -x & \text{or} & -x^{1} & \text{or} & q = 1 \\ \left(\frac{dy}{dx}\right) & 20x^{3} + 2 \times \frac{3}{2}x^{\frac{1}{2}} - 1 \end{array} $	B1 (M1	(2)
		$= \underline{20x^3 + 3x^{\frac{1}{2}} - 1}$	A1A1ftA1	ft (4) [6]
	(a)	$1^{st} B1 \text{for } p = 1.5 \text{ or exact equivalent} \\ 2^{nd} B1 \text{for } q = 1$		
	(b)	M1 for an attempt to differentiate $x^n \to x^{n-1}$ (for any of the 4 terms) 1 st A1 for 20 x^3 (the -3 must 'disappear')		
		2^{nd} A1ft for $3x^{\frac{1}{2}}$ or $3\sqrt{x}$. Follow through their <i>p</i> but they must be differentiating $2x^p$, where <i>p</i> is a <u>fraction</u> , and the coefficient must be simplified if necessary. 3^{rd} A1ft for -1 (<u>not</u> the unsimplified $-x^0$), or follow through for correct differentiation of their $-x^q$ (i.e. coefficient of x^q is -1). If ft is applied, the coefficient must be simplified if necessary.		
		'Simplified' coefficient means $\frac{a}{b}$ where a and b are integers with no common		
		factors. Only a single + or – sign is allowed (e.g. – – must be replaced by +).		
		If there is a 'restart' in part (b) it can be marked independently of part (a), but marks for part (a) cannot be scored for work seen in (b).		
		<u>Multiplying</u> by \sqrt{x} : (assuming this is a restart)		
		e.g. $y = 5x^4 \sqrt{x} - 3\sqrt{x} + 2x^2 - x^{7/2}$ $\left(\frac{dy}{dx}\right) = \frac{45}{2}x^{7/2} - \frac{3}{2}x^{-1/2} + 4x - \frac{3}{2}x^{1/2}$ scores M1 A0 A0 (<i>p</i> not a fraction) A1ft.		
		Extra term included: This invalidates the final mark.		
		e.g. $y = 5x^4 - 3 + 2x^2 - x^{/2} - x^{/2}$ $\begin{pmatrix} dy \\ 20x^3 + 4x - 3x^{1/2} - 1x^{-1/2} \\ y = 1x^{-1/2} \\ y =$		
		$\left(\frac{1}{dx}\right)^{20x} + 4x - \frac{1}{2}x^{2x} - \frac{1}{2}x^{2x}$ scores M1 A1 A0 (<i>p</i> not a fraction) A0. Numerator and denominator differentiated separately: For this, neither of the last two (ft) marks should be awarded.		
		<u>Quotient/product rule</u> : Last two terms must be correct to score the last 2 marks. (If the M mark has not already been earned, it can be given for the quotient/product rule attempt.)		

ist Pape	2009 r	www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics
7.	The equation f for x .	$kx^2 + 4x + (5 - k) = 0$, where k is a constant, has 2 differen	t real solutions
	(a) Show that	t <i>k</i> satisfies	
		$k^2 - 5k + 4 > 0.$	(3)
	(b) Hence fin	d the set of possible values of k .	(4)

Question Number	Scheme	Mark	s
7 (a)	$b^2 - 4ac > 0 \Rightarrow 16 - 4k(5-k) > 0$ or equiv., e.g. $16 > 4k(5-k)$		
	So $k^2 - 5k + 4 > 0$ (Allow any order of terms, e.g. $4 - 5k + k^2 > 0$) (*)	A1cso	(3)
(b)	<u>Critical Values</u> $(k-4)(k-1) = 0$ $k = \dots$ k = 1 or 4	M1 A1	
	Choosing "outside" region	M1	
	k < 1 or $k > 4$	A1	(4) [7]
	For this question, ignore (a) and (b) labels and award marks wherever correct work is se	een.	
(a)	M1 for attempting to use the discriminant of the initial equation (> 0 not required, but substitute of <i>a</i> , <i>b</i> and <i>c</i> in the correct formula is required). If the formula $b^2 - 4ac$ is seen, at least 2 of <i>a</i> , <i>b</i> and <i>c</i> must be correct. If the formula $b^2 - 4ac$ is not seen, all 3 (<i>a</i> , <i>b</i> and <i>c</i>) must be correct. This mark can still be scored if substitution in $b^2 - 4ac$ is within the quadratic formula. This mark can also be scored by comparing b^2 and $4ac$ (with substitution). However, use of $b^2 + 4ac$ is M0. 1 st A1 for fully correct expression, possibly unsimplified, with > symbol. NB must appear before the last line, even if this is simply in a statement such as $b^2 - 4ac > 0$ or 'discriminant positive Condone a bracketing slip, e.g. $16 - 4 \times k \times 5 - k$ if subsequent work is correct and convincing 2^{nd} A1 for a fully correct derivation with no incorrect working seen. Condone a bracketing slip if otherwise correct and convincing.		
(b)	 1st M1 for attempt to solve an appropriate 3TQ 1st A1 for both k = 1 and 4 (only the critical values are required, so accept, e.g. k > 1 and 2nd M1 for choosing the "outside" region. A diagram or table alone is not sufficient. Follow through their values of k. The set of values must be 'narrowed down' to score this M mark listing every k < 1, 1 < k < 4, k > 4 is M0. 2nd A1 for correct answer only, condone "k < 1, k > 4" and even "k < 1 and k > 4", but "1 > k > 4" is A0. ** Often the statement k > 1 and k > 4 is followed by the correct final answer. Allow fu Seeing 1 and 4 used as critical values gives the first M1 A1 by implication. In part (b), condone working with x's except for the final mark, where the set of values not of values of k (i.e. 3 marks out of 4). Use of ≤ (or ≥) in the final answer loses the final mark. 	nd <i>k</i> > 4) ything 11 marks. must be a	. ** 1 set

(1)

(5)

Leave blank

- 8. The point P (1, a) lies on the curve with equation $y = (x + 1)^2(2 x)$.
 - (a) Find the value of *a*.
 - (b) On the axes below sketch the curves with the following equations:

y

- (i) $y = (x + 1)^2(2 x)$,
- (ii) $y = \frac{2}{x}$.

On your diagram show clearly the coordinates of any points at which the curves meet the axes.

(c) With reference to your diagram in part (b) state the number of real solutions to the equation

$$(x+1)^2(2-x) = \frac{2}{x}.$$
(1)

х



Que: Num	stion Iber		Scheme	Mar	ks
8	(a)	$(a=) (1+1)^{2} (2-1) = \underline{4}$ (1, 4)	or $y = 4$ is also acceptable	B1	(1)
	(D)	Y	(i) Shape \bigvee or \bigwedge anywhere	B1	
		2	Min at $(-1,0)$ can be -1 on <i>x</i> -axis. Allow $(0,-1)$ if marked on the <i>x</i> -axis. Marked in the correct place, but 1, is B0.	B1	
		-1, 2	(2, 0) and (0, 2) can be 2 on axes	B1	
			(ii) Top branch in 1 st quadrant with 2 intersections	B1	
			Bottom branch in 3 rd quadrant (ignore any intersections)	B1	(5)
	(c)	(2 intersections therefore) $\underline{2}$ (roots)		B1ft	(1) [7]
	(b)	 1st B1 for shape or Can be anywhere, but there must be one max. and one min. and no further max. and min. turning points. Shape: Be generous, even when the curve seems to be composed of straight line segments, but there must be a discernible 'curve' at the max. and min. 2nd B1 for minimum at (-1,0) (even if there is an additional minimum point shown) 3rd B1 for the sketch meeting axes at (2, 0) and (0, 2). They can simply mark 2 on the axes. The marks for minimum and intersections are dependent upon having a sketch. Answers on the diagram for min. and intersections take precedence over answers seen elsewhere. 			
		other curve. The curve can 'touch' the axes. A curve of (roughly) the correct shape is required, but be very generous, even when the arc appears to turn 'inwards' rather than approaching the axes, and when the curve looks like two straight lines with a small curve at the join. Allow, for example, shapes like these:			
		5 th B1 for a branch fully in the 3 rd quadrant (ignore any intersections with the other curve for this branch). The curve can 'touch' the axes. A curve of (roughly) the correct shape is required, but be very generous, even when the arc appears to turn 'inwards' rather than approaching the axes.			is arc
	(c)	B1ft for a statement about the number of roots - compatible with their sketch. No sketch is B0. The answer 2 <u>incompatible with the sketch</u> is B0 (ignore any algebra seen). If the sketch shows the 2 correct intersections <u>and</u> , for example, one other intersection, the answer here should be 3, not 2, to score the mark.). ne

Winter : Past Pape	2009	www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics C
			Leav
9.	The first term	m of an arithmetic series is a and the common difference is d .	
	The 18th term	m of the series is 25 and the 21st term of the series is $32\frac{1}{2}$.	
	(a) Use this	information to write down two equations for a and d .	(2)
	(b) Show th	hat $a = -17.5$ and find the value of <i>d</i> .	(2)
	The sum of t	the first n terms of the series is 2750.	
	(c) Show th	hat <i>n</i> is given by	
		$n^2 - 15n = 55 \times 40.$	
			(4)
	(d) Hence fi	ind the value of <i>n</i> .	(3)

Question Number		Scheme	Mar	ks	
9	(a) (b)	a + 17d = 25 or equiv. (for 1 st B1), $a + 20d = 32.5$ or equiv. (for 2 nd B1), <u>Solving</u> (Subtract) $3d = 7.5$ so $d = 2.5$ $a = 32.5 - 20 \times 2.5$ so $a = -17.5$ (*)	B1, B1 M1 A1cso	(2) (2)	
	(c)	$2750 = \frac{n}{2} \left[-35 + \frac{5}{2} (n-1) \right]$ { $4 \times 2750 = n(5n-75)$ } $4 \times 550 = n(n-15)$ $n^2 - 15n = 55 \times 40$ (*)	M1A1ft M1 A1cso	(4)	
	(d)	$n^{2} - 15n - 55 \times 40 = 0 \text{or} n^{2} - 15n - 2200 = 0$ $(n - 55)(n + 40) = 0 \qquad n = \dots$ $\underline{n = 55} \text{(ignore - 40)}$	M1 M1 A1	(3) [11]	
	(a)	Mark parts (a) and (b) as 'one part', ignoring labelling. <u>Alternative</u> : $1^{\text{st}} \text{B1:} d = 2.5 \text{ or equiv. or } d = \frac{32.5 - 25}{2}$. No method required, but $a = -17.5$ must not be assumed.			
	(b)	 2nd B1: Either a +17d = 25 or a + 20d = 32.5 seen, or used with a value of d or for 'listing terms' or similar methods, 'counting back' 17 (or 20) terms. M1: In main scheme: for a full method (allow numerical or sign slips) leading to solution for d or a without assuming a = -17.5 In alternative scheme: for using a d value to find a value for a 			
		A1: Finding correct values for both <i>a</i> and <i>d</i> (allowing equiv. fractions such as $d = \frac{15}{6}$),	with no		
	(c)	In the main scheme, if the given <i>a</i> is used to find <i>d</i> from one of the equations, then allow M1A1 if both values are <u>checked</u> in the 2^{nd} equation.			
	(d)	1 st M1 for attempt to form equation with correct S_n formula and 2750, with values of <i>a</i> and <i>d</i> . 1 st A1ft for a correct equation following through their <i>d</i> . 2 nd M1 for expanding and simplifying to a 3 term quadratic. 2 nd A1 for correct working leading to printed result (no incorrect working seen).			
		1 st M1 forming the correct $3TQ = 0$. Can condone missing "= 0" but all terms must be on one side. First M1 can be implied (perhaps seen in (c), but there must be an attempt at (d) for it to be scored). 2 nd M1 for attempt to solve 3TQ, by factorisation, formula or completing the square (see general marking principles at end of scheme). If this mark is earned for the 'completing the square' method or if the factors are written down directly, the 1 st M1 is given by implication. A1 for <i>n</i> = 55 dependent on both Ms. Ignore – 40 if seen. No working or 'trial and improvement' methods in (d) score all 3 marks for the answer 55, otherwise no marks.			

www.mystudybro.com This resource was created and owned by Pearson Edexcel 6663 Leave blank 10. The line l_1 passes through the point A (2, 5) and has gradient $-\frac{1}{2}$. (a) Find an equation of l_1 , giving your answer in the form y = mx + c. (3) The point *B* has coordinates (-2, 7). (b) Show that *B* lies on l_1 . (1) (c) Find the length of AB, giving your answer in the form $k\sqrt{5}$, where k is an integer. (3) The point C lies on l_1 and has x-coordinate equal to p. The length of AC is 5 units. (d) Show that *p* satisfies $p^2 - 4p - 16 = 0.$ (4)



Ques Num	stion ber	Scheme	Marks
10	(a)	$y-5 = -\frac{1}{2}(x-2)$ or equivalent, e.g. $\frac{y-5}{x-2} = -\frac{1}{2}$, $y = -\frac{1}{2}x+6$	M1A1, A1cao (3)
	(b)	$x = -2 \Rightarrow y = -\frac{1}{2}(-2) + 6 = 7$ (therefore <i>B</i> lies on the line)	B1 (1)
		(or equivalent verification methods)	
	(c)	$(AB^{2} =) (2 - 2)^{2} + (7 - 5)^{2}, = 16 + 4 = 20, AB = \sqrt{20} = 2\sqrt{5}$	(3) MI, AI, AI
		<i>C</i> is $(p, -\frac{1}{2}p+6)$, so $AC^2 = (p-2)^2 + \left(-\frac{1}{2}p+6-5\right)^2$	M1
	(d)	Therefore $25 = p^2 - 4p + 4 + \frac{1}{4}p^2 - p + 1$	M1
		$25 = 1.25p^2 - 5p + 5$ or $100 = 5p^2 - 20p + 20$ (or better, RHS simplified to 3 terms)	A1
		Leading to: $0 = p^2 - 4p - 16$ (*)	[11]
	(a)	 M1 A1 The version in the scheme above can be written down directly (for 2 marks), and M1 A0 can be allowed if there is just one slip (sign or number). If the 5 and 2 are the wrong way round the M mark can still be given if a correct formula (e.g. y - y₁ = m(x - x₁)) is seen, otherwise M0. If (2, 5) is substituted into y = mx + c to find c, the M mark is for attempting this and the 1st A mark is for c = 6. Correct answer without working or from a sketch scores full marks. 	
	(b)	A conclusion/comment is not required, except when the method used is to establish that the line through $(-2,7)$ with gradient $-\frac{1}{2}$ has the same eqn. as found in part (a),	
		of to establish that the line through $(-2, 7)$ and $(2, 3)$ has gradient $-\frac{1}{2}$. In these cases	
	(c)	M1 for attempting AB^2 or AB . Allow one slip (sign or number) <u>inside</u> a bracket, i.e. do not allow $(22)^2 - (7-5)^2$.	
		1 st A1 for 20 (condone bracketing slips such as $-2^2 = 4$)	
		2^{nd} A1 for $2\sqrt{5}$ or $k = 2$ (Ignore ± here).	
	(d)	1 st M1 for $(p-2)^2$ + (linear function of p) ² . The linear function may be unsimplified	
		but must be equivalent to $ap + b$, $a \neq 0$, $b \neq 0$. 2^{nd} M1 (dependent on 1^{st} M) for forming an equation in <i>p</i> (using 25 or 5) and attempting (perhaps not very well) to multiply out both brackets. 1^{st} A1 for collecting like <i>p</i> terms and having a correct expression. 2^{nd} A1 for correct work leading to printed answer. Alternative, using the result:	
		Solve the quadratic $(p = 2 \pm 2\sqrt{5})$ and use one or both of the two solutions to find the	
		length of AC^2 or $C_1C_2^2$: e.g. $AC^2 = (2 + 2\sqrt{5} - 2)^2 + (5 - \sqrt{5} - 5)^2$ scores 1 st M1, and 1 st A1 if fully correct.	
		Finding the length of AC or AC^2 for both values of p , or finding C_1C_2 with some evidence of halving (or intending to halve) scores the 2 nd M1.	
		Getting $AC = 5$ for both values of p, or showing $\frac{1}{2}C_1C_2 = 5$ scores the 2 nd A1 (cso).	

11. The curve <i>C</i> has equation $y = 9 - 4x - \frac{8}{x}$, $x > 0$. The point <i>P</i> on <i>C</i> has <i>x</i> -coordinate equal to 2. (a) Show that the equation of the tangent to <i>C</i> at the point <i>P</i> is $y = 1 - 2x$. (b) Find an equation of the normal to <i>C</i> at the point <i>P</i> . (c) Find the area of the tangent <i>A PB</i> .	6663		st Paper This res
11. The curve <i>C</i> has equation $y = 9 - 4x - \frac{8}{x}, x > 0.$ The point <i>P</i> on <i>C</i> has <i>x</i> -coordinate equal to 2. (a) Show that the equation of the tangent to <i>C</i> at the point <i>P</i> is $y = 1 - 2x$. (b) Find an equation of the normal to <i>C</i> at the point <i>P</i> . (c) Find the area of triangle <i>APB</i> . (d) (e) Find the area of triangle <i>APB</i> . (f)	Leave blank		
$y = 9 - 4x - \frac{8}{x}, x > 0.$ The point <i>P</i> on <i>C</i> has <i>x</i> -coordinate equal to 2. (a) Show that the equation of the tangent to <i>C</i> at the point <i>P</i> is $y = 1 - 2x$. (b) Find an equation of the normal to <i>C</i> at the point <i>P</i> . (c) Find the area of triangle <i>APB</i> . (f)			11. The curve <i>C</i> has equation
The point <i>P</i> on <i>C</i> has <i>x</i> -coordinate equal to 2. (a) Show that the equation of the tangent to <i>C</i> at the point <i>P</i> is $y = 1 - 2x$. (b) Find an equation of the normal to <i>C</i> at the point <i>P</i> . (c) Find the area of triangle <i>APB</i> . (4)			У
(a) Show that the equation of the tangent to <i>C</i> at the point <i>P</i> is <i>y</i> = 1 - 2 <i>x</i> . (6) (b) Find an equation of the normal to <i>C</i> at the point <i>P</i> . (3) The tangent at <i>P</i> meets the <i>x</i> -axis at <i>A</i> and the normal at <i>P</i> meets the <i>x</i> -axis at <i>B</i> . (c) Find the area of triangle <i>APB</i> . (c) Find the area of triangle <i>APB</i> . (4) (a) (b) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) (c) Find the area of triangle <i>APB</i> . (c) <			The point <i>P</i> on <i>C</i> has <i>x</i> -co
(6) (b) Find an equation of the normal to <i>C</i> at the point <i>P</i> . (c) (c) Find the area of triangle <i>APB</i> . (d)			(a) Show that the equation
(b) Find an equation of the normal to C at the point P. (3) The tangent at P meets the x-axis at A and the normal at P meets the x-axis at B. (c) Find the area of triangle APB. (d) (d)		(6)	
The tangent at P meets the x-axis at A and the normal at P meets the x-axis at B. (4) (c) Find the area of triangle APB. (4)		(3)	(b) Find an equation of the
(c) Find the area of triangle APB. (4)		2	The tangent at P meets th
		•	(a) Find the area of trian
		(4)	(c) Find the area of trian
24			
			24

Question Number	Scheme	Marks	
11 (a)	$\left(\frac{dy}{dt}\right) = -4 + 8x^{-2}$ (4 or $8x^{-2}$ for M1 sign can be wrong)	M1A1	
	$ \begin{array}{c} (dx) \\ x = 2 \Longrightarrow m = -4 + 2 = -2 \end{array} $	M1	
	$y = 9 - 8 - \frac{8}{2} = -3$ The first 4 marks <u>could</u> be earned in part (b)	B1	
	Equation of tangent is: $y+3 = -2(x-2) \rightarrow y = 1-2x$ (*)	M1 A1cso (6)	
(b)	Gradient of normal = $\frac{1}{2}$	B1ft	
	Equation is: $\frac{y+3}{x-2} = \frac{1}{2}$ or better equivalent, e.g. $y = \frac{1}{2}x - 4$	M1A1	
(c)	$(A:) \frac{1}{2}, \qquad (B:) 8$	(3) B1, B1	
	Area of triangle is: $\frac{1}{2}(x_B \pm x_A) \times y_P$ with values for all of x_B, x_A and y_P	M1	
	$\frac{1}{2}\left(8 - \frac{1}{2}\right) \times 3 = -\frac{45}{4}$ or 11.25	A1 (4) [13]	
(a)	1 st M1 for 4 or $8x^{-2}$ (ignore the signs).		
	2 nd M1 for substituting $r = 2$ into their $\frac{dy}{dx}$ (must be different from their y)		
	dx B1 for $y = -3$ but not if clearly found from the given equation of the tangent		
	3^{rd} M1 for attempt to find the equation of tangent at P, follow through their m and y_P .		
	Apply general principles for straight line equations (see end of scheme). <u>NO DIFFERENTIATION ATTEMPTED</u> : Just assuming $m = -2$ at this stage is 2 nd A1cso for correct work leading to printed answer (allow equivalents with 2x, y, and such as $2x + y - 1 = 0$)	s M0 I 1 terms…	
(b)	B1ft for correct use of the perpendicular gradient rule. Follow through their <i>m</i> , but is there must be clear evidence that the <i>m</i> is thought to be the gradient of the tangent of tangent of the tangent of tangen	f $m \neq -2$ ent.	
	M1 for an attempt to find normal at P using their changed gradient and their y_P .		
	Appry general principles for straight line equations (see end of scheme). A1 for any correct form as specified above (correct answer only).		
(C)	$1^{\text{st}} \text{B1 for } \frac{1}{2} \text{ and } 2^{\text{nd}} \text{B1 for 8.}$		
	M1 for a full method for the area of triangle <i>ABP</i> . Follow through their x_A, x_B and	their y_P , but	
	the mark is to be awarded 'generously', condoning sign errors The final answer must be positive for A1, with negatives in the working condor	ned.	
	Determinant: Area = $\frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = \frac{1}{2} \begin{vmatrix} 2 & -3 & 1 \\ 0.5 & 0 & 1 \\ 8 & 0 & 1 \end{vmatrix} = \dots$ (Attempt to multiply out required for N		
	<u>Alternative</u> : $AP = \sqrt{(2-0.5)^2 + (-3)^2}$, $BP = \sqrt{(2-8)^2 + (-3)^2}$, Area $= \frac{1}{2}AP \times BP =$ M1		
	Intersections with y-axis instead of x-axis: Only the M mark is available B0 B0 M1 A0.		