

Write your name here

Surname	Other names
---------	-------------

**Pearson
Edexcel GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Core Mathematics C2

Advanced Subsidiary

Wednesday 24 May 2017 – Morning
Time: 1 hour 30 minutes

Paper Reference
6664/01

You must have:
Mathematical Formulae and Statistical Tables (Pink)

Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P44824A

©2017 Pearson Education Ltd.

1/1/2/1/1/1/



Pearson

Leave blank

- 1. Find the first 4 terms, in ascending powers of x , of the binomial expansion of

$$\left(3 - \frac{1}{3}x\right)^5$$

giving each term in its simplest form.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme	Marks
1.	$\left(3 - \frac{1}{3}x\right)^5 -$ $3^5 + {}^5C_1 3^4 \left(-\frac{1}{3}x\right) + {}^5C_2 3^3 \left(-\frac{1}{3}x\right)^2 + {}^5C_3 3^2 \left(-\frac{1}{3}x\right)^3 \dots$ First term of 243 $\left({}^5C_1 \times \dots \times x\right) + \left({}^5C_2 \times \dots \times x^2\right) + \left({}^5C_3 \times \dots \times x^3\right) \dots$ $= (243 \dots) - \frac{405}{3}x + \frac{270}{9}x^2 - \frac{90}{27}x^3 \dots$ $= (243 \dots) - 135x + 30x^2 - \frac{10}{3}x^3 \dots$	B1 M1 A1 A1 (4) [4]
Alternative method	$\left(3 - \frac{1}{3}x\right)^5 = 3^5 \left(1 - \frac{x}{9}\right)^5$ $3^5 \left(1 + {}^5C_1 \left(-\frac{1}{9}x\right) + {}^5C_2 \left(-\frac{1}{9}x\right)^2 + {}^5C_3 \left(-\frac{1}{9}x\right)^3 \dots\right)$ Scheme is applied exactly as before	
Notes B1: The constant term should be 243 in their expansion M1: Two of the three binomial coefficients must be correct and must be with the correct power of x. Accept 5C_1 or $\binom{5}{1}$ or 5 as a coefficient, and 5C_2 or $\binom{5}{2}$ or 10 as another and 5C_3 or $\binom{5}{3}$ or 10 as another..... Pascal's triangle may be used to establish coefficients. NB: If they only include the first two of these terms then the M1 may be awarded. A1: Two of the final three terms correct – may be unsimplified i.e. two of $-135x + 30x^2 - \frac{10}{3}x^3$ correct, or two of $-\frac{405}{3}x + \frac{270}{9}x^2 - \frac{90}{27}x^3$ (may be just two terms) A1: All three final terms correct and simplified. (Can be listed with commas or appear on separate lines. Accept in reverse order.) Accept correct alternatives to $-\frac{10}{3}$ e.g. $-3\frac{1}{3}$ or $-3.\bar{3}$ the recurring must be clear. 3.3 is not acceptable. Allow e.g. $+ -135x$		
e.g. The common error $3^5 + {}^5C_1 3^4 \left(-\frac{1}{3}\right)x + {}^5C_2 3^3 \left(-\frac{1}{3}\right)x^2 + {}^5C_3 3^2 \left(-\frac{1}{3}\right)x^3 = (243) - 135x - 90x^2 - 30x^3$ would earn B1, M1, A0, A0, so 2/4 If extra terms are given then isw No negative signs in answer also earns B1, M1, A0, A0 If the series is divided through by 3 at the final stage after an error or omission resulting in all multiple of three coefficients then apply scheme to series before this division and ignore subsequent work (isw) Special Case: Only gives first three terms $= (243 \dots) - 135x + 30x^2 \dots$ or $243 - \frac{405}{3}x + \frac{270}{9}x^2$ Follow the scheme to give B1 M1 A1 A0 special case. (Do not treat as misread.) Answers such as $243 + 405 - \frac{1}{3}x + 270 - \frac{1}{9}x^2 + 90 - \frac{1}{27}x^3 \dots$ gain no credit as the binomial coefficients are not linked to the x terms.		

Leave blank

2. In the triangle ABC , $AB = 16$ cm, $AC = 13$ cm, angle $ABC = 50^\circ$ and angle $BCA = x^\circ$

Find the two possible values for x , giving your answers to one decimal place.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme	Marks
2.	$\frac{\sin x}{16} = \frac{\sin 50^\circ}{13}$ $(\sin x) = \frac{16 \times \sin 50}{13} (= 0.943 \text{ but accept } 0.94)$ $x = \text{awrt } 70.5(3) \text{ and } 109.5 \quad \text{or } 70.6 \text{ and } 109.4$	M1 A1 dM1 A1 (4) [4]
<p style="text-align: center;">Notes</p> <p>M1: use sine formula correctly in any form. Allow awrt 0.77 for $\sin 50^\circ$ A1: give the correct value or correct expression for $\sin x$ (this implies the M1 mark). If it is given as expression they do not need degrees symbol. $\frac{\sin 50 \times 16}{13}$ is fine, If this is given as a decimal allow answers which round to 0.94. Allow awrt -0.323 (radians) here but no further marks are available. If they give this as x (not $\sin x$) and do not recover this is A0 dM1: Correct work leading to $x = \dots$ (via inverse sin) expression or value for $\sin x$ If the previous A mark has been awarded for a correct expression then this is for getting to awrt 70.5 or 109.5 (allow for 70.6 or 109.4). If the previous A mark was not gained, e.g. rounding errors were made in rearranging the correct sine formula then award dM1 for evidence of use of inverse sin in degrees on their value for $\sin x$ (may need to check on calculator). NB 70.5 following a correct sine formula will gain M1A1M1. A1: deduce and state both of the answers $x = 70.5$ and 109.5 (do not need degrees) Accept awrt these. Also accept 70.6 and 109.4. (Second answer is sometimes obtained by a long indirect route but still scores A1) If working in radians throughout, answers are 1.23 and 1.91 and this can be awarded M1 A1 M1 A0 (Working with 50 radians gives probable answers of -0.3288 and 3.47 giving M1A1M0A0) Special case: Wrong labelling of triangle. This simplifies the problem as there is only one solution for angle x. So it is not treated as a misread. If they find the missing side as awrt 12.6 then proceed to find an angle or its sine or cosine then give M1A0M0A0</p>		
<p>Alternative Method using cosine rule Let $BC = a$. M1: uses the cosine rule to form a three term quadratic equation in a (e.g. $a^2 - 32a \cos 50^\circ + 87 = 0$ or $a^2 - \text{awrt } 20.6a + 87 = 0$ though allow slips in signs rearranging) A1: Solves and obtains a correct value for a of awrt 14.6 or awrt 5.95. dM1: A correct full method to find (at least) one of the two angles. May use cosine rule again, or find angle BAC and then use sine rule. As in the main scheme, if the previous A mark has been awarded then they should obtain one of the correct angles for this mark. A1: deduces both correct answer as in main scheme. NB obtaining only one correct angle will usually score M1A1M1A0 in any method.</p>		

Leave blank

3. (a) $y = 5^x + \log_2(x + 1), \quad 0 \leq x \leq 2$

Complete the table below, by giving the value of y when $x = 1$

x	0	0.5	1	1.5	2
y	1	2.821		12.502	26.585

(1)

(b) Use the trapezium rule, with all the values of y from the completed table, to find an approximate value for

$$\int_0^2 (5^x + \log_2(x + 1)) dx$$

giving your answer to 2 decimal places.

(4)

(c) Use your answer to part (b) to find an approximate value for

$$\int_0^2 (5 + 5^x + \log_2(x + 1)) dx$$

giving your answer to 2 decimal places.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme						Marks
<p>3.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	x	0	0.5	1	1.5	2	<p>B1 cao</p> <p style="text-align: right;">(1)</p> <p>B1 oe</p> <p>M1A1ft</p> <p>A1</p> <p style="text-align: right;">(4)</p> <p>B1ft</p> <p style="text-align: right;">(1)</p> <p style="text-align: right;">[6]</p>
	y	1	2.821	6	12.502	26.585	
	<p>{At $x = 1,$ } $y = 6$ (allow 6.000 or even 6.00)</p>						
	<p>$\frac{1}{2} \times 0.5 ;$</p> <p style="text-align: center;">$\{ 1 + 26.585 + 2(2.821 + \text{their } 6 + 12.502) \}$ <u>For structure of {.....} ;</u></p> <p>$\frac{1}{2} \times 0.5 \{ 1 + 26.585 + 2(2.821 + 6 + 12.502) \} \{ = \frac{1}{4}(70.231) = 17.557.. \} = \text{awrt } 17.56$</p>						
Notes							
<p>(a)</p> <p>(b)</p>	<p>B1: 6</p> <p>B1: for using $\frac{1}{2} \times 0.5$ or $\frac{1}{4}$ or equivalent.</p> <p>M1: requires the correct {.....} bracket structure. It needs the first bracket to contain first y value plus last y value and the second bracket to be multiplied by 2 and to be the summation of the remaining y values in the table with no additional values. If the only mistake is a copying error or is to omit one value from 2nd bracket this may be regarded as a slip and the M mark can be allowed (An extra repeated term forfeits the M mark however). M0 if values used in brackets are x values instead of y values</p> <p>A1ft: for the correct bracket {.....} following through candidate's y value found in part (a).</p> <p>A1: for answer which rounds to 17.56</p> <p>NB: Separate trapezia may be used: B1 for 0.25, M1 for $\frac{1}{2} h(a + b)$ used 3 or 4 times (and A1ft if it is all correct) Then A1 as before.</p> <p>Special case: Bracketing mistake $0.25 \times (1 + 26.585) + 2(2.821 + \text{their } 6 + 12.502)$ scores B1 M1 A0 A0 unless the final answer implies that the calculation has been done correctly (then full marks can be given). An answer of 49.542 usually indicates this error.</p>						
<p>(c)</p>	<p>B1ft: 10 + their answer to part (b)</p> <p>(May be obtained by using the trapezium rule again with all values for y increased by 5)</p>						

Leave blank

4.




Diagram not drawn to scale

Figure 1

Figure 1 is a sketch representing the cross-section of a large tent $ABCDEF$.
 AB and DE are line segments of equal length.
 Angle FAB and angle DEF are equal.
 F is the midpoint of the straight line AE and FC is perpendicular to AE .
 BCD is an arc of a circle of radius 3.5 m with centre at F .
 It is given that

$$AF = FE = 3.7\text{m}$$

$$BF = FD = 3.5\text{m}$$

$$\text{angle } BFD = 1.77 \text{ radians}$$

Find

- (a) the length of the arc BCD in metres to 2 decimal places, (2)
- (b) the area of the sector $FBCD$ in m^2 to 2 decimal places, (2)
- (c) the total area of the cross-section of the tent in m^2 to 2 decimal places. (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




Question Number	Scheme	Marks
4. (a)	Usually answered in radians: Uses $BCD = 3.5 \times (\text{angle})$, $= 3.5 \times 1.77 = 6.195$ (m) (accept awrt 6.20)	M1 A1 (2)
(b)	Area = $\frac{1}{2}(3.5)^2 \times 1.77 = 10.84$ (m ²)	M1 A1 (2)
(c)	Area of triangle = $\frac{1}{2} \times 3.7 \times 3.5 \times \sin(\text{angle})$, $= \frac{1}{2} \times 3.7 \times 3.5 \times \sin(\frac{\pi}{2} - \frac{1.77}{2})$ (=awrt 4.1) Total area = "10.84" + 2 × "4.101" = 19.04	M1, A1 M1 A1cao (4) [8]
Notes		
(a)	M1: uses $s = 3.5 \times \theta$ with θ in radians or completely correct work in degrees A1: awrt 6.20 or just 6.2 (do not need to see units) Correct answer can imply the method.	
(b)	M1 for attempt to use $A = \frac{1}{2} \times 3.5^2 \times \theta$ (Accept θ in degrees.) A1 for awrt 10.84 (do not need to see units) isw if correct answer is followed by 10.8. Correct answer can imply the method.	
(c)	M1: Uses area of triangle $\frac{1}{2} \times 3.7 \times 3.5 \times \sin(\text{angle})$ Must be correct method for area of triangle but may be less direct. A1: Correct expression using correct angle i.e. $\frac{\pi}{2} - \frac{1.77}{2}$ or awrt 0.69 or awrt 39 degrees (need at least 2 sf if no other work seen, but may be implied by correct final answer) If correct expression is given then isw (so e.g. isw an answer of 0.0775 implying angle set to degrees on calculator) M1: Adds twice their second calculated area (even if rectangle or segment) to their sector area (may have been slips or errors in one or both formulae – such as missing $\frac{1}{2}$ or mixture of degrees and radians or weak attempt at triangle area) so M0A0M1A0 is a possible mark distribution A1: 19.04 cao (common answer through insufficient accuracy is 19.08 which loses final mark) Special Case. The mark profile M1A0M1A0M1A0M1A0 can be given if the angle is misunderstood as 1.77π or as AFB for example If "10.84" + $3.5 \times 3.7 \sin(\text{angle})$ is used then this can gain both M marks and the A marks if correct. But use of $3.5 \times 3.7 \sin(\text{angle})$ and later doubled and added to "10.84" is 1 st M0, 2 nd M1.	

Question number	Scheme	Marks
5	$x^2 + y^2 - 10x + 6y + 30 = 0$	
(a)	Uses any appropriate method to find the coordinates of the centre, e.g achieves $\underline{(x \pm 5)^2} + \underline{(y \pm 3)^2} = \dots$. Accept $(\pm 5, \pm 3)$ as indication of this.	M1

Question Number	Scheme	Marks
4. (a)	Usually answered in radians: Uses $BCD = 3.5 \times (\text{angle})$, $= 3.5 \times 1.77 = 6.195$ (m) (accept awrt 6.20)	M1 A1 (2)
(b)	Area = $\frac{1}{2}(3.5)^2 \times 1.77 = 10.84$ (m ²)	M1 A1 (2)
(c)	Area of triangle = $\frac{1}{2} \times 3.7 \times 3.5 \times \sin(\text{angle})$, $= \frac{1}{2} \times 3.7 \times 3.5 \times \sin(\frac{\pi}{2} - \frac{1.77}{2})$ (=awrt 4.1) Total area = "10.84" + 2 × "4.101" = 19.04	M1, A1 M1 A1cao (4) [8]
Notes		
(a)	M1: uses $s = 3.5 \times \theta$ with θ in radians or completely correct work in degrees A1: awrt 6.20 or just 6.2 (do not need to see units) Correct answer can imply the method.	
(b)	M1 for attempt to use $A = \frac{1}{2} \times 3.5^2 \times \theta$ (Accept θ in degrees.) A1 for awrt 10.84 (do not need to see units) isw if correct answer is followed by 10.8. Correct answer can imply the method.	
(c)	M1: Uses area of triangle $\frac{1}{2} \times 3.7 \times 3.5 \times \sin(\text{angle})$ Must be correct method for area of triangle but may be less direct. A1: Correct expression using correct angle i.e. $\frac{\pi}{2} - \frac{1.77}{2}$ or awrt 0.69 or awrt 39 degrees (need at least 2 sf if no other work seen, but may be implied by correct final answer) If correct expression is given then isw (so e.g. isw an answer of 0.0775 implying angle set to degrees on calculator) M1: Adds twice their second calculated area (even if rectangle or segment) to their sector area (may have been slips or errors in one or both formulae – such as missing $\frac{1}{2}$ or mixture of degrees and radians or weak attempt at triangle area) so M0A0M1A0 is a possible mark distribution A1: 19.04 cao (common answer through insufficient accuracy is 19.08 which loses final mark) Special Case. The mark profile M1A0M1A0M1A0M1A0 can be given if the angle is misunderstood as 1.77π or as AFB for example If "10.84" + $3.5 \times 3.7 \sin(\text{angle})$ is used then this can gain both M marks and the A marks if correct. But use of $3.5 \times 3.7 \sin(\text{angle})$ and later doubled and added to "10.84" is 1 st M0, 2 nd M1.	

Question number	Scheme	Marks
5	$x^2 + y^2 - 10x + 6y + 30 = 0$	
(a)	Uses any appropriate method to find the coordinates of the centre, e.g achieves $\underline{(x \pm 5)^2} + \underline{(y \pm 3)^2} = \dots$. Accept $(\pm 5, \pm 3)$ as indication of this.	M1

	Centre is $(5, -3)$.	A1	(2)
(b) Way 1	Uses $(x \pm 5)^2 - 5^2 + (y \pm 3)^2 - 3^2 + 30 = 0$ to give $r = \sqrt{25 + 9 - 30}$ or $r^2 = 25 + 9 - 30$ (not $30 - 25 - 9$) $r = 2$	M1 A1cao	(2)
Or Way 2	Using $\sqrt{g^2 + f^2 - c}$ from $x^2 + y^2 + 2gx + 2fy + c = 0$ (Needs formula stated or correct working) $r = 2$	M1 A1	(2)
(c) Way 1	Use $x = 4$ in an equation of circle and obtain equation in y only e.g. $(4 - 5)^2 + (y + 3)^2 = 4$ or $4^2 + y^2 - 10 \times 4 + 6y + 30 = 0$ Solve their quadratic in y and obtain two solutions for y e.g. $(y + 3)^2 = 3$ or $y^2 + 6y + 6 = 0$ so $y = -3 \pm \sqrt{3}$	M1 dM1 A1	(3)
Or Way 2	 Divide triangle PTQ and use Pythagoras with $r^2 - (5 - 4)^2 = h^2$, Find h and evaluate $-3 \pm h$. May recognise $(1, \sqrt{3}, 2)$ triangle. So $y = -3 \pm \sqrt{3}$	M1 dM1 A1	(3)
			[7]

Notes	
(a)	<p>Parts (a) and (b) can be marked together</p> <p>M1 as in scheme and can be <u>implied</u> by $(\pm 5, \pm 3)$ May be awarded for writing LHS as $(x \pm 5)^2 + (y \pm 3)^2 = \dots$</p> <p>or by comparing with $x^2 + y^2 + 2gx + 2fy + c = 0$ to write down centre $(-g, -f)$ directly</p> <p>A1: $(5, -3)$. This correct answer implies M1A1</p>
(b)	<p>M1 for a full correct method leading to $r = \dots$, or $r^2 =$ with their 5, their -3, their 25 and their 9 and their “-30”. Completion of square method errors result in M0 here. Usually $r = 4$ or $r = 16$ imply M0A0</p> <p>A1 2 cao Do not accept $r = \pm 2$ unless it is followed by $(r =) 2$ The correct answer with no wrong work seen implies M1A1</p> <p>Special case: if centre is given as $(-5, -3)$ or $(5, 3)$ or $(-5, 3)$ allow M1A1 for $r = 2$ worked correctly. i.e. $r^2 = "25" + "9" - 30$</p>
(c)	<p>M1: <i>Way 1:</i> Use $x = 4$ in a circle equation (may have wrong centre and/or radius) to obtain an equation in y only</p> <p>or <i>Way 2.</i> Uses geometry to find equation in h (ft on their radius and centre)</p> <p>dM1: (needs first method mark) Solve their quadratic in y or <i>Way 2.</i> Uses their h and their y coordinate correctly</p> <p>A1: cao</p>

Leave blank

6. $f(x) = -6x^3 - 7x^2 + 40x + 21$

(a) Use the factor theorem to show that $(x + 3)$ is a factor of $f(x)$ (2)

(b) Factorise $f(x)$ completely. (4)

(c) Hence solve the equation

$$6(2^{3y}) + 7(2^{2y}) = 40(2^y) + 21$$

giving your answer to 2 decimal places. (3)

Handwriting practice lines consisting of horizontal lines for student answers.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme	Marks
6. (a)	Attempt $f(3)$ or $f(-3)$ Use of long division is M0A0 as factor theorem was required. $f(-3) = 162 - 63 - 120 + 21 = 0$ so $(x + 3)$ is a factor	M1 A1 (2)
(b)	Either (Way 1): $f(x) = (x + 3)(-6x^2 + 11x + 7)$ $= (x + 3)(-3x + 7)(2x + 1)$ or $-(x + 3)(3x - 7)(2x + 1)$	M1A1 M1A1 (4)
	Or (Way 2) Uses trial or factor theorem to obtain $x = -1/2$ or $x = 7/3$ Uses trial or factor theorem to obtain both $x = -1/2$ and $x = 7/3$ Puts three factors together (see notes below) Correct factorisation : $(x + 3)(7 - 3x)(2x + 1)$ or $-(x + 3)(3x - 7)(2x + 1)$ oe	M1 A1 M1 A1 (4)
	Or (Way 3) No working three factors $(x + 3)(-3x + 7)(2x + 1)$ otherwise need working	M1A1M1A1 (4)
(c)	$2^y = \frac{7}{3}$, $\rightarrow \log(2^y) = \log\left(\frac{7}{3}\right)$ or $y = \log_2\left(\frac{7}{3}\right)$ or $\frac{\log(7/3)}{\log 2}$ $\{y = 1.222392421...\} \Rightarrow y = \text{awrt } 1.22$	B1, M1 A1 (3) [9]

Notes

- (a) M1 for attempting either $f(3)$ or $f(-3)$ – with **numbers substituted into expression**
A1 for calculating $f(-3)$ **correctly to 0**, and they must state $(x + 3)$ **is a factor** for A1 (or equivalent ie. QED, \square or a tick). A conclusion may be implied by a preamble, “if $f(-3) = 0$, $(x+3)$ is a factor”.
- $-6(-3)^3 - 7(-3)^2 + 40(-3) + 21 = 0$ so $(x + 3)$ is a factor of $f(x)$ is M1A1 providing bracketing is correct.
- (b) 1st M1: attempting to divide by $(x + 3)$ leading to a 3TQ beginning with the correct term, usually $-6x^2$. This may be done by a variety of methods including long division, comparison of coefficients, inspection etc. Allow for work in part (a) if the result is used in (b).
1st A1: usually for $(-6x^2 + 11x + 7) \dots$ Credit when seen and use isw if miscopied
2nd M1: for a **valid*** attempt to factorise their quadratic (* see notes on page 6 - General Principles for Core Mathematics Marking section 1)
2nd A1 is cao and needs all three factors together fully factorised. Accept e.g. $-3(x + 3)(x - \frac{7}{3})(2x + 1)$ but $(x + 3)(x - \frac{7}{3})(-6x - 3)$ and $(x + 3)(3x - 7)(-2x - 1)$ are A0 as not fully factorised.
Ignore subsequent work (such as a solution to a quadratic equation.)
Way 2: The second M mark needs three roots together so $\pm 6(x - \alpha)(x - \beta)(x + 3)$ or equivalent where they obtained α and β by trial, so if correct roots identified, then $(x + 3)(3x - 7)(2x + 1)$ can gain M1A1M1A0.
N.B. Replacing $(-6x^2 + 11x + 7)$ (already awarded M1A1) by $(6x^2 - 11x - 7)$ giving $(x + 3)(3x - 7)(2x + 1)$ can have M1A0 for factorization so M1A1M1A0
- (c) B1: $2^y = \frac{7}{3}$
M1: Attempt to take logs to solve $2^y = \alpha$ or $2^y = 1/\alpha$, where $\alpha > 0$ and α was a root of their factorization.
A1: for an answer that rounds to 1.22. If other answers are included (and not “rejected”) such as $\ln(-3)$ or -1 lose final A mark
- Special case:** Those who deal **throughout** with $f(x) = 6x^3 + 7x^2 - 40x - 21$
They may have full credit in part (a). In part (b) they can achieve a maximum of M1A0M1A0 unless they return the negative sign to give the correct answer. This is then full marks. Part (c) is fine. So they could lose 2 marks on the factorisation. (Like a misread)

Leave blank

7. (i) $2\log(x + a) = \log(16a^6)$, where a is a positive constant

Find x in terms of a , giving your answer in its simplest form. (3)

(ii) $\log_3(9y + b) - \log_3(2y - b) = 2$, where b is a positive constant

Find y in terms of b , giving your answer in its simplest form. (4)

Lined writing area for student response.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme	Marks
<p>7. (i)</p> <p>(ii) Way 1</p> <p>Way 2</p>	<p>Use of power rule so $\log(x+a)^2 = \log 16a^6$ or $2\log(x+a) = 2\log 4a^3$ or $\log(x+a) = \log(16a^6)^{\frac{1}{2}}$</p> <p>Removes logs and square roots, or halves then removes logs to give $(x+a) = 4a^3$</p> <p>Or $x^2 + 2ax + a^2 - 16a^6 = 0$ followed by factorisation or formula to give $x = \sqrt{16a^6} - a$</p> <p>$(x =) 4a^3 - a$ (depends on previous M's and must be this expression or equivalent)</p> <p>$\log_3 \frac{(9y+b)}{(2y-b)} = 2$ Applies quotient law of logarithms</p> <p>$\frac{(9y+b)}{(2y-b)} = 3^2$ Uses $\log_3 3^2 = 2$</p> <p>$(9y+b) = 9(2y-b) \Rightarrow y =$ Multiplies across and makes y the subject</p> <p>$y = \frac{10}{9}b$</p> <p>Or : $\log_3(9y+b) = \log_3 9 + \log_3(2y-b)$ 2nd M mark</p> <p>$\log_3(9y+b) = \log_3 9(2y-b)$ 1st M mark</p> <p>$(9y+b) = 9(2y-b) \Rightarrow y = \frac{10}{9}b$ Multiplies across and makes y the subject</p>	<p>M1</p> <p>M1</p> <p>A1cao (3)</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1cso (4)</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1cso (4)</p> <p>[7]</p>
Notes		
<p>(i)</p> <p>(ii)</p>	<p>1st M1: Applies power law of logarithms correctly to one side of the equation</p> <p>M1: Correct log work in correct order. If they square and obtain a quadratic the algebra should be correct. The marks is for $x+a = \sqrt{16a^6}$ isw so allow $x+a = \pm 4a^3$ for Method mark. Also allow $x+a = 4a^4$ or $x+a = \pm 4a^{5.5}$ or even $x+a = 16a^3$ as there is evidence of attempted square root.</p> <p>May see the correct $x+a = 10^{(\log 4 + 3\log a)}$ so $x = -a + 10^{(\log 4 + 3\log a)}$ which gains M1A0 unless followed by the answer in the scheme.</p> <p>A1: Do not allow $x = \pm 4a^3 - a$ for accuracy mark. You may see the factorised $a(2a+1)(2a-1)$ o.e.</p> <p>M1: Applying the subtraction or addition law of logarithms correctly to make two log terms into one log term in y</p> <p>M1: Uses $\log_3 3^2 = 2$</p> <p>3rd M1: Obtains correct linear equation in y usually the one in the scheme and attempts $y =$</p> <p>A1cso: $y = \frac{10}{9}b$ or correct equivalent after completely correct work.</p> <p>Special case:</p> <p>$\frac{\log_3(9y+b)}{\log_3(2y-b)} = 2$ is M0 unless clearly crossed out and replaced by the correct $\log_3 \frac{(9y+b)}{(2y-b)} = 2$</p> <p>Candidates may then write $\frac{(9y+b)}{(2y-b)} = 3^2$ and proceed to the <i>correct</i> answer – allow M0M1M1A0 as the answer requires a completely correct solution.</p>	

Leave
blank

8. (a) Show that the equation

$$\cos^2 x = 8 \sin^2 x - 6 \sin x$$

can be written in the form

$$(3 \sin x - 1)^2 = 2$$

(3)

(b) Hence solve, for $0 \leq x < 360^\circ$,

$$\cos^2 x = 8 \sin^2 x - 6 \sin x$$

giving your answers to 2 decimal places.

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme		Marks
<p>8. (a)</p>	<p>Way 1</p> $1 - \sin^2 x = 8\sin^2 x - 6\sin x$ <p>E.g. $9\sin^2 x - 6\sin x = 1$ or $9\sin^2 x - 6\sin x - 1 = 0$ or $9\sin^2 x - 6\sin x + 1 = 2$</p> <p>So $9\sin^2 x - 6\sin x + 1 = 2$ or $(3\sin x - 1)^2 - 2 = 0$ so $(3\sin x - 1)^2 = 2$ or $2 = (3\sin x - 1)^2$*</p>	<p>Way 2</p> $2 = (3\sin x - 1)^2$ gives $9\sin^2 x - 6\sin x + 1 = 2$ so $\sin^2 x + 8\sin^2 x - 6\sin x + 1 = 2$ <p>so $8\sin^2 x - 6\sin x = 1 - \sin^2 x$</p> $8\sin^2 x - 6\sin x = \cos^2 x$ *	<p>B1</p> <p>M1</p> <p>A1cso*</p> <p>(3)</p> <p>M1</p> <p>A1</p> <p>dM1A1 A1</p> <p>(5) [8]</p>
<p>(b)</p>	<p>Way 1: $(3\sin x - 1) = (\pm)\sqrt{2}$</p> $\sin x = \frac{1 \pm \sqrt{2}}{3}$ or awrt 0.8047 and awrt -0.1381 <p>$x = 53.58, 126.42$ (or 126.41), 352.06, 187.94</p>		<p>Way 2: Expands $(3\sin x - 1)^2 = 2$ and uses quadratic formula on 3TQ</p> <p>A1</p> <p>dM1A1 A1</p> <p>(5) [8]</p>
Notes			
<p>(a)</p>	<p>Way 1 B1: Uses $\cos^2 x = 1 - \sin^2 x$</p> <p>M1: Collects $\sin^2 x$ terms to form a three term quadratic or into a suitable completed square format. May be sign slips in the collection of terms. A1*: cso This needs an intermediate step from 3 term quadratic and no errors in answer and printed answer stated but allow $2 = (3\sin x - 1)^2$. If sin is used throughout instead of sinx it is A0.</p> <p>Way 2 B1: Needs correct expansion and split M1: Collects $1 - \sin^2 x$ together A1*: Conclusion and no errors seen</p>		
<p>(b)</p>	<p>M1: Square roots both sides(Way 1), or expands and uses quadratic formula (Way 2) Attempts at factorization after expanding are M0. A1: Both correct answers for sinx (need plus and minus). Need not be simplified. dM1: Uses inverse sin to give one of the given correct answers 1st A1: Need two correct angles (allow awrt) Note that the scheme allows 126.41 in place of 126.42 though 126.42 is preferred A1: All four solutions correct (Extra solutions in range lose this A mark, but outside range - ignore) (Premature approximation:- in the final three marks lose first A1 then ft other angles for second A mark) Do not require degrees symbol for the marks Special case: Working in radians M1A1A0 for the <i>correct</i> 0.94, 2.21, 6.14, 3.28</p>		

Question Number	Scheme	Marks
9.(a)	$a = 7k - 5, ar = 5k - 7$ and $ar^2 = 2k + 10$	B1
	(So $r =$) $\frac{5k - 7}{7k - 5} = \frac{2k + 10}{5k - 7}$ or $(7k - 5)(2k + 10) = (5k - 7)^2$ or equivalent	M1
	See $(5k - 7)^2 = 25k^2 - 70k + 49$	M1
	$14k^2 + 60k - 50 = 25k^2 - 70k + 49 \rightarrow 11k^2 - 130k + 99 = 0^*$	A1cso * (4)
(b)	$(k - 11)(11k - 9)$ so $k =$	M1
	$k = 9/11$ only* (after rejecting 11) N.B. Special case $k = 9/11$ can be verified in (b) (1 mark only)	A1*
	$11 \times \left(\frac{9}{11}\right)^2 - 130 \times \left(\frac{9}{11}\right) + 99 = \frac{81}{11} - \frac{1170}{11} + \frac{1089}{11} = 0$ M1A0	(2)
(c)	$a = \frac{8}{11}$	B1
	$\frac{5 \times \frac{9}{11} - 7}{7 \times \frac{9}{11} - 5}$ or $\frac{2 \times \frac{9}{11} + 10}{5 \times \frac{9}{11} - 7}$ so $r = -4$	B1
	(i) Fourth term = $ar^3 = -\frac{512}{11}$	M1A1
	(ii) $S_{10} = \frac{a(1 - r^{10})}{(1 - r)} = \frac{\frac{8}{11}(1 - (-4)^{10})}{(1 - (-4))} = -152520$	M1A1
		(6) [12]

Notes

(a) Mark parts (a) and (b) together

B1: Correct statement (needs all three terms)– **this may be omitted and implied** by correct statement in k only, as candidates may use geometric mean, or may use ratio of terms being equal and give a correct line 2 without line 1. (This would earn the B1M1 immediately)

M1: Valid Attempt to eliminate a and r and to obtain equation in k only

M1: Correct expansion of $(5k - 7)^2 = 25k^2 - 70k + 49$ - may have four terms $(5k - 7)^2 = 25k^2 - 35k - 35k + 49$

A1cso: No incorrect work seen. The printed answer is obtained including “=0”.

(b) M1: Attempt to solve quadratic by usual methods (factorisation, completion of square or formula – see notes at start of mark scheme) or see $9/11$ substituted and given as “=0” for M1A0

A1*: $9/11$ **only** and 11 should be seen and rejected. Accept $9/11$ underlined or $k = 9/11$ written on following line.

Alternatively $(k - 11)$ may be seen in the factorisation and a statement ‘ k not integer’ given with $k = 9/11$ stated.

(c) Mark parts (i) and (ii) together

B1: $a = \frac{8}{11}$ or any equivalent (If not stated explicitly or used in formula may be implied by correct answer to (ii))

B1: Substitutes $k = 9/11$ completely and obtain $r = -4$ (If not stated explicitly, may be implied by correct answer to (i) or (ii))

(i) M1: Use of correct formula with $n = 4$ a and/or r may still be in terms of k or uses $(2k + 10) \times r$. May assume $r = k$.

A1: Correct exact answer

(ii) M1: Use of correct formula with $n = 10$ a and/or r may still be in terms of k May assume $r = k$ A1: -152520 cao

NB Correct formula **with negative sign** in numerator followed by the incorrect $(8/11)(1 + 4^{10}) / (1 - (-4))$ usually found equal to 152520.2909 with no negative sign can be allowed M1A0 but if the incorrect numerical expression appears on its own with no formula then M0A0

Listing terms can get: B1 (first term) B1 M1A1 (implied by correct 4th term) M1A1 (implied by -152520)

Leave blank

10.




Figure 2

Figure 2 shows a sketch of part of the curve with equation

$$y = 4x^3 + 9x^2 - 30x - 8, \quad -0.5 \leq x \leq 2.2$$

The curve has a turning point at the point A .

- (a) Using calculus, show that the x coordinate of A is 1 (3)

The curve crosses the x -axis at the points $B(2, 0)$ and $C\left(-\frac{1}{4}, 0\right)$

The finite region R , shown shaded in Figure 2, is bounded by the curve, the line AB , and the x -axis.

- (b) Use integration to find the area of the finite region R , giving your answer to 2 decimal places. (7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question Number	Scheme	Marks
<p>10. (a)</p> <p>(b) Way 1</p>	$\frac{dy}{dx} = 12x^2 + 18x - 30$ <p>Either</p> <p>Substitute $x = 1$ to give $\frac{dy}{dx} = 12 + 18 - 30 = 0$</p> <p>So turning point (all correct work so far)</p> <p>When $x = 1$, $y = 4 + 9 - 30 - 8 = -25$</p> <p>Area of triangle $ABP = \frac{1}{2} \times 1 \times 25 = 12.5$ (Where P is at $(1, 0)$)</p> <p>Way 1: $\int (4x^3 + 9x^2 - 30x - 8) dx = x^4 + \frac{9}{3}x^3 - \frac{30x^2}{2} - 8x \{+ c\}$ or $x^4 + 3x^3 - 15x^2 - 8x \{+ c\}$</p> $\left[x^4 + 3x^3 - 15x^2 - 8x \right]_{-\frac{1}{4}}^1 = (1 + 3 - 15 - 8) - \left(\left(-\frac{1}{4} \right)^4 + 3 \left(-\frac{1}{4} \right)^3 - 15 \left(-\frac{1}{4} \right)^2 - 8 \left(-\frac{1}{4} \right) \right)$ $= (-19) - \frac{261}{256} \text{ or } -19 - 1.02$ <p>So Area = "their 12.5" + "their 20 $\frac{5}{256}$" or "12.5" + "20.02" or "12.5" + "their $\frac{5125}{256}$"</p> <p>= 32.52 (NOT - 32.52)</p>	<p>M1</p> <p>A1</p> <p>A1cso (3)</p> <p>B1</p> <p>B1</p> <p>M1A1</p> <p>dM1</p> <p>ddM1</p> <p>A1 (7) [10]</p>
	<p>Less efficient alternative methods for first two marks in part (b) with Way 1 or 2</p> <p>For first mark: Finding equation of the line AB as $y = 25x - 50$ as this implies the -25</p> <p>For second mark: Integrating to find triangle area</p> $\int_1^2 (25x - 50) dx = \left[\frac{25}{2}x^2 - 50x \right]_1^2 = -50 + 37.5 = -12.5 \quad \text{so area is } 12.5$ <p>Then mark as before if they use Method in original scheme</p>	<p>B1</p> <p>B1</p>
<p>(b) Way 2</p>	<p>Way 2: Those who use area for original curve between -1/4 and 2 and subtract area between line and curve between 1 and 2 have a correct (long) method.</p> <p>The first B1 (if $y = -25$ is not seen) is for equation of straight line $y = 25x - 50$</p> <p>The second B1 may be implied by final answer correct, or 4.5 seen for area of "segment shaped" region between line and curve, or by area between line and axis/triangle found as 12.5</p> $\int (4x^3 + 9x^2 - 55x + 42) dx = x^4 + \frac{9}{3}x^3 - \frac{55x^2}{2} + 42x \{+ c\} \text{ (or integration as in Way 1)}$ <p>The dM1 is for correct use of the different correct limits for each of the two areas: i.e.</p> $\left[x^4 + 3x^3 - 15x^2 - 8x \right]_{-\frac{1}{4}}^2 = (16 + 24 - 60 - 16) - \left(\left(-\frac{1}{4} \right)^4 + 3 \left(-\frac{1}{4} \right)^3 - 15 \left(-\frac{1}{4} \right)^2 - 8 \left(-\frac{1}{4} \right) \right)$ <p>And $\left[x^4 + \frac{9}{3}x^3 - \frac{55x^2}{2} + 42x \right]_1^2 = 16 + 24 - 110 + 84 - (1 + 3 - 27.5 + 42)$</p> <p>So Area = their $\left[x^4 + 3x^3 - 15x^2 - 8x \right]_{-\frac{1}{4}}^2$ minus their $\left[x^4 + \frac{9}{3}x^3 - \frac{55x^2}{2} + 42x \right]_1^2$</p> <p>i.e. "their 37.0195" - "their 4.5" (with both sets of limits correct for the integral)</p> <p>Reaching = 32.52 (NOT - 32.52)</p> <p>See over for special case with wrong limits</p>	<p>B1</p> <p>B1</p> <p>M1A1</p> <p>dM1</p> <p>ddM1</p> <p>A1</p>

<p>NB: Those who attempt curve – line wrongly with limits $-1/4$ to 2 may earn M1A1 for correct integration of their cubic. Usually e.g.</p> $\int (4x^3 + 9x^2 - 55x + 42)dx = x^4 + \frac{9}{3}x^3 - \frac{55x^2}{2} + 42x \{+ c\}$ <p>(They will not earn any of the last 3 marks)</p> <p>They may also get first B1 mark for the correct equation of the straight line (usually seen but may be implied by correct line –curve equation) and second B1 if they also use limits 1 and 2 to obtain 4.5 (or find the triangle area 12.5).</p>	<p>M1A1</p>
---	-------------

<p>Notes</p>	
<p>(a)</p> <p>(b)</p>	<p>M1: Attempt at differentiation - all powers reduced by 1 with $8 \rightarrow 0$.</p> <p>A1: the derivative must be correct and uses derivative = 0 to find x or substitutes $x = 1$ to give 0. Ignore any reference to the other root ($-5/2$) for this mark.</p> <p>A1cso: obtains $x = 1$ from correct work, or deduces turning point (if substitution used – may be implied by a preamble e.g. $dy/dx = 0$ at T.P.)</p> <p>N.B. If their factorisation or their second root is incorrect then award A0cso.</p> <p>If however their factorisation/roots are correct, it is not necessary for them to comment that -2.5 is outside the range given.</p> <p>Way 1:</p> <p>B1: Obtains $y = -25$ when $x = 1$ (may be seen anywhere – even in (a)) or finds correct equation of line is $y = 25x - 50$</p> <p>B1: Obtains area of triangle = 12.5 (may be seen anywhere). Allow -12.5. Accept $\frac{1}{2} \times 1 \times 25$</p> <p>M1: Attempt at integration of cubic; two correct terms for their integration. No limits needed</p> <p>A1: completely correct integral for the cubic (may be unsimplified)</p> <p>dM1: We are looking for the start of a correct method here (dependent on previous M). It is for substituting 1 and $-1/4$ and subtracting. May use 2 and $-1/4$ and also 2 and 1 AND subtract (which is equivalent)</p> <p>ddM1 (depends on both method marks) Correct method to obtain shaded area so adds two positive numbers (areas) together – one is area of triangle, the other is area of region obtained from integration of correct function with correct limits (may add two negatives then makes positive)</p> <p>Way 2: This is a long method and needs to be a correct method</p> <p>B1: Finds $y = -25$ at $x = 1$, or correct equation of line is $y = 25x - 50$</p> <p>B1: May be implied where WAY 2 is used and final correct answer obtained so award of final A1 results in the award of this B1. It may also be implied by correct integration of line equation or of curve minus line expression between limits 1 and 2. So if only slip is final subtraction (giving final A0, this mark may still be awarded) So may be implied by 4.5 seen for area of “segment shaped” region between line and curve.</p> <p>M1: Attempt at integration of given cubic or after attempt at subtracting their line equation (no limits needed). Two correct terms needed</p> <p>A1: Completely correct integral for their cubic (may be unsimplified) – may have wrong coefficients of x and wrong constant term through errors in subtraction</p> <p>dM1: Use limits for original curve between $-1/4$ and 2 and use limits of 1 and 2 for area between line and curve– needs completely correct limits– see scheme- this is dependent on two integrations</p> <p>ddM1: (depends on both method marks) Subtracts “<i>their 37.0195</i>” – “<i>their 4.5</i>” Needs consistency of signs.</p> <p>A1: 32.52 or awrt 32.52 e.g. $32 \frac{133}{256}$ NB: This correct answer implies the second B mark</p> <p>(Trapezium rule gets no marks after first two B marks) The first two B marks may be given wherever seen. The integration of a cubic gives the following M1 and correct integration of their cubic</p> $\int (4x^3 + 9x^2 + Ax + B)dx = x^4 + \frac{9}{3}x^3 + \frac{Ax^2}{2} + Bx \{+ c\}$ gives the A1