



Leave  
blank

1. The first three terms of a geometric series are

$$18, 12 \text{ and } p$$

respectively, where  $p$  is a constant.

Find

(a) the value of the common ratio of the series, (1)

(b) the value of  $p$ , (1)

(c) the sum of the first 15 terms of the series, giving your answer to 3 decimal places. (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>1. (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	$\{r = \} \frac{2}{3}$ $\{p = \} 8$ $\{S_{15} = \} \frac{18(1 - (\frac{2}{3})^{15})}{1 - \frac{2}{3}}$ $\{S_{15} = 53.87668...\} \Rightarrow S_{15} = \text{awrt } 53.877$	<p>B1</p> <p>(1)</p> <p>B1 cao</p> <p>(1)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>[4]</p>
<b>Notes for Question 1</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	<p>B1: Accept <math>\frac{12}{18}</math>, 0.6 or 0.6 recurring, or even 0.667 (3sf) but not 0.6 or 0.67</p> <p>B1: accept 8 only</p> <p>M1: Applies this formula <math>S_{15} = \frac{18(1 - (\text{their } r)^{15})}{1 - (\text{their } r)}</math>, can be implied by their answer. For this mark they may use any value for <math>r</math> except <math>r = 1</math> or <math>r = 0</math> (even <math>3/2</math> or <math>-6</math> may be used)</p> <p>A1: Answers which round to 53.877</p>	
<p><b>Alternative method for (c)</b></p>	<p>M1: (Adding terms is an unlikely method for this question) Need to see 15 terms listed as <math>18+12+\dots+0.06165877</math> or can be implied by correct answer</p> <p>A1: awrt 53.877</p> <p><b>Answer only</b> : 53.9 is M0A0 with no working, but 53.877 with no working is M1A1</p>	



Question Number	Scheme	Marks
<p><b>2. (a)</b></p>	<p><math>(2 + 3x)^4</math> - Mark (a) and (b) together  <math>2^4 + {}^4C_1 2^3(3x) + {}^4C_2 2^2(3x)^2 + {}^4C_3 2^1(3x)^3 + (3x)^4</math>                      First term of 16  <math>({}^4C_1 \times \dots \times x) + ({}^4C_2 \times \dots \times x^2) + ({}^4C_3 \times \dots \times x^3) + ({}^4C_4 \times \dots \times x^4)</math>  <math>= (16 + ) 96x + 216x^2 + 216x^3 + 81x^4</math>                      Must use Binomial – otherwise A0,                      A0</p>	<p>B1                      M1                      A1 A1                        (4)                      B1ft                      (1)                      5</p>
<p><b>Alternative method (a)</b></p>	<p><math>(2 + 3x)^4 = 2^4(1 + \frac{3x}{2})^4</math>  <math>2^4(1 + {}^4C_1(\frac{3x}{2}) + {}^4C_2(\frac{3x}{2})^2 + {}^4C_3(\frac{3x}{2})^3 + (\frac{3x}{2})^4)</math>                      Scheme is applied exactly as before</p>	
<b>Notes for Question 2</b>		
<p><b>(a)</b></p>	<p>B1: The constant term should be 16 in their expansion                      M1: Two binomial coefficients must be correct and must be with the correct power of <math>x</math>. Accept <math>{}^4C_1</math> or <math>\binom{4}{1}</math> or 4 as a coefficient, and <math>{}^4C_2</math> or <math>\binom{4}{2}</math> or 6 as another..... Pascal's triangle may be used to establish coefficients.                      A1: Any two of the final four terms correct (i.e. two of <math>96x + 216x^2 + 216x^3 + 81x^4</math>) in expansion following Binomial Method.                      A1: All four of the final four terms correct in expansion. (Accept answers without + signs, can be listed with commas or appear on separate lines)</p>	
<p><b>(b)</b></p>	<p>B1ft: Award for correct answer as printed above or <b>ft their previous answer</b> provided it has <b>five terms</b> ft and must be <b>subtracting the <math>x</math> and <math>x^3</math> terms</b>                      Allow terms in (b) to be in descending order and allow <math>+96x</math> and <math>+216x^3</math> in the series. (Accept answers without + signs, can be listed with commas or appear on separate lines)</p>	
<p>e.g. The common error <math>2^4 + {}^4C_1 2^3 3x + {}^4C_2 2^2 3x^2 + {}^4C_3 2^1 3x^3 + 3x^4 = (16) + 96x + 72x^2 + 24x^3 + 3x^4</math> would earn B1, M1, A0, A0, and if followed by <math>= (16) - 96x + 72x^2 - 24x^3 + 3x^4</math> gets B1ft so 3/5  <b>Fully correct answer with no working can score B1 in part (a) and B1 in part (b).</b> The question stated use the Binomial theorem and if there is no evidence of its use then M mark and hence A marks cannot be earned.  <b>Squaring the bracket and squaring again</b> may also earn <b>B1 M0 A0 A0 B1 if correct</b>  <b>Omitting the final term</b> but otherwise correct is B1 M1 A1 A0 B0ft so 3/5                      If the series is divided through by 2 or a power of 2 at the final stage after an error or omission resulting in all even coefficients then apply scheme to series before this division and ignore subsequent work (isw)</p>		

Leave blank

3.  $f(x) = 2x^3 - 5x^2 + ax + 18$

where  $a$  is a constant.

Given that  $(x - 3)$  is a factor of  $f(x)$ ,

(a) show that  $a = -9$  (2)

(b) factorise  $f(x)$  completely. (4)

Given that

$$g(y) = 2(3^{3y}) - 5(3^{2y}) - 9(3^y) + 18$$

(c) find the values of  $y$  that satisfy  $g(y) = 0$ , giving your answers to 2 decimal places where appropriate. (3)

Handwritten solution area with horizontal lines.



Question Number	Scheme	Marks
3. (a)	Either (Way 1) : Attempt $f(3)$ or $f(-3)$ $f(3) = 54 - 45 + 3a + 18 = 0 \Rightarrow 3a = -27 \Rightarrow a = -9^*$	M1 A1 * cso <b>(2)</b>
	Or (Way 2): Assume $a = -9$ and attempt $f(3)$ or $f(-3)$ $f(3) = 0$ so $(x - 3)$ is factor	M1 A1* cso <b>(2)</b>
(b)	Either (Way 1): $f(x) = (x - 3)(2x^2 + x - 6)$ $= (x - 3)(2x - 3)(x + 2)$	M1A1 M1A1 <b>(4)</b>
	Or (Way 2) Uses trial or factor theorem to obtain $x = -2$ <b>or</b> $x = 3/2$ Uses trial or factor theorem to obtain both $x = -2$ <b>and</b> $x = 3/2$ Puts three factors together (see notes below) Correct factorisation : $(x - 3)(2x - 3)(x + 2)$ or $(3 - x)(3 - 2x)(x + 2)$ or $2(x - 3)(x - \frac{3}{2})(x + 2)$ oe	M1 A1 M1 A1 <b>(4)</b>
	Or (Way 3) No working three factors $(x - 3)(2x - 3)(x + 2)$ otherwise need working	M1A1M1A1
(c)	$\{3^y = 3 \Rightarrow\} \underline{y = 1}$ or $g(1) = 0$	B1
	$\{3^y = 1.5 \Rightarrow\} \log(3^y) = \log 1.5$ or $y = \log_3 1.5$	M1
	$\{y = 0.3690702\dots\} \Rightarrow y = \text{awrt } 0.37$	A1 (3) <b>[9]</b>
<b>Notes for Question 3</b>		
(a)	M1 for attempting either $f(3)$ or $f(-3)$ – with <b>numbers substituted into expression</b> A1 for applying $f(3)$ <b>correctly</b> , setting the result <b>equal to 0</b> , and manipulating this correctly to give the result given on the paper i.e. $a = -9$ . (Do not accept $x = -9$ ) Note that the answer is given in part (a). If they <b>assume</b> $a = -9$ and <b>verify</b> by factor theorem or division they must state $(x - 3)$ <b>is a factor</b> for A1 (or equivalent such as QED or a tick).	
(b)	1 <sup>st</sup> M1: attempting to divide by $(x - 3)$ leading to a 3TQ beginning with the correct term, usually $2x^2$ . (Could divide by $(3 - x)$ , in which case the quadratic would begin $-2x^2$ .) This may be done by a variety of methods including long division, comparison of coefficients, inspection etc. 1 <sup>st</sup> A1: usually for $2x^2 + x - 6 \dots$ Credit when seen and use isw if miscopied 2 <sup>nd</sup> M1: for a <b>valid</b> * attempt to factorise their quadratic (* see notes on page 6 - General Principles for Core Mathematics Marking section 1) 2 <sup>nd</sup> A1 is cao and needs all three factors together. Ignore subsequent work (such as a solution to a quadratic equation.) NB: $(x - 3)(x - \frac{3}{2})(x + 2)$ is M1A1M0A0, $(x - 3)(x - \frac{3}{2})(2x + 4)$ is M1A1M1A0, but $2(x - 3)(x - \frac{3}{2})(x + 2)$ is M1A1M1A1.	
(c)	B1: $\underline{y = 1}$ seen as a solution – may be spotted as answer – no working needed. Allow also for $g(1) = 0$ . M1: Attempt to take logs to solve $3^y = \alpha$ or even $3^{ky} = \alpha$ , but not $6^y = \alpha$ where $\alpha > 0$ and $\alpha \neq 3$ & was a root of $f(x) = 0$ (ft their factorization) A1: for an answer that rounds to 0.37. If a third answer is included (and not “rejected”) such as $\ln(-2)$ lose final A mark	

4. 
$$y = \frac{5}{(x^2 + 1)}$$

(a) Complete the table below, giving the missing value of  $y$  to 3 decimal places.

$x$	0	0.5	1	1.5	2	2.5	3
$y$	5	4	2.5		1	0.690	0.5

(1)




Figure 1

Figure 1 shows the region  $R$  which is bounded by the curve with equation  $y = \frac{5}{(x^2 + 1)}$ , the  $x$ -axis and the lines  $x = 0$  and  $x = 3$

(b) Use the trapezium rule, with all the values of  $y$  from your table, to find an approximate value for the area of  $R$ .

(4)

(c) Use your answer to part (b) to find an approximate value for

$$\int_0^3 \left( 4 + \frac{5}{(x^2 + 1)} \right) dx$$

giving your answer to 2 decimal places.

(2)

---



---



---



---





Question Number	Scheme	Marks																
<p>4.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<table border="1" data-bbox="209 304 1243 376"> <tr> <td>x</td> <td>0</td> <td>0.5</td> <td>1</td> <td>1.5</td> <td>2</td> <td>2.5</td> <td>3</td> </tr> <tr> <td>y</td> <td>5</td> <td>4</td> <td>2.5</td> <td>1.538</td> <td>1</td> <td>0.690</td> <td>0.5</td> </tr> </table> <p>{At <math>x = 1.5,</math>} <math>y = 1.538</math> (only)</p> <p><math>\frac{1}{2} \times 0.5;</math></p> <p><math>\{5 + 0.5 + 2(4 + 2.5 + \text{their } 1.538 + 1 + 0.690)\}</math>      <u>For structure of</u> <math>\{.....\};</math></p> <p><math>\frac{1}{2} \times 0.5 \times \{ (5 + 0.5) + 2(4 + 2.5 + \text{their } 1.538 + 1 + 0.690) \} = \frac{1}{4}(24.956) = 6.239 = \text{awrt } 6.24</math></p> <p>Adds Area of Rectangle or first integral = <math>3 \times 4</math> or <math>[4x]_0^3</math> to <b>previous answer</b></p> <p>So required estimate = <math>\{ "6.239" + 12 = "18.239" \} = \text{"awrt } 18.24"</math> (or <math>12 + \text{previous answer}</math>).</p> <p>N.B. <math>7 \times 4 + \text{previous answer}</math> is M0A0 (added 4 seven times because 7 numbers in table)</p>	x	0	0.5	1	1.5	2	2.5	3	y	5	4	2.5	1.538	1	0.690	0.5	<p>B1 cao</p> <p>[1]</p> <p>B1 oe</p> <p>M1A1ft</p> <p>A1</p> <p>[4]</p> <p>M1</p> <p>A1ft</p> <p>[2]</p> <p>7</p>
x	0	0.5	1	1.5	2	2.5	3											
y	5	4	2.5	1.538	1	0.690	0.5											
<b>Notes for Question 4</b>																		
<p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>B1: 1.538</p> <p>B1: for using <math>\frac{1}{2} \times 0.5</math> or <math>\frac{1}{4}</math> or equivalent.</p> <p>M1: requires the correct <math>\{.....\}</math> bracket structure. It needs the first bracket to contain first y value <b>plus</b> last y value and the second bracket to be multiplied by 2 and to be the summation of the remaining y values in the table with no additional values. If the only mistake is a copying error or is to omit one value from 2nd bracket this may be regarded as a slip and the M mark can be allowed ( An extra repeated term forfeits the M mark however). M0 if values used in brackets are x values instead of y values</p> <p>A1ft: for the correct bracket <math>\{.....\}</math> following through candidate's y value found in part (a).</p> <p>A1: for answer which rounds to 6.24.</p> <p>NB: Separate trapezia may be used : B1 for 0.25, M1 for <math>\frac{1}{2}h(a + b)</math> used 5 or 6 times (and A1ft if it is all correct ) Then A1 as before.</p> <p>Special case: Bracketing mistake <math>0.25 \times (5 + 0.5) + 2(4 + 2.5 + \text{their } 1.538 + 1 + 0.690)</math> scores B1 M1 A0</p> <p>A0 unless the final answer implies that the calculation has been done correctly (then full marks can be given). An answer of 20.831 usually indicates this error.</p> <p>M1: Relates <b>previous answer ( not integral of previous answer)</b> to this question by integrating 4 between limits, and adding, or by using geometry to find rectangle and adding.</p> <p>A1ft: for <math>12 + \text{answer to (b)}</math></p>																	
<p>Alternative method</p> <p>(c)</p>	<p>Those who do a trapezium rule for part (b)- using the table from (a) with 4 added to each cell of the table</p> <p>Get: M1 for <math>"\text{their } \frac{1}{4}" \times \{ 9 + 4.5 + 2(8 + 6.5 + \text{their } 5.538 + 5 + 4.690) \} = (\text{structure must be correct} - \text{allow one copying error only})</math></p> <p>And A1ft: for awrt 18.24 (or <math>12 + \text{previous answer}</math>).</p>																	

5.




Figure 2

Figure 2 shows a plan view of a garden.  
The plan of the garden *ABCDEA* consists of a triangle *ABE* joined to a sector *BCDE* of a circle with radius 12m and centre *B*.  
The points *A*, *B* and *C* lie on a straight line with  $AB = 23\text{ m}$  and  $BC = 12\text{ m}$ .

Given that the size of angle *ABE* is exactly 0.64 radians, find

(a) the area of the garden, giving your answer in  $\text{m}^2$ , to 1 decimal place, (4)

(b) the perimeter of the garden, giving your answer in metres, to 1 decimal place. (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p>5. (a)</p>	<p>Mark (a) and (b) together.</p> <p><b>Usually answered in radians:</b> Uses either <math>\frac{1}{2}ab\sin(\text{angle})</math> or <math>\frac{1}{2}(12)^2(\text{angle})</math> or both</p> <p>Area = <math>\frac{1}{2}(23)(12)\sin 0.64</math> or <math>\frac{1}{2}(12)^2(\pi - 0.64)</math> {= 82.41297091... or 180.1146711...}</p> <p>Area = <math>\frac{1}{2}(23)(12)\sin 0.64 + \frac{1}{2}(12)^2(\pi - 0.64)</math> {= 82.41297091... + 180.1146711...}</p> <p>{Area = 262.527642...} = awrt 262.5 (m<sup>2</sup>) or 262.4(m<sup>2</sup>) or 262.6 (m<sup>2</sup>)</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>(4)</p>
	<p><b>Notes for Question 5</b></p>	
<p>(a)</p>	<p>M1: uses either area of triangle formula as given in mark scheme, or area of sector or both (may be implied by answer)</p> <p>A1: one correct area expression (with <b>correct angle</b> used) <math>\frac{1}{2}(23)(12)\sin 0.64</math> or <math>\frac{1}{2}(12)^2(\pi - 0.64)</math> or see awrt 82.4 or awrt 180 (180 may be split as 226.2(semicircle) minus 46.1(small sector))</p> <p>A1: two correct area expressions (with correct angles) <b>added together</b> (allow 2.5 as implying <math>\pi - 0.64</math>) or see awrt 82.4 + awrt 180 ( or 226 - 46 )</p>	<p>M1, A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(5)</p> <p>[9]</p>
<p><b>Degrees</b></p>		
<p>(a)</p>	<p>Uses either <math>\frac{1}{2}ab\sin(\text{angle})</math> or <math>\frac{\text{angle in degrees}}{360} \times \pi(12)^2</math> or both for M1</p> <p>Area = <math>\frac{1}{2}(23)(12)\sin 36.7</math> or <math>\frac{(180-36.7)}{360} \times \pi(12)^2</math> {= awrt 82.4... or 180} A1</p> <p>Area = <math>\frac{1}{2}(23)(12)\sin 36.7 + \frac{(180-36.7)}{360} \times \pi(12)^2</math> {= awrt 82.4... + 180} A1</p> <p>Final mark as before</p>	
<p>(b)</p>	<p><math>CDE = \frac{\text{Angle in degrees}}{360} \times 24\pi = \frac{180-36.7}{360} \times 24\pi</math> { <math>\Rightarrow CDE = 30.01268...</math> } M1, A1</p> <p>Final three marks as before</p>	

6.




Figure 3

Figure 3 shows a sketch of part of the curve  $C$  with equation

$$y = x(x + 4)(x - 2)$$

The curve  $C$  crosses the  $x$ -axis at the origin  $O$  and at the points  $A$  and  $B$ .

- (a) Write down the  $x$ -coordinates of the points  $A$  and  $B$ . (1)

The finite region, shown shaded in Figure 3, is bounded by the curve  $C$  and the  $x$ -axis.

- (b) Use integration to find the total area of the finite region shown shaded in Figure 3. (7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>6. (a)</b></p> <p><b>(b)</b></p>	<p>Seeing <math>-4</math> and <math>2</math>.</p> <p><math>x(x+4)(x-2) = x^3 + 2x^2 - 8x</math> or <math>x^3 - 2x^2 + 4x^2 - 8x</math> (without simplifying)</p> <p><math>\int (x^3 + 2x^2 - 8x) dx = \frac{x^4}{4} + \frac{2x^3}{3} - \frac{8x^2}{2} \{+c\}</math> or <math>\frac{x^4}{4} - \frac{2x^3}{3} + \frac{4x^3}{3} - \frac{8x^2}{2} \{+c\}</math></p> <p><math>\left[ \frac{x^4}{4} + \frac{2x^3}{3} - \frac{8x^2}{2} \right]_{-4}^0 = (0) - \left( 64 - \frac{128}{3} - 64 \right)</math> or <math>\left[ \frac{x^4}{4} + \frac{2x^3}{3} - \frac{8x^2}{2} \right]_0^2 = \left( 4 + \frac{16}{3} - 16 \right) - (0)</math></p> <p>One integral <math>= \pm 42\frac{2}{3}</math> (42.6 or awrt 42.7) or other integral <math>= \pm 6\frac{2}{3}</math> (6.6 or awrt 6.7)</p> <p>Hence Area = "their <math>42\frac{2}{3}</math>" + "their <math>6\frac{2}{3}</math>" or Area = "their <math>42\frac{2}{3}</math>" - "their <math>6\frac{2}{3}</math>"</p> <p><math>= 49\frac{1}{3}</math> or 49.3 or <math>\frac{148}{3}</math> (NOT <math>-\frac{148}{3}</math>)</p> <p>(An answer of <math>= 49\frac{1}{3}</math> may not get the final two marks – check solution carefully)</p>	<p>B1 (1)</p> <p><u>B1</u></p> <p>M1A1ft</p> <p>dM1</p> <p>A1</p> <p>dM1</p> <p>A1</p> <p>(7)</p> <p><b>[8]</b></p>
<b>Notes for Question 6</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p>	<p>B1: Need both <math>-4</math> and <math>2</math>. May see <math>(-4,0)</math> and <math>(2,0)</math> (correct) but allow <math>(0,-4)</math> and <math>(0,2)</math> or <math>A = -4, B = 2</math> or indeed any indication of <math>-4</math> and <math>2</math> – check graph also</p> <p>B1: Multiplies out cubic correctly (terms may not be collected, but if they are, mark collected terms here)</p> <p>M1: Tries to integrate their expansion with <math>x^n \rightarrow x^{n+1}</math> for at least one of the terms</p> <p>A1ft: completely correct integral <b>following through</b> from their CUBIC expansion (if only quadratic or quartic this is A0)</p> <p>dM1: (dependent on previous M) substituting EITHER <math>-a</math> and <math>0</math> and subtracting either way round OR similarly for <math>0</math> and <math>b</math>. <b>If their limits <math>-a</math> and <math>b</math> are used in ONE integral, apply the Special Case below.</b></p> <p>A1: Obtain <b>either</b> <math>\pm 42\frac{2}{3}</math> (or 42.6 or awrt 42.7) <b>from the integral from <math>-4</math> to <math>0</math> or <math>\pm 6\frac{2}{3}</math></b> (6.6 or awrt 6.7) <b>from the integral from <math>0</math> to <math>2</math></b>; NO follow through on their cubic (allow decimal or improper equivalents <math>\frac{128}{3}</math> or <math>\frac{20}{3}</math>) isw such as subtracting from rectangles. This will be penalized in the next two marks, which will be M0A0.</p> <p>dM1 (depends on first method mark) <b>Correct method to obtain shaded area</b> so adds two positive numbers (areas) together or uses their <b>positive</b> value minus <b>their negative</b> value, <b>obtained from</b> two separate <b>definite integrals</b>.</p> <p>A1: Allow 49.3, 49.33, 49.333 etc. Must follow correct logical work with <b>no errors</b> seen.</p> <p>For full marks on this question there must be <b>two</b> definite integrals, from <math>-4</math> to <math>0</math> and from <math>0</math> to <math>2</math>, though the evaluations for <math>0</math> may not be seen.</p> <p>(Trapezium rule <b>gets no marks after first two B marks</b>)</p>	
<p><b>(b)</b></p>	<p><b>Special Case: one integral only from <math>-a</math> to <math>b</math>:</b> B1M1A1 available as before, then</p> <p><math>\left[ \frac{x^4}{4} + \frac{2x^3}{3} - \frac{8x^2}{2} \right]_{-4}^2 = \left( 4 + \frac{16}{3} - 16 \right) - \left( 64 - \frac{128}{3} - 64 \right) = -6\frac{2}{3} + 42\frac{2}{3} = \dots</math> <b>dM1</b> for correct use of their limits <math>-a</math> and <math>b</math> and <b>subtracting</b> either way round.</p> <p>A1 for 36: NO follow through. Final M and A marks not available. Max 5/7 for part (b)</p>	

Leave  
blank

7. (i) Find the exact value of  $x$  for which

$$\log_2(2x) = \log_2(5x + 4) - 3 \quad (4)$$

- (ii) Given that

$$\log_a y + 3\log_a 2 = 5$$

express  $y$  in terms of  $a$ .  
Give your answer in its simplest form. (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>7. (i)</b> <b>Method 1</b></p>	<p><math>\log_2\left(\frac{2x}{5x+4}\right) = -3</math> or <math>\log_2\left(\frac{5x+4}{2x}\right) = 3</math>, or <math>\log_2\left(\frac{5x+4}{x}\right) = 4</math> (see special case 2)</p> <p><math>\left(\frac{2x}{5x+4}\right) = 2^{-3}</math> or <math>\left(\frac{5x+4}{2x}\right) = 2^3</math> or <math>\left(\frac{5x+4}{x}\right) = 2^4</math> or <math>\left(\log_2\left(\frac{2x}{5x+4}\right)\right) = \log_2\left(\frac{1}{8}\right)</math></p> <p><math>16x = 5x + 4 \Rightarrow x =</math> (depends on previous Ms and must be this equation or equivalent)</p> <p><math>x = \frac{4}{11}</math> or exact recurring decimal <math>0.\dot{3}\dot{6}</math> after correct work</p>	<p>M1</p> <p>M1</p> <p>dM1</p> <p>A1 <b>cs</b> <b>(4)</b></p>
<p><b>7(i)</b> <b>Method 2</b></p>	<p><math>\log_2(2x) + 3 = \log_2(5x + 4)</math></p> <p>So <math>\log_2(2x) + \log_2(8) = \log_2(5x + 4)</math> (3 replaced by <math>\log_2 8</math>)</p> <p>Then <math>\log_2(16x) = \log_2(5x + 4)</math> (addition law of logs)</p> <p>Then final M1 A1 as before</p>	<p>2<sup>nd</sup> M1</p> <p>1<sup>st</sup> M1</p> <p>dM1A1</p>
<p><b>(ii)</b></p>	<p><math>\log_a y + \log_a 2^3 = 5</math></p> <p><math>\log_a 8y = 5</math> Applies product law of logarithms.</p> <p><math>y = \frac{1}{8}a^5</math> <math>y = \frac{1}{8}a^5</math></p>	<p>M1</p> <p>dM1</p> <p>A1cao <b>(3)</b> <b>[7]</b></p>
<b>Notes for Question 7</b>		
<p><b>(i)</b></p>	<p>1<sup>st</sup> M1: Applying the subtraction or addition law of logarithms correctly to make <b>two log terms in x into one log term in x</b></p> <p>2<sup>nd</sup> M1: For RHS of either <math>2^{-3}</math>, <math>2^3</math>, <math>2^4</math> or <math>\log_2\left(\frac{1}{8}\right)</math>, <math>\log_2 8</math> or <math>\log_2 16</math> i.e. using connection between log base 2 and 2 to a power. This may follow an earlier error. <b>Use of <math>3^2</math> is M0</b></p> <p>3<sup>rd</sup> dM1: Obtains <b>correct</b> linear equation in x. usually the one in the scheme and attempts <math>x =</math></p> <p>A1: cso Answer of <math>4/11</math> with <b>no</b> suspect log work preceding this.</p>	
<p><b>(ii)</b></p>	<p>M1: Applies power law of logarithms to replace <math>3\log_a 2</math> by <math>\log_a 2^3</math> or <math>\log_a 8</math></p> <p>dM1: (should not be following M0) Uses addition law of logs to give <math>\log_a 2^3 y = 5</math> or <math>\log_a 8y = 5</math></p>	
<p><b>(i)</b></p>	<p><b>Special case 1:</b> <math>\log_2(2x) = \log_2(5x + 4) - 3 \Rightarrow \frac{\log_2(2x)}{\log_2(5x + 4)} = -3 \Rightarrow \frac{2x}{5x + 4} = 2^{-3} \Rightarrow x = \frac{4}{11}</math> or</p> <p><math>\log_2(2x) = \log_2(5x + 4) - 3 \Rightarrow \frac{\log_2(2x)}{\log_2(5x + 4)} = -3 \Rightarrow \log_2 \frac{2x}{5x + 4} = -3 \Rightarrow \frac{2x}{5x + 4} = 2^{-3} \Rightarrow x = \frac{4}{11}</math> each</p> <p>attempt scores M0M1M1A0 – special case</p> <p><b>Special case 2:</b></p> <p><math>\log_2(2x) = \log_2(5x + 4) - 3 \Rightarrow \log_2 2 + \log_2 x = \log_2(5x + 4) - 3</math>, is M0 until the two log terms are combined to give <math>\log_2\left(\frac{5x+4}{x}\right) = 3 + \log_2 2</math>. This earns M1</p> <p>Then <math>\left(\frac{5x+4}{x}\right) = 2^4</math> or <math>\log_2\left(\frac{5x+4}{x}\right) = \log_2 2^4</math> gets second M1. Then scheme as before.</p>	

Leave blank

8. (i) Solve, for  $-180^\circ \leq x < 180^\circ$ ,

$$\tan(x - 40^\circ) = 1.5$$

giving your answers to 1 decimal place.

(3)

(ii) (a) Show that the equation

$$\sin\theta \tan\theta = 3\cos\theta + 2$$

can be written in the form

$$4\cos^2\theta + 2\cos\theta - 1 = 0$$

(3)

(b) Hence solve, for  $0 \leq \theta < 360^\circ$ ,

$$\sin\theta \tan\theta = 3\cos\theta + 2$$

showing each stage of your working.

(5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





Question Number	Scheme	Marks
<p><b>8. (i)</b></p> <p><b>(ii)(a)</b></p> <p><b>(b)</b></p>	<p><math>( \alpha  = 56.3099\dots)</math>  <math>x = \{\alpha + 40 = 96.309993\dots\} = \text{awrt } \mathbf{96.3}</math>  <math>x - 40^\circ = -180 + "56.3099" \dots</math> or <math>x - 40^\circ = -\pi + "0.983" \dots</math>  <math>x = \{-180 + 56.3099\dots + 40 = -83.6901\dots\} = \text{awrt } \mathbf{-83.7}</math></p> <p><math>\sin \theta \left( \frac{\sin \theta}{\cos \theta} \right) = 3 \cos \theta + 2</math>  <math>\left( \frac{1 - \cos^2 \theta}{\cos \theta} \right) = 3 \cos \theta + 2</math>  <math>1 - \cos^2 \theta = 3 \cos^2 \theta + 2 \cos \theta \Rightarrow 0 = 4 \cos^2 \theta + 2 \cos \theta - 1^*</math></p> <p><math>\cos \theta = \frac{-2 \pm \sqrt{4 - 4(4)(-1)}}{8}</math>  <b>or</b> <math>4(\cos \theta \pm \frac{1}{4})^2 \pm q \pm 1 = 0</math>, or <math>(2 \cos \theta \pm \frac{1}{2})^2 \pm q \pm 1 = 0</math>, <math>q \neq 0</math> so <math>\cos \theta = \dots</math>                  One solution is <math>72^\circ</math> or <math>144^\circ</math>, Two solutions are <math>72^\circ</math> and <math>144^\circ</math>  <math>\theta = \{72, 144, 216, 288\}</math></p>	<p>B1 M1 A1 <b>(3)</b></p> <p>M1 dM1 A1 cso * <b>(3)</b></p> <p>M1 A1, A1 M1 A1 <b>(5)</b> <b>[11]</b></p>
<b>Notes for Question 8</b>		
<p><b>(i)</b></p> <p><b>(ii) (a)</b></p> <p><b>(b)</b></p>	<p>B1: 96.3 by any or no method                  M1: Takes 180 degrees from arctan (1.5) or from their "96.3" May be implied by A1. (Could be obtained by adding 180, then subtracting 360).                  A1: awrt -83.7                  Extra answers: ignore extra answers outside range. Any extra answers in range lose final A mark (if earned)                  Working in radians – could earn M1 for <math>x - 40^\circ = -\pi + "0.983" \dots</math> so B0M1A0</p> <p>M1: uses <math>\tan \theta = \frac{\sin \theta}{\cos \theta}</math> or equivalent in equation (not just <math>\tan = \frac{\sin}{\cos}</math>, with no argument)                  dM1: uses <math>\sin^2 \theta = 1 - \cos^2 \theta</math> (quoted correctly) in equation                  A1: completes proof correctly, with no errors to give printed answer*. Need at least three steps in proof and need to achieve the correct quadratic with all terms on one side and "=0"</p> <p>M1: Attempts to solve quadratic by correct quadratic formula, or completion of the square .                  Factorisation attempts score M0.                  1<sup>st</sup> A1: Either 72 <b>or</b> 144, 2<sup>nd</sup> A1: both 72 <b>and</b> 144 (allow 72.0 etc.)                  M1: 360 – "a previous solution" (provided that cos was being used) (not dependent on previous M)                  A1: All four solutions correct (Extra solutions in range lose this A mark, but outside range - ignore)  <b>(Premature approximation:</b> e.g. 71.9, 144.1, 288.1 and 215.9 – lose first A1 then fit other angles)                  Do <b>not</b> require degrees symbol for the marks  <b>Special case: Working in radians</b>                  M1: as before, A1 for either <math>\theta = \frac{2}{5}\pi</math> or <math>\theta = \frac{4}{5}\pi</math> or decimal equivalents, and 2<sup>nd</sup> A1: both                  M1: <math>2\pi - \alpha_1</math> or <math>2\pi - \alpha_2</math> then A0 so 4/5</p>	



Question Number	Scheme	Marks
<p>9. (a)</p> <p>(b)</p>	$\left\{ \frac{dy}{dx} = \right\} 2x - 16x^{-\frac{1}{2}}$ $2x - 16x^{-\frac{1}{2}} = 0 \Rightarrow x^{\frac{3}{2}} = , x^{-\frac{3}{2}} = , \text{ or } 2x - = 16x^{-\frac{1}{2}} \text{ then squared then obtain } x^3 =$ <p>[or <math>2x - 16x^{-\frac{1}{2}} = 0 \Rightarrow x = 4</math> (no wrong work seen)]</p> $(x^{\frac{3}{2}} = 8 \Rightarrow) x = 4$ $x = 4, y = 4^2 - 32\sqrt{4} + 20 = -28 \text{ (ignore } y = 100 \text{ as second answer)}$ $\left\{ \frac{d^2y}{dx^2} = \right\} 2 + 8x^{-\frac{3}{2}}$ $\left( \frac{d^2y}{dx^2} > 0 \Rightarrow \right) y \text{ is a minimum ( there should be no wrong reasoning)}$	<p>M1 A1</p> <p>M1</p> <p>A1</p> <p>M1 A1</p> <p>(6)</p> <p>M1 A1</p> <p>A1</p> <p>(3)</p> <p>[9]</p>
(b)	<p><b>Alternative Method: Gradient Test:</b>  M1 for finding the gradient either side of their <math>x</math>-value from part (a).  A1 for <u>both gradients calculated correctly to 1 significant figure, then using <math>&lt; 0</math> and <math>&gt; 0</math> respectively maybe by use of sketch or table.</u> (See appendix for gradient values. This is <b>not ft their <math>x</math></b>)  A1 states minimum needs M1A1 to have been awarded.</p>	
<b>Notes for Question 9</b>		
(a)	<p>1<sup>st</sup> M1: At least one term differentiated correctly, so <math>x^2 \rightarrow 2x</math>, or <math>32\sqrt{x} \rightarrow 16x^{-\frac{1}{2}}</math>, or <math>20 \rightarrow 0</math></p> <p>A1: This answer or equivalent e.g. <math>2x - \frac{16}{\sqrt{x}}</math></p> <p>2<sup>nd</sup> M1: Sets their <math>\frac{dy}{dx}</math> to 0, and solves to give <math>x^{\frac{3}{2}} = , x^{-\frac{3}{2}} = \text{ or } x^3 =</math> after correct squaring or spots <math>x = 4</math></p> <p>(NB <math>\left\{ \frac{d^2y}{dx^2} = 0 \right\}</math> so <math>2 + 8x^{-\frac{3}{2}} = 0</math> is <b>M0</b> )</p> <p>N.B. Common error: Putting derivative = 0 and merely obtaining <math>x = 0</math> is M0A0, then M0A0 for next two marks. (The first two marks in (a) and marks for second derivative may be earned in part (b).)  A1: <math>x = 4</math> cao [ <math>x = -4</math> is A0 and <math>x = \pm 4</math> is also A0 ]</p> <p>3<sup>rd</sup> M1: Substitutes <b>their positive</b> found <math>x</math> (<b>NOT zero</b>) into <math>y = x^2 - 32\sqrt{x} + 20, x &gt; 0</math>. <b>Should follow attempting to set <math>\frac{dy}{dx} = 0</math> and not setting <math>\frac{d^2y}{dx^2} = 0</math></b></p> <p>A1: -28 cao (Does not need to be written as coordinates)</p>	
(b)	<p>M1: Attempts differentiation of their first derivative with at least one term differentiated correctly. Should be seen or referred to (in part (b)) in determining the nature of the stationary point.</p> <p>A1: Answer in scheme or equivalent</p> <p>A1: States minimum (Second derivative should be correct- can follow incorrect positive <math>x</math>. Needs M1A1 to have been awarded- should not follow incorrect reasoning – (need not say <math>\frac{d^2y}{dx^2} &gt; 0</math> but should not have said <math>\frac{d^2y}{dx^2} = 0</math> for example )</p>	

Leave blank

10.




Figure 4

The circle  $C$  has radius 5 and touches the  $y$ -axis at the point  $(0, 9)$ , as shown in Figure 4.

(a) Write down an equation for the circle  $C$ , that is shown in Figure 4. **(3)**

A line through the point  $P(8, -7)$  is a tangent to the circle  $C$  at the point  $T$ .

(b) Find the length of  $PT$ . **(3)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>10. (a)</b></p> <p><b>(b)</b></p>	<p>Equation of form <math>(x \pm 5)^2 + (y \pm 9)^2 = k</math> , <math>k &gt; 0</math>                      Equation of form <math>(x - a)^2 + (y - b)^2 = 5^2</math> , with values for <math>a</math> and <math>b</math>  <math>(x + 5)^2 + (y - 9)^2 = 25 = 5^2</math>   <math>P(8, -7)</math>. Let centre of circle = <math>X(-5, 9)</math>  <math>PX^2 = (8 - "-5")^2 + (-7 - "9")^2</math> or <math>PX = \sqrt{(8 - "-5")^2 + (-7 - 9)^2}</math>  <math>(PX = \sqrt{425}</math> or <math>5\sqrt{17}</math> ) <math>PT^2 = (PX)^2 - 5^2</math> with numerical <math>PX</math>  <math>PT \{ = \sqrt{400} \} = 20</math> (allow 20.0)</p>	<p>M1 M1 A1 <b>(3)</b>  M1 dM1 A1 cso <b>(3)</b> <b>[6]</b></p>
<p><b>Alternative 2 for (a)</b></p>	<p>Equation of the form <math>x^2 + y^2 \pm 10x \pm 18y + c = 0</math>                      Uses <math>a^2 + b^2 - 5^2 = c</math> with their <math>a</math> and <math>b</math> or substitutes <math>(0, 9)</math> giving <math>+9^2 \pm 2b \times 9 + c = 0</math>  <math>x^2 + y^2 + 10x - 18y + 81 = 0</math></p>	<p>M1 M1 A1 <b>(3)</b></p>
<p><b>Alternative 2 for (b)</b></p>	<p>An attempt to find the point <math>T</math> may result in pages of algebra, but solution needs to reach <math>(-8, 5)</math> or <math>\left(\frac{-8}{17}, 11\frac{2}{17}\right)</math> to get first M1 (even if gradient is found first)                       M1: Use either of the correct points with <math>P(8, -7)</math> and distance between two points formula                      A1: 20</p>	<p>M1  dM1 A1cso <b>(3)</b></p>
<p><b>Alternative 3 for (b)</b></p>	<p>Substitutes <math>(8, -7)</math> into circle <b>equation</b> so <math>PT^2 = 8^2 + (-7)^2 + 10 \times 8 - 18 \times (-7) + 81</math>                      Square roots to give <math>PT \{ = \sqrt{400} \} = 20</math></p>	<p>M1 dM1A1 <b>(3)</b></p>
<b>Notes for Question 10</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p>	<p><b>The three marks in (a) each require a circle equation – (see special cases which are not circles)</b>                      M1: Uses coordinates of centre to obtain LHS of circle equation (RHS must be <math>r^2</math> or <math>k &gt; 0</math> or a positive value)                      M1: Uses <math>r = 5</math> to obtain RHS of circle equation as 25 or <math>5^2</math>                      A1: correct circle equation in any equivalent form  <b>Special cases</b> <math>(x \pm 5)^2 + (x \pm 9)^2 = (5^2)</math> is <b>not a circle</b> equation so M0M0A0                      Also <math>(x \pm 5)^2 + (y - 9) = (5^2)</math> <b>And</b> <math>(x \pm 5)^2 - (y \pm 9)^2 = (5^2)</math> <b>are not circles</b> and gain M0M0A0  <b>But</b> <math>(x - 0)^2 + (y - 9)^2 = 5^2</math> gains M0M1A0</p> <p>M1: Attempts to find distance from their <b>centre of circle</b> to <math>P</math> (or square of this value). If this is called <math>PT</math> and given as answer this is M0. Solution may use letter other than <math>X</math>, as centre was not labelled in the question.                      N.B. Distance from <math>(0, 9)</math> to <math>(8, -7)</math> is incorrect method and is M0, followed by M0A0.                      dM1: Applies the <b>subtraction</b> form of Pythagoras to find <math>PT</math> or <math>PT^2</math> (depends on previous method mark for distance from <b>centre to P</b>) or uses appropriate complete method involving trigonometry                      A1: 20 cso</p>	