

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	4	/	0	1	Signature	

Paper Reference(s)

6664/01

# Edexcel GCE

# Core Mathematics C2

## Advanced Subsidiary

## Wednesday 9 June 2010 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

### Materials required for examination

---

### Mathematical Formulae (Pink)

### Items included with question papers

Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 10 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

## Advice to Candidates

---

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.  
©2010 Edexcel Limited.

Printer's Log. No.

Printer's Log. No.  
H35384A

W850/R6664/57570 4/5



*Turn over*

**edexcel**   
advancing learning, changing lives

Leave  
blank

$$y = 3^x + 2x$$

$x$	0	0.2	0.4	0.6	0.8	1
$y$	1	1.65				5

(2)

value for  $\int_0^1 (3^x + 2x) \, dx$ .

(4)

[illegible]

June 2010  
Core Mathematics C2 6664  
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) 2.35, 3.13, 4.01 (One or two correct B1 B0, all correct B1 B1)  <u>Important:</u> If part (a) is blank, or if answers have been crossed out and no replacement answers are visible, please send to Review as 'out of clip'.</p>	<p>B1 B1 (2)</p>
	<p>(b) <math>\frac{1}{2} \times 0.2 \dots\dots</math> (or equivalent numerical value)  <math>k\{(1+5)+2(1.65+p+q+r)\}</math>, <math>k</math> constant, <math>k \neq 0</math> (See notes below)  <math>= 2.828</math> (awrt 2.83, allowed even after minor slips in values)  The fractional answer <math>\frac{707}{250}</math> (or other fraction wrt 2.83) is also acceptable.  Answers with no working score no marks.</p>	<p>B1 M1 A1 A1 (4) 6</p>
	<p>(a) Answers must be given to 2 decimal places.  <u>No marks</u> for answers given to only 1 decimal place.</p> <p>(b) The <math>p</math>, <math>q</math> and <math>r</math> below are positive numbers, none of which is equal to any of: 1, 5, 1.65, 0.2, 0.4, 0.6 or 0.8  M1 A1: <math>k\{(1+5)+2(1.65+p+q+r)\}</math>  M1 A0: <math>k\{(1+5)+2(1.65+p+q)\}</math> or <math>k\{(1+5)+2(p+q+r)\}</math>  M0 A0: <math>k\{(1+5)+2(1.65+p+q+r+other\ value(s))\}</math>  Note that if the only mistake is to <u>omit</u> a value from the second bracket, this is considered as a slip and the M mark is allowed.  <u>Bracketing mistake:</u> i.e. <math>\frac{1}{2} \times 0.2(1+5)+2(1.65+2.35+3.13+4.01)</math>  instead of <math>\frac{1}{2} \times 0.2\{(1+5)+2(1.65+2.35+3.13+4.01)\}</math>, so that only the <math>(1+5)</math> is multiplied by 0.1 scores B1 M1 A0 A0 <u>unless</u> the final answer implies that the calculation has been done correctly (then full marks can be given).  <u>Alternative:</u>  Separate trapezia may be used, and this can be marked equivalently.</p>	

Leave  
blank

2.

$$f(x) = 3x^3 - 5x^2 - 58x + 40$$

(a) Find the remainder when  $f(x)$  is divided by  $(x-3)$ .

(2)

Given that  $(x - 5)$  is a factor of  $f(x)$ ,

(b) find all the solutions of  $f(x) = 0$ .

(5)



Question Number	Scheme	Marks
2	(a) Attempting to find $f(3)$ or $f(-3)$ $f(3) = 3(3)^3 - 5(3)^2 - (58 \times 3) + 40 = 81 - 45 - 174 + 40 = -98$	M1 A1 (2)
	(b) $\{3x^3 - 5x^2 - 58x + 40 = (x - 5)\} (3x^2 + 10x - 8)$ Attempt to <u>factorise</u> 3-term quadratic, or to use the quadratic formula (see general principles at beginning of scheme). This mark may be implied by the correct solutions to the quadratic. $(3x - 2)(x + 4) = 0 \quad x = \dots \quad \underline{\text{or}} \quad x = \frac{-10 \pm \sqrt{100 + 96}}{6}$ $\frac{2}{3}$ (or exact equiv.), $-4$ , $5$ (Allow 'implicit' solns, e.g. $f(5) = 0$ , etc.) Completely correct solutions without working: full marks.	M1 A1 M1 A1 ft A1 (5) 7
	(a) <u>Alternative (long division):</u> Divide by $(x - 3)$ to get $(3x^2 + ax + b)$ , $a \neq 0$ , $b \neq 0$ . [M1] $(3x^2 + 4x - 46)$ , and $-98$ seen. [A1] (If continues to say 'remainder = 98', isw)	
	(b) 1st M requires use of $(x - 5)$ to obtain $(3x^2 + ax + b)$ , $a \neq 0$ , $b \neq 0$ . (Working need not be seen... this could be done 'by inspection'.) $(3x^2 + 10x - 8) \longleftarrow$ 2 <sup>nd</sup> M for the attempt to <u>factorise</u> their 3-term quadratic, or to solve it using the quadratic formula. Factorisation: $(3x^2 + ax + b) = (3x + c)(x + d)$ , where $ cd  =  b $ . A1ft: Correct factors for their 3-term quadratic <u>followed by a solution</u> (at least one value, which might be incorrect), <u>or</u> numerically correct expression from the quadratic formula for their 3-term quadratic. <u>Note</u> therefore that if the quadratic is correctly factorised but no solutions are given, the last 2 marks will be lost. <u>Alternative (first 2 marks):</u> $(x - 5)(3x^2 + ax + b) = 3x^3 + (a - 15)x^2 + (b - 5a)x - 5b = 0$ , then compare coefficients to find <u>values</u> of $a$ and $b$ . [M1] $a = 10$ , $b = -8$ [A1] <u>Alternative 1:</u> (factor theorem) M1: Finding that $f(-4) = 0$ A1: Stating that $(x + 4)$ is a factor. M1: Finding third factor $(x - 5)(x + 4)(3x \pm 2)$ . A1: Fully correct factors (no ft available here) <u>followed by a solution</u> , (which might be incorrect). A1: All solutions correct. <u>Alternative 2:</u> (direct factorisation) M1: Factors $(x - 5)(3x + p)(x + q)$ A1: $pq = -8$ M1: $(x - 5)(3x \pm 2)(x \pm 4)$ Final A marks as in Alternative 1.	
Throughout this scheme, allow $\left(x \pm \frac{2}{3}\right)$ as an alternative to $(3x \pm 2)$ .		

Leave  
blank

$$y = x^2 - k\sqrt{x}, \text{ where } k \text{ is a constant.}$$

(2)

(2)



Question Number	Scheme	Marks
3	(a) $\left(\frac{dy}{dx} = \right) 2x - \frac{1}{2}kx^{\frac{1}{2}}$ (Having an extra term, e.g. +C, is A0)	M1 A1 (2)
	(b) Substituting $x = 4$ into their $\frac{dy}{dx}$ and ‘compare with zero’ (The mark is allowed for : $<, >, =, \leq, \geq$ ) $8 - \frac{k}{4} < 0 \quad k > 32 \quad (\text{or } 32 < k) \quad \underline{\text{Correct inequality needed}}$	M1 A1 (2) 4
	(a) M: $x^2 \rightarrow cx$ or $k\sqrt{x} \rightarrow cx^{\frac{1}{2}}$ ( $c$ constant, $c \neq 0$ )  (b) Substitution of $x = 4$ into $y$ scores M0. However, $\frac{dy}{dx}$ is sometimes <u>called</u> $y$ , and in this case the M mark can be given.  $\frac{dy}{dx} = 0$ may be ‘implied’ for M1, when, for example, a value of $k$ or an inequality solution for $k$ is found.  <u>Working</u> must be seen to justify marks in (b), i.e. $k > 32$ alone is M0 A0.	

Leave  
blank

- Given that the coefficient of  $x^2$  in this expansion is 525,

- (b) find the possible values of  $a$ . (2)

[illegible]



Question Number	Scheme	Marks
4	<p>(a) <math>(1+ax)^7 = 1 + 7ax \dots</math> or <math>1 + 7(ax) \dots</math> (<u>Not</u> unsimplified versions)</p> <p><math>+ \frac{7 \times 6}{2}(ax)^2 + \frac{7 \times 6 \times 5}{6}(ax)^3</math> Evidence from <u>one</u> of these terms is enough</p> <p><math>+ 21a^2x^2</math> or <math>+ 21(ax)^2</math> or <math>+ 21(a^2x^2)</math></p> <p><math>+ 35a^3x^3</math> or <math>+ 35(ax)^3</math> or <math>+ 35(a^3x^3)</math></p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(4)</p>
	<p>(b) <math>21a^2 = 525</math></p> <p><math>a = \pm 5</math> (Both values are required)</p> <p>(The answer <math>a = 5</math> with no working scores M1 A0)</p>	<p>M1</p> <p>A1</p> <p>(2)</p> <p>6</p>
	<p>(a) The terms can be ‘listed’ rather than added.</p> <p>M1: Requires correct structure: a correct binomial coefficient in any form (perhaps from Pascal’s triangle) with the correct power of <math>x</math>. Allow missing <math>a</math>’s and wrong powers of <math>a</math>, e.g.</p> $\frac{7 \times 6}{2}ax^2, \quad \frac{7 \times 6 \times 5}{3 \times 2}x^3$ <p>However, <math>21 + a^2x^2 + 35 + a^3x^3</math> or similar is M0.</p> <p><math>1 + 7ax + 21 + a^2x^2 + 35 + a^3x^3 = 57 + \dots</math> scores the B1 (isw).</p> <p><math>\binom{7}{2}</math> and <math>\binom{7}{3}</math> or equivalent such as <math>{}^7C_2</math> and <math>{}^7C_3</math> are acceptable,</p> <p>but <u>not</u> <math>\left(\frac{7}{2}\right)</math> or <math>\left(\frac{7}{3}\right)</math> (unless subsequently corrected).</p> <p>1<sup>st</sup> A1: Correct <math>x^2</math> term. 2<sup>nd</sup> A1: Correct <math>x^3</math> term (The binomial coefficients <u>must</u> be simplified).</p> <div style="border: 1px solid black; padding: 5px;"> <p><u>Special case:</u></p> <p>If <math>(ax)^2</math> and <math>(ax)^3</math> are seen within the working, but then lost...</p> <p>... A1 A0 can be given if <math>21ax^2</math> and <math>35ax^3</math> are <u>both</u> achieved.</p> </div> <p><u><math>a</math>’s omitted throughout:</u></p> <p>Note that only the M mark is available in this case.</p> <p>(b) M: Equating their coefficient of <math>x^2</math> to 525.</p> <p>An equation in <math>a</math> or <math>a^2</math> alone is required for this M mark, but allow ‘recovery’ that shows <u>the required coefficient</u>, e.g.</p> <p><math>21a^2x^2 = 525 \Rightarrow 21a^2 = 525</math> is acceptable,</p> <p>but <math>21a^2x^2 = 525 \Rightarrow a^2 = 25</math> is not acceptable.</p> <p>After <math>21ax^2</math> in the answer for (a), allow ‘recovery’ of <math>a^2</math> in (b) so that full marks are available for (b) (but not retrospectively for (a)).</p>	

Leave  
blank

- (1)

- (5)



Question Number	Scheme	Marks
5	(a) $\tan \theta = \frac{2}{5}$ (or 0.4) (i.s.w. if a value of $\theta$ is subsequently found) Requires the correct value with no incorrect working seen.	B1 (1)
	(b) awrt 21.8 ( $\alpha$ ) (Also allow awrt 68.2, ft from $\tan \theta = \frac{5}{2}$ in (a), but no other ft) (This value must be seen in part (b). It may be implied by a correct solution, e.g. 10.9)	B1
	180 + $\alpha$ (= 201.8), or 90 + ( $\alpha/2$ ) (if division by 2 has already occurred) ( $\alpha$ found from $\tan 2x = \dots$ or $\tan x = \dots$ or $\sin 2x = \pm \dots$ or $\cos 2x = \pm \dots$ )	M1
	360 + $\alpha$ (= 381.8), or 180 + ( $\alpha/2$ ) ( $\alpha$ found from $\tan 2x = \dots$ or $\sin 2x = \dots$ or $\cos 2x = \dots$ ) OR 540 + $\alpha$ (= 561.8), or 270 + ( $\alpha/2$ ) ( $\alpha$ found from $\tan 2x = \dots$ )	M1
	Dividing at least one of the angles by 2 ( $\alpha$ found from $\tan 2x = \dots$ or $\sin 2x = \dots$ or $\cos 2x = \dots$ ) $x = 10.9, 100.9, 190.9, 280.9$ (Allow awrt)	M1 A1 (5) 6

(b) Extra solution(s) in range: Loses the final A mark.

Extra solutions outside range: Ignore (whether correct or not).

Common answers:

10.9 and 100.9 would score B1 M1 M0 M1 A0 (Ensure that these M marks are awarded)

10.9 and 190.9 would score B1 M0 M1 M1 A0 (Ensure that these M marks are awarded)

Alternatives:

$$\begin{aligned}
 \text{(i)} \quad 2 \cos 2x - 5 \sin 2x = 0 \quad R \cos(2x + \lambda) = 0 \quad \lambda = 68.2 \Rightarrow 2x + 68.2 = 90 & \quad \text{B1} \\
 2x + \lambda = 270 & \quad \text{M1} \\
 2x + \lambda = 450 \quad \text{or} \quad 2x + \lambda = 630 & \quad \text{M1} \\
 \text{Subtracting } \lambda \text{ and dividing by 2 (at least once)} & \quad \text{M1}
 \end{aligned}$$

$$\text{(ii)} \quad 25 \sin^2 2x = 4 \cos^2 2x = 4(1 - \sin^2 2x)$$

$$29 \sin^2 2x = 4 \quad 2x = 21.8 \quad \text{B1}$$

The M marks are scored as in the main scheme, but extra solutions will be likely, losing the A mark.

Using radians:

B1: Can be given for awrt 0.38 ( $\beta$ )

M1: For  $\pi + \beta$  or  $180 + \beta$

M1: For  $2\pi + \beta$  or  $3\pi + \beta$  (Must now be consistently radians)

M1: For dividing at least one of the angles by 2

A1: For this mark, the answers must be in degrees.

(Correct) answers only (or by graphical methods):

B and M marks can be awarded by implication, e.g.


10.9 scores B1 M0 M0 M1 A0

10.9, 100.9 scores B1 M1 M0 M1 A0

10.9, 100.9, 190.9, 280.9 scores full marks.

Using 11, etc. instead of 10.9 can still score the M marks by implication.

6.



### Figure 1

Figure 1 shows the sector  $OAB$  of a circle with centre  $O$ , radius 9 cm and angle 0.7 radians.

- (a) Find the length of the arc  $AB$ . (2)

- (b) Find the area of the sector  $OAB$ . (2)

The line  $AC$  shown in Figure 1 is perpendicular to  $OA$ , and  $OBC$  is a straight line.

- (c) Find the length of  $AC$ , giving your answer to 2 decimal places. (2)

The region  $H$  is bounded by the arc  $AB$  and the lines  $AC$  and  $CB$ .

- (d) Find the area of  $H$ , giving your answer to 2 decimal places. (3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Question Number	Scheme	Marks
6	(a) $r\theta = 9 \times 0.7 = 6.3$ (Also allow 6.30, or awrt 6.30)	M1 A1 (2)
	(b) $\frac{1}{2}r^2\theta = \frac{1}{2} \times 81 \times 0.7 = 28.35$ (Also allow 28.3 or 28.4, or awrt 28.3 or 28.4) (Condone $28.35^2$ written instead of $28.35 \text{ cm}^2$ )	M1 A1 (2)
	(c) $\tan 0.7 = \frac{AC}{9}$ $AC = 7.58$ (Allow awrt) <u>NOT</u> 7.59 (see below)	M1 A1 (2)
	(d) Area of triangle $AOC = \frac{1}{2}(9 \times \text{their } AC)$ (or other complete method) Area of $R = "34.11" - "28.35"$ (triangle – sector) or (sector – triangle) (needs a <u>value</u> for each) $= 5.76$ (Allow awrt)	M1 M1 A1 (3) 9
	(a) M: Use of $r\theta$ (with $\theta$ in radians), or equivalent (could be working in degrees with a correct degrees formula).  (b) M: Use of $\frac{1}{2}r^2\theta$ (with $\theta$ in radians), or equivalent (could be working in degrees with a correct degrees formula).  (c) M: Other methods must be fully correct, e.g. $\frac{AC}{\sin 0.7} = \frac{9}{\sin\left(\frac{\pi}{2} - 0.7\right)}$ $(\pi - 0.7)$ instead of $\left(\frac{\pi}{2} - 0.7\right)$ here is <u>not</u> a fully correct method.  <u>Premature approximation (e.g. taking angle C as 0.87 radians):</u> This will often result in loss of A marks, e.g. $AC = 7.59$ in (c) is A0.	



Question Number	Scheme	Marks
7	(a) $2\log_3(x-5) = \log_3(x-5)^2$ $\log_3(x-5)^2 - \log_3(2x-13) = \log_3 \frac{(x-5)^2}{2x-13}$ $\log_3 3 = 1$ seen or used correctly $\log_3\left(\frac{P}{Q}\right) = 1 \Rightarrow P = 3Q \quad \left\{ \frac{(x-5)^2}{2x-13} = 3 \Rightarrow (x-5)^2 = 3(2x-13) \right\}$ $x^2 - 16x + 64 = 0$ (*)	B1 M1 B1 M1 A1 cso (5)
	(b) $(x-8)(x-8) = 0 \Rightarrow x = 8$ <u>Must</u> be seen in part (b). Or: Substitute $x = 8$ into original equation and verify. Having additional solution(s) such as $x = -8$ loses the A mark. $x = 8$ with no working scores both marks.	M1 A1 (2) 7

(a) Marks may be awarded if equivalent work is seen in part (b).

1st M:  $\log_3(x-5)^2 - \log_3(2x-13) = \frac{\log_3(x-5)^2}{\log_3(2x-13)}$  is M0

$2\log_3(x-5) - \log_3(2x-13) = 2\log \frac{x-5}{2x-13}$  is M0

2<sup>nd</sup> M: After the first mistake above, this mark is available only if there is 'recovery' to the required

$\log_3\left(\frac{P}{Q}\right) = 1 \Rightarrow P = 3Q$ . Even then the final mark (cso) is lost.

'Cancelling logs', e.g.  $\frac{\log_3(x-5)^2}{\log_3(2x-13)} = \frac{(x-5)^2}{2x-13}$  will also lose the 2<sup>nd</sup> M.

A typical wrong solution:

$\log_3 \frac{(x-5)^2}{2x-13} = 1 \Rightarrow \log_3 \frac{(x-5)^2}{2x-13} = 3 \Rightarrow \frac{(x-5)^2}{2x-13} = 3 \Rightarrow (x-5)^2 = 3(2x-13)$



(Wrong step here)

This, with no evidence elsewhere of  $\log_3 3 = 1$ , scores B1 M1 B0 M0 A0

However,  $\log_3 \frac{(x-5)^2}{2x-13} = 1 \Rightarrow \frac{(x-5)^2}{2x-13} = 3$  is correct and could lead to full marks.


(Here  $\log_3 3 = 1$  is implied).

No log methods shown:

It is not acceptable to jump immediately to  $\frac{(x-5)^2}{2x-13} = 3$ . The only mark this scores is the 1<sup>st</sup> B1 (by generous implication).

(b) M1: Attempt to solve the given quadratic equation (usual rules), so the factors  $(x-8)(x-8)$  with no solution is M0.

**8.**



### Figure 2

Figure 2 shows a sketch of part of the curve  $C$  with equation

$$y = x^3 - 10x^2 + kx,$$

where  $k$  is a constant.

The point  $P$  on  $C$  is the maximum turning point.

Given that the  $x$ -coordinate of  $P$  is 2,

- (a) show that  $k = 28$ .

(3)

The line through  $P$  parallel to the  $x$ -axis cuts the  $y$ -axis at the point  $N$ .

The region  $R$  is bounded by  $C$ , the  $y$ -axis and  $PN$ , as shown shaded in Figure 2.

- (b) Use calculus to find the exact area of  $R$ .

(6)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.



Question Number	Scheme	Marks
8	<p>(a) <math>\frac{dy}{dx} = 3x^2 - 20x + k</math> (Differentiation is required)</p> <p>At <math>x = 2</math>, <math>\frac{dy}{dx} = 0</math>, so <math>12 - 40 + k = 0</math> <math>k = 28</math> (*)</p> <p><u>N.B. The ' = 0 ' must be seen at some stage to score the final mark.</u></p> <p><u>Alternatively:</u> (using <math>k = 28</math>)</p> <p><math>\frac{dy}{dx} = 3x^2 - 20x + 28</math> (M1 A1)</p> <p>'Assuming' <math>k = 28</math> only scores the final cso mark if there is justification that <math>\frac{dy}{dx} = 0</math> at <math>x = 2</math> represents the <u>maximum</u> turning point.</p>	<p>M1 A1</p> <p>A1 cso</p> <p>(3)</p>
	<p>(b) <math>\int (x^3 - 10x^2 + 28x) dx = \frac{x^4}{4} - \frac{10x^3}{3} + \frac{28x^2}{2}</math> Allow <math>\frac{kx^2}{2}</math> for <math>\frac{28x^2}{2}</math></p> <p><math>\left[ \frac{x^4}{4} - \frac{10x^3}{3} + 14x^2 \right]_0^2 = \dots</math> <math>\left( = 4 - \frac{80}{3} + 56 = \frac{100}{3} \right)</math></p> <p>(With limits 0 to 2, substitute the limit 2 into a 'changed function')</p> <p>y-coordinate of <math>P = 8 - 40 + 56 = 24</math> <u>Allow if seen in part (a)</u> (The B1 for 24 may be scored by implication from later working)</p> <p>Area of rectangle = <math>2 \times</math> (their y - coordinate of <math>P</math>)</p> <p>Area of <math>R =</math> (their 48) <math>-</math> <math>\left( \text{their } \frac{100}{3} \right) = \frac{44}{3} \left( 14\frac{2}{3} \text{ or } 14.\dot{6} \right)</math></p> <p>If the subtraction is the 'wrong way round', the final A mark is lost.</p>	<p>M1 A1</p> <p>M1</p> <p>B1</p> <p>M1 A1</p> <p>(6) 9</p>
	<p>(a) M: <math>x^n \rightarrow cx^{n-1}</math> (<math>c</math> constant, <math>c \neq 0</math>) for one term, seen <u>in part (a)</u>.</p> <p>(b) 1<sup>st</sup> M: <math>x^n \rightarrow cx^{n+1}</math> (<math>c</math> constant, <math>c \neq 0</math>) for one term. Integrating the <u>gradient function</u> loses this M mark.</p> <p>2ndM: Requires use of limits 0 and 2, with 2 substituted into a 'changed function'. (It may, for example, have been differentiated).</p> <p>Final M: Subtract their values either way round. This mark is dependent on the use of calculus and a correct method attempt for the area of the rectangle.</p> <p>A1: Must be <u>exact</u>, not 14.67 or similar, but isw after seeing, say, <math>\frac{44}{3}</math>.</p> <p><u>Alternative:</u> (effectively finding area of rectangle by integration)</p> <p><math>\int \{24 - (x^3 - 10x^2 + 28x)\} dx = 24x - \left( \frac{x^4}{4} - \frac{10x^3}{3} + \frac{28x^2}{2} \right)</math>, etc.</p> <p>This can be marked equivalently, with the 1<sup>st</sup> A being for integrating the same 3 terms correctly. The 3<sup>rd</sup> M (for subtraction) will be scored at the same stage as the 2<sup>nd</sup> M. If the subtraction is the 'wrong way round', the final A mark is lost.</p>	



Question Number	Scheme	Marks
9	(a) $25\,000 \times 1.03 = 25750$ $\left\{ 25000 + 750 = 25750, \text{ or } 25000 \frac{(1 - 0.03^2)}{1 - 0.03} = 25750 \right\}$ (*)	B1 (1)
	(b) $r = 1.03$ Allow $\frac{103}{100}$ or $1\frac{3}{100}$ but no other alternatives	B1 (1)
	(c) $25000r^{N-1} > 40\,000$ (Either letter $r$ or their $r$ value) Allow '=' or '<' $r^M > 1.6 \Rightarrow \log r^M > \log 1.6$ Allow '=' or '<' (See below) OR (by change of base), $\log_{1.03} 1.6 < M \Rightarrow \frac{\log 1.6}{\log 1.03} < M$ $(N - 1)\log 1.03 > \log 1.6$ (Correct bracketing required) (*) Accept work for part (c) seen in part (d)	M1 M1 A1 cso (3)
	(d) Attempt to evaluate $\frac{\log 1.6}{\log 1.03} + 1$ {or $25000(1.03)^{15}$ and $25000(1.03)^{16}$ } $N = 17$ (not 16.9 and not e.g. $N \geq 17$ ) Allow '17 <sup>th</sup> year' Accept work for part (d) seen in part (c)	M1 A1 (2)
	(e) Using formula $\frac{a(1 - r^n)}{1 - r}$ with values of $a$ and $r$ , and $n = 9, 10$ or $11$ $\frac{25000(1 - 1.03^{10})}{1 - 1.03}$ 287 000 (must be rounded to the nearest 1 000) Allow 287000.00	M1 A1 A1 (3) 10

(c) 2<sup>nd</sup> M: Requires  $\frac{40000}{25000}$  to be dealt with, and 'two' logs introduced.

With, say,  $N$  instead of  $N - 1$ , this mark is still available.

Jumping straight from  $1.03^{N-1} > 1.6$  to  $(N - 1)\log 1.03 > \log 1.6$  can score only M1 M0 A0.

(The intermediate step  $\log 1.03^{N-1} > \log 1.6$  must be seen).

Longer methods require correct log work throughout for 2<sup>nd</sup> M, e.g.:

$$\log(25000r^{N-1}) > \log 40\,000 \Rightarrow \log 25\,000 + \log r^{N-1} > \log 40\,000 \Rightarrow$$

$$\log r^{N-1} > \log 40\,000 - \log 25\,000 \Rightarrow \log r^{N-1} > \log 1.6$$

(d) Correct answer with no working scores both marks.

Evaluating  $\log\left(\frac{1.6}{1.03}\right) + 1$  does not score the M mark.

(e) M1 can also be scored by a "year by year" method, with terms added.

(Allow the M mark if there is evidence of adding 9, 10 or 11 terms).

1<sup>st</sup> A1 is scored if the 10 correct terms have been added (allow terms to be to the nearest 100).

To the nearest 100, these terms are:

25000, 25800, 26500, 27300, 28100, 29000, 29900, 30700, 31700, 32600

No working shown: Special case: 287 000 scores 1 mark, scored on ePEN as 1, 0, 0.

(Other answers with no working score no marks).

**10.** The circle  $C$  has centre  $A(2, 1)$  and passes through the point  $B(10, 7)$ .

- (a) Find an equation for  $C$ .

(4)

The line  $l_1$  is the tangent to  $C$  at the point  $B$ .

- (b) Find an equation for  $l_1$ .

(4)

The line  $l_2$  is parallel to  $l_1$  and passes through the mid-point of  $AB$ .

Given that  $l_2$  intersects  $C$  at the points  $P$  and  $Q$ ,

- (c) find the length of  $PQ$ , giving your answer in its simplest surd form.

(3)



Question Number	Scheme	Marks
10	<p>(a) <math>(10-2)^2 + (7-1)^2</math> or <math>\sqrt{(10-2)^2 + (7-1)^2}</math>  <math>(x \pm 2)^2 + (y \pm 1)^2 = k</math> (<math>k</math> a positive <u>value</u>)  <math>(x-2)^2 + (y-1)^2 = 100</math> (Accept <math>10^2</math> for 100)                      (Answer only scores full marks)</p>	<p>M1 A1                      M1                      A1                      (4)</p>
	<p>(b) (Gradient of radius) <math>= \frac{7-1}{10-2} = \frac{6}{8}</math> (or equiv.) Must be seen in part (b)                      Gradient of tangent <math>= -\frac{4}{3}</math> (Using perpendicular gradient method)  <math>y-7 = m(x-10)</math> Eqn., in any form, of a line through (10, 7) with any numerical gradient (except 0 or <math>\infty</math>)  <math>y-7 = -\frac{4}{3}(x-10)</math> or equiv (ft gradient of <u>radius</u>, dep. on <u>both</u> M marks)  <math>\{3y = -4x + 61\}</math> (N.B. The A1 is only available as <u>ft</u> after B0)                      The unsimplified version scores the A mark (isw if necessary... subsequent mistakes in simplification are not penalised here.                      The equation must at some stage be <u>exact</u>, not, e.g. <math>y = -1.3x + 20.3</math></p>	<p>B1                      M1                      M1                      A1ft                      (4)</p>
	<p>(c) <math>\sqrt{r^2 - \left(\frac{r}{2}\right)^2}</math> Condone sign slip if there is evidence of correct use of Pythag.  <math>= \sqrt{10^2 - 5^2}</math> or numerically exact equivalent  <math>PQ (= 2\sqrt{75}) = 10\sqrt{3}</math> Simplest surd form <math>10\sqrt{3}</math> required for final mark</p>	<p>M1                      A1                      A1                      (3)                      11</p>
	<p>(b) 2<sup>nd</sup> M: Using (10, 7) to find the equation, in any form, of a straight line through (10, 7), with any numerical gradient (except 0 or <math>\infty</math>).  <u>Alternative</u>: 2<sup>nd</sup> M: Using (10, 7) and an <math>m</math> value in <math>y = mx + c</math> to find a value of <math>c</math>.                      (b) <u>Alternative</u> for first 2 marks (differentiation):  <math>2(x-2) + 2(y-1)\frac{dy}{dx} = 0</math> or equiv. B1                      Substitute <math>x = 10</math> and <math>y = 7</math> to find a value for <math>\frac{dy}{dx}</math> M1                      (This M mark can be awarded generously, even if the attempted 'differentiation' is not 'implicit').                      (c) <u>Alternatives</u>:                      To score M1, must be a <u>fully</u> correct method to obtain <math>\frac{1}{2}PQ</math> or <math>PQ</math>.                      1<sup>st</sup> A1: For alternative methods that find <math>PQ</math> directly, this mark is for an <u>exact numerically correct version</u> of <math>PQ</math>.</p>	