	C	•	u	L	•	U		•	•	•	•	,
								6	6	6	35	5

Write your name here		
Surname	Other na	mes
Pearson	Centre Number	Candidate Number
Edexcel GCE		
Coro Mat	homatic	
Advanced	Incinatic	5 5
Advanced Tuesday 20 June 2017 –	Afternoon	Paper Reference
Advanced Tuesday 20 June 2017 – Time: 1 hour 30 minute	Afternoon es	Paper Reference 6665/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.





Turn over 🕨



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

	Leave
1. Express $\frac{4x}{x^2 - 9} - \frac{2}{x + 3}$ as a single fraction in its simplest form. (4)	

Question Number	Scheme	Marks
1.	$x^2 - 9 = (x + 3)(x - 3)$	B1
	$\frac{4x}{x^2 - 9} - \frac{2}{(x+3)} = \frac{4x - 2(x-3)}{(x+3)(x-3)}$	M1
	$=\frac{2x+6}{(x+3)(x-3)}$	A1
	$=\frac{2(x+3)}{(x+3)(x-3)}$	
	$=\frac{2}{(x-3)}$	A1
		(4)

B1 $x^2 - 9 = (x+3)(x-3)$ This can occur anywhere.

M1 For combining the two fractions with a common denominator. The denominator must be correct and at least one numerator must have been adapted. Accept as separate fractions. Condone missing brackets.

For example accept
$$\frac{4x}{x^2 - 9} - \frac{2}{x + 3} = \frac{4x(x + 3) - 2(x^2 - 9)}{(x + 3)(x^2 - 9)}$$

accept separately
$$\frac{4x}{(x + 3)(x - 3)} - \frac{2}{(x + 3)} = \frac{4x}{(x + 3)(x - 3)} - \frac{2x - 3}{(x + 3)(x - 3)}$$
 condoning missing bracket
condone
$$\frac{4x}{x^2 - 9} - \frac{2}{x + 3} = \frac{4x(x + 3) - 2}{(x + 3)(x^2 - 9)}$$
.....as only one numerator has been adapted
A correct intermediate form of $\frac{\text{simplified linear}}{\text{simplified quadratic}}$

Accept
$$\frac{2x+6}{(x+3)(x-3)}$$
, $\frac{2x+6}{x^2-9}$, and even $\frac{(2x+6)(x+3)}{(x^2-9)(x+3)}$,

A1 Further factorises and cancels (which may be implied) to reach the answer $\frac{2}{x-3}$

Do not penalise correct solutions that include incomplete lines Eg $\frac{4x-2(x-3)}{(x+3)(x-3)} = \frac{4x-2x+6}{...} = \frac{2x+6}{(x+3)(x-3)} = \frac{2}{x-3}$

This is not a "show that" question.

A1

Note: Watch out for an answer of $\frac{2}{x+3}$ probably scored from $\frac{4x-2(x-3)}{(x+3)(x-3)} = \frac{2x-6}{(x+3)(x-3)} = \frac{2(x-3)}{(x+3)(x-3)}$

This would score B1 M1 A0 A0

Paper	This resource was created and owned by Pearson Edexcel	Mathoma	66
			Leav
			blan
2. Find the	e exact solutions, in their simplest form, to the equations		
(a) e^{3x}	-9 = 8		
(a) C	- 0	(3)	
		(0)	
(b) ln(2	$2y + 5) = 2 + \ln(4 - y)$		
		(4)	

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Qu Nı	uestion umber	Scheme	Marks
2	2.(a)	$e^{3x-9} = 8 \Longrightarrow 3x-9 = \ln 8$	M1
		$\Rightarrow x = \frac{\ln 8 + 9}{3}, = \ln 2 + 3$	A1, A1
			(3)
	(b)	$\ln(2y+5) = 2 + \ln(4-y)$ (2y+5)	
		$\ln\left(\frac{2y+3}{4-y}\right) = 2$	M1
		$\left(\frac{2y+5}{4-y}\right) = e^2$	M1
		$2y+5 = e^2(4-y) \Longrightarrow 2y + e^2y = 4e^2 - 5 \Longrightarrow y = \frac{4e^2 - 5}{2+e^2}$	dM1, A1
			(4)
(n)			7 marks
A1 Alt I (a) $e^{3x-9} =$ Alt II (a) $e^{x-3} = \frac{1}{2}$	$cso \ln 2$ $8 \Rightarrow \frac{e^{3x}}{e^{9}}$ $a)$ $\sqrt[3]{8} \Rightarrow x$	2+3. Accept $\ln 2e^3$ $= 8 \Rightarrow e^{3x} = 8e^9 \Rightarrow 3x = \ln(8e^9)$ for M1 (Condone slips on index work and lack of $-3 = \ln(\sqrt[3]{8})$ for M1 (Condone slips on the 9. Eg $e^{x-9} = 2 \Rightarrow x-9 = \ln 2$)	bracket)
(b) M1	Uses a d	correct method to combine two terms to create a single ln term.	
	Eg. Sco	ore for $2 + \ln(4 - y) = \ln(e^2(4 - y))$ or $\ln(2y + 5) - \ln(4 - y) = \ln\left(\frac{2y + 5}{4 - y}\right)$	
M1	Condon Scored the ln te	he slips on the signs and coefficients of the terms, but not on the e^2 for an attempt to undo the ln's to get an equation in y This must be awarded after an a terms. Award for $\ln(g(y)) = 2 \Longrightarrow g(y) = e^2$ and can be scored eg where $g(y) = 2y$	attempt to combine $+5 - (4 - y)$
lM1	It canno Depend factoris	be awarded for just $2y+5 = e^2 + 4 - y$ where the candidate attempts to undo term lent upon both previous M's. It is for making y the subject. Expect to see both terms is ded (may be implied) before reaching y =. Condone slips, for eg, on signs. $y = 2.615$	by term by collected and scores this.
A 1	$y = \frac{4\epsilon}{2}$	$e^2 - 5$ or equivalent such as $y = 4 - \frac{13}{2 + e^2}$ ISW after you see the correct answ	ver.
Special	Case: 1	$\ln(2y+5) - \ln(4-y) = 2 \Longrightarrow \frac{\ln(2y+5)}{\ln(4-y)} = 2 \Longrightarrow \frac{2y+5}{4-y} = e^2 \Longrightarrow \text{Correct answer score}$	e M0 M1 M1 A0



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



6665

Question Number	Scheme	Marks
3. (a)	y3	B1
		(1)
(b)	$y=3+\sqrt{x+2} \Rightarrow y-3=\sqrt{x+2} \Rightarrow x=(y-3)^2-2$	M1 A1
	$\Rightarrow g^{-1}(x) = (x-3)^2 - 2, \text{ with } x3$	A1
		(3)
(c)	$g(x) = x \Longrightarrow 3 + \sqrt{x+2} = x$	
	$\Rightarrow x+2=(x-3)^2 \Rightarrow x^2-7x+7=0$	M1, A1
	$\Rightarrow x = \frac{7 \pm \sqrt{21}}{2} \Rightarrow x = \frac{7 + \sqrt{21}}{2} \text{ only}$	M1, A1
		(4)
(d)	$a = \frac{7 + \sqrt{21}}{2}$	B1 ft
		(1)
		9 marks
(c) Alt	Solves $g^{-1}(x) = x \Longrightarrow (x-3)^2 - 2 = x$	
	$\Rightarrow x^2 - 7x + 7 = 0$	M1, A1
	$\Rightarrow x = \frac{7 \pm \sqrt{21}}{2} \Rightarrow x = \frac{7 + \sqrt{21}}{2} \text{ only}$	dM1, A1
		(4)

(a)

- B1 States the correct range for g Accept g(x) ... 33g... 3, Range... $3, [3, \infty)$ Range is greater than or equal to 3 Condone f ... 3 Do not accept $g(x) > 3, x ... 3, (3, \infty)$
- (b)

A1 Achieves $x = (y-3)^2 - 2$ or if swapped $y = (x-3)^2 - 2$ or equivalent such as $x = y^2 - 6y + 7$

A1 Requires a correct function in x + correct domain or a correct function in x with a correct follow through on the range in (a) but do not follow through on $x \in \mathbb{R}$

M1 Attempts to make x or a swapped y the subject of the formula. The minimum expectation is that the 3 is moved over followed by an attempt to square both sides. Condone for this mark $\sqrt{x+2} = y \pm 3 \Longrightarrow x+2 = y^2 \pm 9$

(c)

M1 Sets $3+\sqrt{x+2} = x$, moves the 3 over and then attempts to square both sides. Can be scored for $\sqrt{x+2} = x-3 \Rightarrow x+2 = x^2 \pm 9$

A1 $x^2 - 7x + 7 = 0$. The = 0 may be implied by subsequent working

M1 Correct method of solving their 3TQ by the formula/ completing the square. The equation must have real roots. It is dependent upon them having attempted to set $3 + \sqrt{x+2} = x$ and proceeding to a quadratic. You may just see both roots written down which is fine.

Allow for this mark decimal answers Eg 5.79 and 1.21 for $x^2 - 7x + 7 = 0$ You may need to check with a calc.

A1 $(x) = \frac{7 + \sqrt{21}}{2}$ or exact equivalent **only**.

This answer following the correct quadratic would imply the previous M

Allow
$$x = \frac{7}{2} + \sqrt{\frac{21}{4}}$$
 but **DO NOT** allow $x = \frac{7 \pm \sqrt{21}}{2}$

.

(c) can of course be attempted by solving

$$+\sqrt{x+2} = "(x-3)^{2} - 2" \Longrightarrow x^{4} - 12x^{3} + 44x^{2} - 49x + 14 = 0$$

$$\vdots \qquad \qquad \Rightarrow (x^{2} - 7x + 7)(x^{2} - 5x + 2) = 0$$

The scheme can be applied to this

(d)

B1ft $(a) = \frac{7 + \sqrt{21}}{2}$ or a. You may condone $x = \frac{7 + \sqrt{21}}{2}$. You may allow this following a re - start.

You may allow the correct decimal answer, awrt 5.79, following exact/decimal work in part (c) or a restart. Follow through on their root, including decimals, coming from the **positive** root with the **positive** sign in (c). Eg In (c) . $x^2 - 7x + 11 = 0 \Rightarrow x = \frac{7 \pm \sqrt{5}}{2}$ So the correct follow through would be $x = \frac{7 + \sqrt{5}}{2}$

If they only had one root in (c) then follow through on this as long as it is positive.

SC. If they give the correct roots in parts (c) and (d) without considering the correct answer then award B1 in (d) following the A0 in (c). So $(x) = \frac{7 \pm \sqrt{21}}{2}$ as their answer in part (c), allow $(x/a) = \frac{7 \pm \sqrt{21}}{2}$ for B1 in (d).

6665 Leave

blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

www.mystudybro.com This resource was created and owned by Pearson Edexcel (a) Write $5\cos\theta - 2\sin\theta$ in the form $R\cos(\theta + \alpha)$, where R and α are constants, 4. R > 0 and $0 \leq \alpha < \frac{\pi}{2}$ Give the exact value of R and give the value of α in radians to 3 decimal places. (3) (b) Show that the equation $5\cot 2x - 3\csc 2x = 2$ can be rewritten in the form $5\cos 2x - 2\sin 2x = c$ where c is a positive constant to be determined. (2) (c) Hence or otherwise, solve, for $0 \le x < \pi$, $5 \cot 2x - 3 \csc 2x = 2$ giving your answers to 2 decimal places. (Solutions based entirely on graphical or numerical methods are not acceptable.) (4)





Question Number	Scheme	Marks	
4. (a)	$R = \sqrt{29}$	B1	
	$\tan \alpha = \frac{2}{5} \Longrightarrow \alpha = \text{awrt } 0.381$	M1A1	
	2 2 2		(3)
(b)	$5\cot 2x - 3\csc 2x = 2 \Longrightarrow 5\frac{\cos 2x}{\sin 2x} - \frac{3}{\sin 2x} = 2$	M1	
	$\Rightarrow 5\cos 2x - 2\sin 2x = 3$	A1	(2)
(c)	$5\cos 2x - 2\sin 2x = 3 \Longrightarrow \cos(2x + 0.381) = \frac{3}{\sqrt{29}}$	M1	(-)
	$2x + 0.381 = \arccos\left(\frac{3}{\sqrt{29}}\right) \Longrightarrow x = \dots$	dM1	
	x = awrt 0.30, 2.46	A1A1	
			(4)
		(9 marks)	
Alt I (c)	$5\cos 2x - 2\sin 2x = 3 \Longrightarrow 10\cos^2 x - 5 - 4\sin x \cos x = 3$		
	$\Rightarrow 4 \tan^2 x + 2 \tan x - 1 = 0$	M1	
	$\Rightarrow \tan x = \frac{-1 \pm \sqrt{5}}{4} \Rightarrow x =$	dM1	
	x = awrt 0.30, 2.46	A1A1	(4)
Alt II (c)	$5\cos 2x - 2\sin 2x = 3 \Longrightarrow (5\cos 2x)^2 = (3 + 2\sin 2x)^2 \& \cos^2 2x = 1 - \sin^2 2x$		
	$\Rightarrow 29\sin^2 2x + 12\sin 2x - 16 = 0$	M1	
	$\Rightarrow \sin 2x = \frac{-12 \pm \sqrt{2000}}{58} \Rightarrow 2x = \Rightarrow x =$	dM1	
	x = awrt 0.30, 2.46	A1A1	
			(4)

(a)

B1 $R = \sqrt{29}$

Condone $R = \pm \sqrt{29}$ (Do not allow decimals for this mark Eg 5.39 but remember to isw after $\sqrt{29}$) M1 $\tan \alpha = \pm \frac{2}{5}$, $\tan \alpha = \pm \frac{5}{2} \Longrightarrow \alpha = ...$

If *R* is used to find α accept $\sin \alpha = \pm \frac{2}{R}$ or $\cos \alpha = \pm \frac{5}{R} \Longrightarrow \alpha = ...$

A1 $\alpha = awrt \ 0.381$ Note that the degree equivalent $\alpha = awrt \ 21.8^{\circ}$ is A0

- (b)
- M1 Replaces $\cot 2x$ by $\frac{\cos 2x}{\sin 2x}$ and $\csc 2x$ by $\frac{1}{\sin 2x}$ in the lhs Do not be concerned by the coefficients 5 and -3. Replacing $\cot 2x$ by $\frac{1}{\tan 2x}$ does not score marks until the $\tan 2x$ has been replaced by $\frac{\sin 2x}{\cos 2x}$ They may state $\times \sin 2x \Rightarrow 5 \cos 2x - 3 = 2 \sin 2x$ which implies this mark
- A1 cso $5\cos 2x 2\sin 2x = 3$ There is no need to state the value of 'c' The notation must be correct. They cannot mix variables within their equation

Do not accept for the final A1 $\tan 2x = \frac{\sin 2x}{\cos 2x}$ within their equations

(c)

M1 Attempts to use part (a) and (b). They must be using their *R* and
$$\alpha$$
 from part (a) and their *c* from part (b)
Accept $\cos(2x\pm'\alpha') = \frac{c'}{R'}$ Condone $\cos(\theta\pm'\alpha') = \frac{c'}{R'}$ or $\exp\cos(x\pm'\alpha') = \frac{c'}{R'}$ for the first M

dM1 Score for dealing with the cos, the α and the 2 **correctly** and in that order to reach x = ...Don't be concerned if they change the variable in the question and solve for $\theta =$ (as long as all operations have been undone). You may not see any working. It is implied by one correct answer. You may need to check with a calculator.

Eg for an incorrect $\alpha \cos(2x+1.19) = \frac{3}{\sqrt{29}} \Rightarrow x = -0.105$ would score M1 dM1 A0 A0

- A1 One solution correct, usually x = 0.3/0.30 or x = 2.46 or in degrees 17.2° or $141.(0)^{\circ}$
- A1 Both solutions correct awrt x = awrt 0.30, 2.46 and no extra values in the range. Condone candidates who write 0.3 and 2.46 without any (more accurate) answers In degrees accept awrt 1 dp 17.2°, 141.(0)° and no extra values in the range.

Special case: For candidates who are misreading the question and using their part (a) with 2 on the rhs. They will be allowed to score a maximum of SC M1 dM1 A0 A0

M1 Attempts to use part (a) with 2. They must be using their R and α from part (a)

Accept
$$\cos(2x \pm \alpha') = \frac{2}{R'}$$
 Condone $\cos(\theta \pm \alpha') = \frac{2}{R'}$ or $\exp\cos(x \pm \alpha') = \frac{2}{R'}$ for the first M

dM1 Score for dealing with the cos, the α and the 2 **correctly** and in that order to reach x = ...You may not see any working. It is implied by one correct answer. You may need to check with a calculator.

Eg for an correct
$$\alpha$$
 and $R \cos(2x+0.381) = \frac{2}{\sqrt{29}} \Rightarrow x = 0.405$

Alt to part (c)

M1 Attempts both double angle formulae condoning sign slips on $\cos 2x$, divides by $\cos^2 x$

and forms a quadratic in tan by using the identity $\pm 1 \pm \tan^2 x = \sec^2 x$

- dM1 Attempts to solve their quadratic in tanx leading to a solution for x.
- A1 A1 As above

.....

6665

blank





Figure 2 shows a sketch of part of the curve C with equation

$$y = 2\ln(2x+5) - \frac{3x}{2}, \qquad x > -2.5$$

The point *P* with *x* coordinate -2 lies on *C*.

(a) Find an equation of the normal to C at P. Write your answer in the form ax + by = c, where a, b and c are integers.

(5)

The normal to C at P cuts the curve again at the point Q, as shown in Figure 2.

(b) Show that the x coordinate of Q is a solution of the equation

$$x = \frac{20}{11}\ln(2x+5) - 2 \tag{3}$$

The iteration formula

$$x_{n+1} = \frac{20}{11}\ln(2x_n + 5) - 2$$

can be used to find an approximation for the x coordinate of Q.

(c) Taking $x_1 = 2$, find the values of x_2 and x_3 , giving each answer to 4 decimal places.

(2)





Question Number	Scheme	Marks
5. (a)	At P $x = -2 \Longrightarrow y = 3$	B1
	$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{4}{2x+5} - \frac{3}{2}$	M1, A1
	$\left. \frac{\mathrm{d}y}{\mathrm{d}x} \right _{x=-2} = \frac{5}{2} \Rightarrow \text{ Equation of normal is } y'' = -\frac{2}{5} \left(x - (-2) \right)$	M1
	$\Rightarrow 2x + 5y = 11$	A1
		(5)
(b)	Combines $5y+2x=11$ and $y=2\ln(2x+5)-\frac{3x}{2}$ to form equation in x	
	$5\left(2\ln(2x+5)-\frac{3x}{2}\right)+2x=11$	M1
	$\Rightarrow x = \frac{20}{11}\ln(2x+5) - 2$	dM1 A1*
		(3)
(c)	Substitutes $x_1 = 2 \Longrightarrow x_2 = \frac{20}{11} \ln 9 - 2$	M1
	Awrt $x_2 = 1.9950$ and $x_3 = 1.9929$.	A1
		(2)
		(10 marks)

(a)

B1 y = 3 at point *P*. This may be seen embedded within their equation which may be a tangent

M1 Differentiates
$$\ln(2x+5) \rightarrow \frac{A}{2x+5}$$
 or equivalent. You may see $\ln(2x+5)^2 \rightarrow \frac{A(2x+5)}{(2x+5)^2}$

- A1 $\frac{dy}{dx} = \frac{4}{2x+5} \frac{3}{2}$ oe. It need not be simplified.
- M1 For using a correct method of finding the equation of the normal using their numerical value of $-\frac{dx}{dy}\Big|_{x=-2}$ as

the gradient. Allow for $(y-3') = -\frac{dx}{dy}\Big|_{x=-2}(x--2)$, oe.

At least one bracket must be correct for their (-2,3)

If the form y = mx + c is used it is scored for proceeding as far as c = ...

A1 $\pm k(5y+2x=11)$ It must be in the form ax+by=c as stated in the question

Score this mark once it is seen. Do not withhold it if they proceed to another form, y = mx + c for example If a candidate uses a graphical calculator to find the gradient they can score a maximum of B1 M0 A0 M1 A1

M1 For combining 'their' **linear** 5y + 2x = 11 with $y = 2\ln(2x+5) - \frac{3x}{2}$ to form equation in just *x*,

condoning slips on the rearrangement of their 5y + 2x = 11. Eg $2\ln(2x+5) - \frac{3x}{2} = \frac{11\pm 2x}{5}$ is OK

- dM1 Collects the two terms in x and proceeds to $ax = b \ln(2x+5) + c$ Allow numerical slips
- A1* This is a given answer. All aspects must be correct including bracketing
- (c)
- M1 Score for substituting $x_1 = 2 \Rightarrow x_2 = \frac{20}{11} \ln(2 \times 2 + 5) 2$ or exact equivalent This may implied by $x_2 = 1.99...$
- A1 Both values correct. Allow awrt $x_2 = 1.9950$ and $x_3 = 1.9929$ but condone $x_2 = 1.995$ Ignore subscripts. Mark on the first and second values given.

6665 Leave

blank

DO NOT WRITE IN THIS AREA

6. Given that *a* and *b* are positive constants,

- (a) on separate diagrams, sketch the graph with equation
 - (i) y = |2x a|
 - (ii) y = |2x a| + b

Show, on each sketch, the coordinates of each point at which the graph crosses or meets the axes.

(4)

Given that the equation

$$\left|2x-a\right|+b=\frac{3}{2}x+8$$

has a solution at x = 0 and a solution at x = c,

(b) find c in terms of a.

(4)



Question Number	Scheme	Marks	
6.(a)(i)	(0,a) V shape on x - axis or coordinates $\left(\frac{1}{2}a,0\right)$ and (0,a) $O\left(\frac{1}{2}a,0\right)$ x Correct shape, position and coordinates	B1 B1	
(ii)	(0, $a+b$) (0, $a+b$) Their "V" shape translated up or (0, $a+b$) Correct shape, position and (0, $a+b$)	B1ft B1	
	\rightarrow \rightarrow γ		(4)
(b)	States or uses $a + b = 8$ Attempts to solve $ 2x - a + b = \frac{3}{2}x + 8$ in either x or with $x = c$	B1	
	$2c-a+b=\frac{3}{2}c+8 \Longrightarrow kc=f(a,b)$	M1	
	Combines $kc = f(a,b)$ with $a+b=8 \implies c=4a$	dM1 A1 (8 marks)	(4)

(a)(i)

- B1 V shape sitting anywhere on the *x* axis or for $(\frac{1}{2}a, 0)$ and (0, a) lying on the curve. Condone non -symmetrical graphs and ones lying on just one side of the *y* -axis
- B1 V shape sitting on the positive x-axis at $(\frac{1}{2}a, 0)$, cutting the y-axis at (0, a) and lying in both quadrants 1 and 2 Accept $\frac{1}{2}a$ and a marked on the correct axis. Condone say (a, 0) for (0, a) as long as it is on the correct axis. Condone a dotted line appearing on the diagram as many reflect y = 2x - a to sketch y = |2x - a|If it is a solid line then it would not score the shape mark.

(a)(ii)

- B1ft Follow through on (a)(i). Their graph translated up. Allow on U shapes and non symmetrical graphs. Alternatively score for the (0, a+b) lying on the curve
- B1 V shape lying in quadrants 1 and 2 with the vertex in quadrant 1 cutting the *y* axis at (0, a+b)Ignore any coordinates given for the vertex.

(b)

A1

- B1 States or uses a+b=8 or exact equivalent. Condone use of capital letters throughout It is not scored for just |0-a|+b=8
- M1 This M is for an understanding of the modulus.

It is scored for an attempt at solving $(2x-a)+b = \frac{3}{2}x+8$ or $-(2x-a)+b = \frac{3}{2}x+8$ in either x or with x replaced by c. The signs of the 2x and the a must be different. $|2x-a| \neq 2x+a$

You may see
$$(2x-a)+b = \frac{3}{2}x+8 \Longrightarrow kx = f(a,b)$$

You may see $-2x + a + b = \frac{3}{2}x + 8 \Longrightarrow kx = f(a,b)$

You may see $(2x-a)+b = \frac{3}{2}x+8 \Rightarrow kx = f(a,b)$ being solved with *b* replaced with **their** a+b=8You may see $-2c+a+b = \frac{3}{2}c+8 \Rightarrow kc = f(a,b)$ being solved with *b* replaced with **their** a+b=8

dM1 This dM mark is scored for combining b = 8 - a with $(2x - a) + b = \frac{3}{2}x + 8$ (or their kx = f(a, b) resulting from that equation) resulting in a link between x and a Both equations must have been correct initially. Alternatively for combining b = 8 - a with their $2c - a + b = \frac{3}{2}c + 8$ (or their kc = f(a, b) resulting from that equation) resulting in a link between c and a

You may condone sign slips in finding the link between x (or c) and aIf you see an approach that involves making |2x-a| the subject followed by squaring, and you feel that it deserves credit, please send to review. The solution proceeds as follows

Look for
$$|2x-a| = \frac{3}{2}x+8-b \Rightarrow |2x-a| = \frac{3}{2}x+a \Rightarrow (2x-a)^2 = \left(\frac{3}{2}x+a\right)^2 \Rightarrow 7x\left(\frac{1}{4}x-a\right) = 0$$

 $c = 4a$ ONLY

Special Case where they have the roots linked with the incorrect branch of the curve.

They have x = 0 as the solution to $2x - a + b = \frac{3}{2}x + 8 \Rightarrow -a + b = 8$(1) They have x = c as the solution to $-2x + a + b = \frac{3}{2}x + 8 \Rightarrow \frac{7}{2}x = a + b - 8$(2) Solve (1) and (2) $\Rightarrow x = \frac{4}{7}a$ Hence $\Rightarrow c = \frac{4}{7}a$ This would score P0 M1 dM0 A0 approach be sourced SC P0 M1 dM1. A0

This would score B0 M1 dM0 A0 anyway but should be awarded SC B0, M1 dM1, A0 for above work leading to either $x = \frac{4}{7}a$ or $c = \frac{4}{7}a$

Immer 2017 st Paper	www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics	0 666
		Le	eave ank
7. (i) Gi	ven $y = 2x(x^2 - 1)^5$, show that		
(a)	$\frac{dy}{dx} = g(x)(x^2 - 1)^4$ where $g(x)$ is a function to be determined.		
	dx	(4)	
(b)	Hence find the set of values of x for which $\frac{dy}{dx} \ge 0$		
	ux	(2)	
(ii) Gi	ven		
	$x = \ln(\sec 2y), \qquad 0 < y < \frac{\pi}{4}$		
fin	$d \frac{dy}{dx}$ as a function of x in its simplest form		
1111	$\frac{d}{dx}$ as a function of x in its simplest form.	(4)	

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6665

Question Number	Scheme	Marks	
7(i) (a)	$y = 2x(x^{2}-1)^{5} \Longrightarrow \frac{dy}{dx} = (x^{2}-1)^{5} \times 2 + 2x \times 10x(x^{2}-1)^{4}$	M1A1	
	$\Rightarrow \frac{dy}{dx} = (x^2 - 1)^4 (2x^2 - 2 + 20x^2) = (x^2 - 1)^4 (22x^2 - 2)$	M1 A1	
			(4)
(b)	$\frac{\mathrm{d}y}{\mathrm{d}x} \dots 0 \Longrightarrow (22x^2 - 2) \dots 0 \Longrightarrow \text{ critical values of } \pm \frac{1}{\sqrt{11}}$	M1	
	$x \dots \frac{1}{\sqrt{11}} x_{,,} = -\frac{1}{\sqrt{11}}$	A1	
			(2)
(ii)	$x = \ln(\sec 2y) \Rightarrow \frac{dx}{dy} = \frac{1}{\sec 2y} \times 2\sec 2y \tan 2y$	B1	
	$\Rightarrow \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{2\tan 2y} = \frac{1}{2\sqrt{\sec^2 2y - 1}} = \frac{1}{2\sqrt{\mathrm{e}^{2x} - 1}}$	M1 M1 A1	
			(4)
		10 ma	arks
Alt 1 (ii)	$x = \ln(\sec 2y) \Longrightarrow \sec 2y = e^x$		
	$\Rightarrow 2\sec 2y\tan 2y\frac{\mathrm{d}y}{\mathrm{d}x} = \mathrm{e}^x$	B1	
	$\Rightarrow \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{\mathrm{e}^x}{2\sec 2y\tan 2y} = \frac{\mathrm{e}^x}{2\mathrm{e}^x\sqrt{\sec^2 2y - 1}} = \frac{1}{2\sqrt{\mathrm{e}^{2x} - 1}}$	M1M1A1	
			(4)
Alt 2 (ii)	$y = \frac{1}{2} \arccos\left(e^{-x}\right) \Longrightarrow \frac{dy}{dx} = -\frac{1}{2} \times \frac{1}{\sqrt{1 - \left(e^{-x}\right)^2}} \times -e^{-x}$	B1M1M1	
	$\Rightarrow \frac{\mathrm{d}y}{\mathrm{d}x} = \frac{1}{2\sqrt{\mathrm{e}^{2x} - 1}}$	A1	
			(4)
1		1	

M1 Attempts the product rule to differentiate $2x(x^2-1)^5$ to a form $A(x^2-1)^5 + Bx^n(x^2-1)^4$ where n = 1 or 2. and A, B > 0 If the rule is stated it must be correct, and not with a "-" sign.

Summer 2017 Past Paper (Mark Scheme)

A1 Any unsimplified but correct form
$$\left(\frac{dy}{dx}\right) = 2(x^2 - 1)^3 + 20x^2(x^2 - 1)^4$$

M1 For taking a common factor of $(x^2 - 1)^4$ out of a suitable expression
Look for $A(x^2 - 1)^3 \pm Bx^n(x^2 - 1)^4 = (x^2 - 1)^4 \left\{A(x^2 - 1) \pm Bx^n\right\}$ but you may condone missing brackets
It can be scored from a $u^2 \cdot u^n$ or similar.
A1 $\left(\frac{dy}{dx}\right) = (x^2 - 1)^4 (22x^2 - 2)$ Expect $g(x)$ to be simplified but accept $\frac{dy}{dx} = (x^2 - 1)^4 2(11x^2 - 1)$
There is no need to state $g(x)$ and remember to isw after a correct answer. This must be in part (a).
(i)(b)
M1 Sets their $\frac{dy}{dx} = 0$ or $\frac{dy}{dx} = 0$ and proceeds to find one of the critical values for **their** $g(x)$ or their
 $\frac{dy}{dx} = 0$ rearranged and $+(x^2 - 1)^4$ if $g(x)$ not found. $g(x)$ should be at least a 2TQ with real roots. If $g(x)$ is
factorised, the usual rules apply. The M cannot be awarded from work just on $(x^2 - 1)^4$. 0 is $x = \pm 1$
You may see and accept decimals for the M.
A1 cao $x \dots \frac{1}{\sqrt{11}} x_n - \frac{1}{\sqrt{11}}$ or exact equivalent only. Condone $x \dots \frac{1}{\sqrt{11}} x_n - \frac{1}{\sqrt{11}} \bigcup \left[\frac{\sqrt{11}}{11}, \infty\right]\right)$
Condone the word "and" appearing between the two sets of values.
Withhold the final mark if $x \dots \frac{1}{\sqrt{11}} x_n - \frac{1}{\sqrt{11}}$, appears with values not in this region $g(x_n), 1, x_m - 1$
(ii)
B1 Differentiates and achieves a correct line involving $\frac{dy}{dx}$ or $\frac{dx}{dy}$
Accept $\frac{dx}{dx} = \frac{1}{\sec 2y} \times 2\sec 2y \tan 2y, \frac{dx}{dy} = -\frac{1}{\cos 2y} \times -2\sin 2y - 2\sec 2y \tan 2y \frac{dy}{dx} = e^x$
M1 For inverting their expression for $\frac{dx}{dy}$ to achieve an expression for $\frac{dy}{dx}$.
The variables (on the rhs) must be consistent, you may condone slips on the coefficients but not the terms.
In the alternative mothod it is for correctly changing the subject
M1 Scored for using $\tan^2 2y = \pm 1 \pm \sec^2 2y$ and $\sec 2y = e^x$ to achieve $\frac{dy}{dx}$ or $\frac{dx}{dy}$ in terms of x .
Alternatively they could use $\sin^2 2y + \cos^2 2y = 1$ with $\cos 2y = e^{-x}$ to achieve $\frac{dy}{dx}$ or $\frac{dx}{dy}$ in terms of x .
For the M mark you may

Allow a misread on $x = \ln(\sec y)$ for the two method marks only



The number of rabbits on an island is modelled by the equation

$$P = \frac{100e^{-0.1t}}{1 + 3e^{-0.9t}} + 40, \qquad t \in \mathbb{R}, t \ge 0$$

where P is the number of rabbits, t years after they were introduced onto the island.

A sketch of the graph of P against t is shown in Figure 3.

(a) Calculate the number of rabbits that were introduced onto the island.

(b) Find
$$\frac{\mathrm{d}P}{\mathrm{d}t}$$
 (3)

The number of rabbits initially increases, reaching a maximum value P_T when t = T

(c) Using your answer from part (b), calculate

- (i) the value of T to 2 decimal places,
- (ii) the value of P_T to the nearest integer.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(4)

(1)

For t > T, the number of rabbits decreases, as shown in Figure 3, but never falls below k, where k is a positive constant.

(d) Use the model to state the maximum value of k.

Question	Scheme	Marks	
Number			
8 (a)	$P_0 = \frac{100}{1+3} + 40 = 65$	B1	
	1 - 5		(1)
	$\frac{\mathrm{d}}{\mathrm{d}t}\mathrm{e}^{kt} = C\mathrm{e}^{kt}$	M1	
(b)	$\frac{\mathrm{d}P}{\mathrm{d}t} = \frac{\left(1+3\mathrm{e}^{-0.9t}\right) \times -10\mathrm{e}^{-0.1t} - 100\mathrm{e}^{-0.1t} \times -2.7\mathrm{e}^{-0.9t}}{\left(1+3\mathrm{e}^{-0.9t}\right)^2}$	M1 A1	
			(3)
(c)(i)	At maximum $-10e^{-0.1t} - 30e^{-0.1t} \times e^{-0.9t} + 270e^{-0.1t} \times e^{-0.9t} = 0$		
	$e^{-0.1t} \left(-10 + 240 e^{-0.9t} \right) = 0$		
	$e^{-0.9t} = \frac{10}{240}$ oe $e^{0.9t} = 24$	M1	
	$-0.9t = \ln\left(\frac{1}{2}\right) \Rightarrow t = \frac{10}{2}\ln(24) = 3.53$	M1. A1	
	(24) 9 $($,	
(c) (ii)	Sub $t = 3.53 \Longrightarrow P_T = 102$	A1	
			(4)
(d)	40	B1	
			(1)
		9 marks	

(a)

B1 $(P_0 =)65$

(b)

M1 For sight of $\frac{d}{dt}e^{kt} = Ce^{kt}$ (Allow C =1)This may be within an incorrect product or quotient rule

M1 Scored for a full application of the quotient rule. If the formula is quoted it should be correct.

The denominator should be present even when the correct formula has been quoted.

In cases where a formula has not been quoted it is very difficult to judge that a correct formula has been used (due to the signs between the terms). So.....

if the formula has not been quoted look for the **order** of the terms

$$\frac{(1+3e^{-0.9t}) \times pe^{-0.1t} - qe^{-0.1t} \times e^{-0.9t}}{(1+3e^{-0.9t})^2}$$
$$\frac{(1+3e^{-0.9t}) \times pe^{-0.1t} + qe^{-0.1t} \times e^{-0.9t}}{(1+3e^{-0.9t})^2}$$

For the product rule. Look for $ae^{-0.1t}(1+3e^{-0.9t})^{-1} \pm be^{-0.1t}e^{-0.9t}(1+3e^{-0.9t})^{-2}$ either way around

Penalise if an incorrect formula is quoted . Condone missing brackets in both cases.

A1 A correct **unsimplified** answer.

Eg using quotient rule
$$\left(\frac{dP}{dt}\right) = \frac{-10e^{-0.1t}\left(1+3e^{-0.9t}\right)+270e^{-0.1t}e^{-0.9t}}{\left(1+3e^{-0.9t}\right)^2}$$
 oe $\frac{-10e^{-0.1t}+240e^{-1t}}{\left(1+3e^{-0.9t}\right)^2}$ simplified
Eg using product rule $\left(\frac{dP}{dt}\right) = -10e^{-0.1t}\left(1+3e^{-0.9t}\right)^{-1}+270e^{-0.1t}e^{-0.9t}\left(1+3e^{-0.9t}\right)^{-2}$ oe
Remember to isw after a correct (unsimplified) answer.
There is no need to have the $\frac{dP}{dt}$ and it could be called $\frac{dy}{dx}$

(c)(i) Do NOT allow any marks in here without sight/implication of $\frac{dP}{dt} = 0$, $\frac{dP}{dt} < 0$ OR $\frac{dP}{dt} > 0$

The question requires the candidate to find *t* using part (b) so it is possible to do this part using inequalities using the same criteria as we apply for the equality. All marks in (c) can be scored from an incorrect denominator (most likely *v*), no denominator, or using a numerator the wrong way around ie uv'-u'v

- M1 Sets their $\frac{dP}{dt} = 0$ or the numerator of their $\frac{dP}{dt} = 0$, factorises out or cancels a term in $e^{-0.1t}$ to reach a form $Ae^{\pm 0.9t} = B$ oe. Alternatively they could combine terms to reach $Ae^{-t} = Be^{-0.1t}$ or equivalent Condone a double error on $e^{-0.1t} \times e^{-0.9t} = e^{-0.1t \times -0.9t}$ or similar before factorising. Look for correct indices. If they use the product rule then expect to see their $\frac{dP}{dt} = 0$ followed by multiplication of $(1 + 3e^{-0.9t})^2$ before similar work to the quotient rule leads to a form $Ae^{\pm 0.9t} = B$ M1 Having set the numerator of their $\frac{dP}{dt} = 0$ and obtained either $e^{\pm kt} = C$ (k may be incorrect) or $Ae^{-t} = Be^{-0.1t}$ it is awarded for the correct order of operations, taking ln's leading to t = ..It cannot be awarded from impossible equations Eg $e^{\pm 0.9t} = -0.3$
- A1 cso t = awrt 3.53 Accept $t = \frac{10}{9} \ln(24)$ or exact equivalent.
- (c)(ii)

A1 awrt 102 following 3.53 The M's must have been awarded. This is not a B mark.

- (d)
- B1 Sight of 40
 - Condone statements such as $P \rightarrow 40 \ k... 40$ or likewise

ummer 201 ' st Paper	7 www.mystudybro.com This resource was created and owned by Pearson Edexcel	Mathematics C
יי מאבו	This resource was created and owned by rearson Edekter	Leave
9. (a) P	Prove that	
	$\sin 2x - \tan x \equiv \tan x \cos 2x, \qquad x \neq (2n+1)90^\circ, \qquad n \in \mathbb{Z}$	
		(4)
(b) C	Given that $x \neq 90^{\circ}$ and $x \neq 270^{\circ}$, solve, for $0 \leq x < 360^{\circ}$,	
	$\sin 2x - \tan x = 3\tan x \sin x$	
C	Give your answers in degrees to one decimal place where appropriate.	
(Solut	tions based entirely on graphical or numerical methods are not acceptable	.)
		(5)
30		1
50		

Question Number	Scheme	Marks
9(a)	$\sin 2x - \tan x = 2\sin x \cos x - \tan x$	M1
	$=\frac{2\sin x\cos^2 x}{\cos x}-\frac{\sin x}{\cos x}$	M1
	$=\frac{\sin x}{\cos x} \times (2\cos^2 x - 1)$	
	$= \tan x \cos 2x$	dM1 A1*
		(4)
(b)	$\tan x \cos 2x = 3\tan x \sin x \Longrightarrow \tan x (\cos 2x - 3\sin x) = 0$	
	$\cos 2x - 3\sin x = 0$	M1
	$\Rightarrow 1 - 2\sin^2 x - 3\sin x = 0$	M1
	$\Rightarrow 2\sin^2 x + 3\sin x - 1 = 0 \Rightarrow \sin x = \frac{-3 \pm \sqrt{17}}{4} \Rightarrow x = \dots$	M1
	Two of $x = 16.3^{\circ}, 163.7^{\circ}, 0, 180^{\circ}$	A1
	All four of $x = 16.3^{\circ}, 163.7^{\circ}, 0, 180^{\circ}$	A1
		(5)
		(9 marks)

(a)

M1 Uses a correct double angle identity involving $\sin 2x$ Accept $\sin(x+x) = \sin x \cos x + \cos x \sin x$

M1 Uses $\tan x = \frac{\sin x}{\cos x}$ with $\sin 2x = 2\sin x \cos x$ and attempts to combine the two terms using a common denominator. This can be awarded on two separate terms with a common denominator. Alternatively uses $\sin x = \tan x \cos x$ and attempts to combine two terms using factorisation of $\tan x$ dM1 Both M's must have been scored. Uses a correct double angle identity involving $\cos 2x$.

A1* A fully correct solution with no errors or omissions. All notation must be correct and variables must be consistent

Withhold this mark if for instance they write $\tan x = \frac{\sin x}{\cos x}$

If the candidate $\times \cos x$ on line 1 and/or $\div \sin x$ they cannot score any more than one mark unless they are working with both sides of the equation or it is fully explained.

(b)

M1 The tan x must be cancelled or factorised out to produce $\cos 2x - 3\sin x = 0$ or $\frac{\cos 2x}{\sin x} = 3$ oe Condone slips

M1 Uses $\cos 2x = 1 - 2\sin^2 x$ to form a 3TQ=0 in $\sin x$ The = 0 may be implied by later work

M1 Uses the formula/completion of square or GC with invsin to produce at least one value for x It may be implied by one correct value.

This mark **can** be scored from factorisation of their 3TQ in sin *x* **but only if** their quadratic factorises.

- A1 Two of $x = 0,180^\circ$, awrt 16.3°, awrt 163.7° or in radians two of awrt 0.28, 2.86, 0 and π or 3.14 This mark can be awarded as a SC for those students who just produce $0,180^\circ$ (or 0 and π) from tan x = 0 or $\sin x = 0$.
- A1 All four values in degrees $x = 0,180^{\circ}$, awrt 16.3°, awrt 163.7° and no extra's inside the range 0, $x < 360^{\circ}$. Condone 0 = 0.0 and $180^{\circ} = 180.0^{\circ}$ Ignore any roots outside range.

Alternatives to parts (a) and (b)

(a) Alt 1	$\tan x \cos 2x = \tan x \left(2\cos^2 x - 1 \right)$	M1	
	$= 2\tan x \cos^2 x - \tan x$		
	$=2\frac{\sin x}{\cos x}\cos^2 x - \tan x$	M1	
	$=2\sin x\cos x - \tan x$		
	$=\sin 2x - \tan x$	dM1 A1	
			(4)

a) Alt 1 Starting from the rhs

M1 Uses a correct double angle identity for $\cos 2x$. Accept any correct version including $\cos(x+x) = \cos x \cos x - \sin x \sin x$

M1 Uses
$$\tan x = \frac{\sin x}{\cos x}$$
 with $\cos 2x = 2\cos^2 x - 1$ and attempts to multiply out the bracket

- dM1 Both M's must have been scored. It is for using $2\sin x \cos x = \sin 2x$
- A1* A fully correct solution with no errors or omissions. All notation must be correct and variables must be consistent. See Main scheme for how to deal with candidates who $\div \tan x$

(a) Alt 2	$\sin 2x - \tan x \equiv \tan x \cos 2x$	
	$2\sin x \cos x - \tan x \equiv \tan x (2\cos^2 x - 1)$	M1
	$2\sin x \cos x - \tan x \equiv 2\tan x \cos^2 x - \tan x$	
	$2\sin x \cos x \equiv 2\frac{\sin x}{\cos x} \cos^2 x$	M1
	$2\sin x \cos x \equiv 2\sin x \cos x$	dM1
	+statement that it must be true	A1*

a) Alt 2 Candidates who use both sides

M1 Uses a correct double angle identity involving $\sin 2x$ or $\cos 2x$. Can be scored from either side Accept $\sin(x+x) = \sin x \cos x + \cos x \sin x$ or $\cos(x+x) = \cos x \cos x - \sin x \sin x$

M1 Uses $\tan x = \frac{\sin x}{\cos x}$ with $\cos 2x = 2\cos^2 x - 1$ and cancels the $\tan x$ term from both sides

dM1 Uses a correct double angle identity involving $\sin 2x$ Both previous M's must have been scored

A1* A fully correct solution with no errors or omissions AND statement "hence true", "a tick", "QED". W⁵ All notation must be correct and variables must be consistent

.....

It is possible to solve part (b) without using the given identity. There are various ways of doing this, one of which is shown below.

$$\sin 2x - \tan x = 3 \tan x \sin x \Longrightarrow 2 \sin x \cos x - \frac{\sin x}{\cos x} = 3 \frac{\sin x}{\cos x} \sin x$$

$$2 \sin x \cos^2 x - \sin x = 3 \sin^2 x \qquad \text{M1 Equation in } \sin x \text{ and } \cos x$$

$$2 \sin x (1 - \sin^2 x) - \sin x = 3 \sin^2 x \qquad \text{M1 Equation in } \sin x \text{ only}$$

$$(2 \sin^2 x + 3 \sin x - 1) \sin x = 0$$

$$x = \dots \qquad \text{M1 Solving equation to find at least one } x$$

$$Two \text{ of } x = 16.3^\circ, 163.7^\circ, 0, 180^\circ \qquad \text{A1}$$

$$All \text{ four of } x = 16.3^\circ, 163.7^\circ, 0, 180^\circ \text{ and no extras A1}$$