

Question Number	Scheme		Marks
<p>1. (a)</p> <p>(b)</p>	$\left\{ \frac{1}{\sqrt{(9-10x)}} \right\} (9-10x)^{-\frac{1}{2}}$ $= (9)^{-\frac{1}{2}} \left(1 - \frac{10x}{9}\right)^{-\frac{1}{2}} = \frac{1}{3} \left(1 - \frac{10x}{9}\right)^{-\frac{1}{2}}$ $= \left\{ \frac{1}{3} \right\} \left[1 + \left(-\frac{1}{2}\right)(kx) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} (kx)^2 + \dots \right]$ $= \left\{ \frac{1}{3} \right\} \left[1 + \left(-\frac{1}{2}\right) \left(\frac{-10x}{9}\right) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} \left(\frac{-10x}{9}\right)^2 + \dots \right]$ $= \frac{1}{3} \left[1 + \frac{5}{9}x + \frac{25}{54}x^2 + \dots \right]$ $= \frac{1}{3} + \frac{5}{27}x + \frac{25}{162}x^2 + \dots$ $\frac{3+x}{\sqrt{(9-10x)}} = (3+x)(9-10x)^{-\frac{1}{2}}$ $= (3+x) \left(\frac{1}{3} + \frac{5}{27}x + \left\{ \frac{25}{162}x^2 + \dots \right\} \right)$ $= 1 + \frac{5}{9}x + \frac{25}{54}x^2 + \frac{1}{3}x + \frac{5}{27}x^2 + \dots$ $= 1 + \frac{8}{9}x + \frac{35}{54}x^2 + \dots$	<p>$(9-10x)^{-\frac{1}{2}}$ or uses power of $-\frac{1}{2}$</p> <p>$(9)^{-\frac{1}{2}}$ or $\frac{1}{3}$</p> <p>At least two correct terms. See notes</p> <p><i>Can be implied by later work</i> See notes</p> <p>Multiplies out to give exactly one constant term, exactly 2 terms in x and exactly 2 terms in x^2. Ignore terms in x^3. Can be implied.</p>	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1; A1</p> <p>[5]</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p> <p>8</p>
Question 1 Notes			
(a)	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>Writes down $(9-10x)^{-\frac{1}{2}}$ or uses power of $-\frac{1}{2}$.</p> <p>This mark can be implied by a constant term of $(9)^{-\frac{1}{2}}$ or $\frac{1}{3}$.</p> <p>$(9)^{-\frac{1}{2}}$ or $\frac{1}{3}$ outside brackets or $\frac{1}{3}$ as candidate's constant term in their binomial expansion.</p> <p>Expands $(\dots + kx)^{-\frac{1}{2}}$ to give any 2 terms out of 3 terms simplified or an un-simplified, $1 + (-\frac{1}{2})(kx)$ or $(-\frac{1}{2})(kx) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} (kx)^2$ or $1 + \dots + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!} (kx)^2$, where $k \neq 1$.</p> <p>$\frac{1}{3} + \frac{5}{27}x$ (simplified fractions)</p> <p>Accept only $\frac{25}{162}x^2$</p>	

1. (a) ctd	Note	<p>You cannot recover correct work for part (a) in part (b). i.e. if the correct answer to (a) appears as part of their solution in part (b), it cannot be credited in part (a).</p>
	SC	<p>If a candidate <i>would otherwise score</i> A0A0 then allow Special Case 1st A1 for either</p>
		<p>SC: $\frac{1}{3}\left[1 + \frac{5}{9}x; \dots\right]$ or SC: $\lambda\left[1 + \frac{5}{9}x + \frac{25}{54}x^2 + \dots\right]$ or SC: $\left[\lambda + \frac{5\lambda}{9}x + \frac{25\lambda}{54}x^2 + \dots\right]$</p>
		<p>(where λ can be 1 or omitted), with each term in the [.....] is a simplified fraction</p>
	SC	<p>Special case for the M1 mark</p>
		<p>Award Special Case M1 for a correct simplified or un-simplified $1 + n(kx) + \frac{n(n-1)}{2!}(kx)^2$</p>
		<p>expansion with a value of $n \neq -\frac{1}{2}$, $n \neq$ positive integer and a consistent (kx). Note that (kx)</p>
		<p>must be consistent (on the RHS, not necessarily the LHS) in a candidate's expansion.</p>
		<p>Note that $k \neq 1$.</p>
	Note	<p>Candidates who write $\left\{\frac{1}{3}\right\}\left[1 + \left(-\frac{1}{2}\right)\left(\frac{10x}{9}\right) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!}\left(\frac{10x}{9}\right)^2 + \dots\right]$</p>
		<p>where $k = \frac{10}{9}$ and not $-\frac{10}{9}$ and achieve $\frac{1}{3} - \frac{5}{27}x; + \frac{25}{162}x^2 + \dots$ will get B1B1M1A0A1.</p>
(b)	M1	<p>Writes down $(3 + x)$(their part (a) answer, at least 2 of the 3 terms.)</p>
	Note	<p>$(3 + x)\left(\frac{1}{4} + \frac{5}{4}x + \dots\right)$ or $(3 + x)\left(\frac{1}{3} + \frac{5}{27}x + \frac{25}{162}x^2 + \dots\right)$ are fine for M1.</p>
	Note	<p>This mark can also be implied by candidate multiplying out to find two terms (or coefficients) in x.</p>
	M1	<p>Multiplies out to give exactly one constant term, exactly 2 terms in x and exactly 2 terms in x^2.</p>
	Note	<p>This M1 mark can be implied. You can also ignore x^3 terms.</p>
	A1	<p>$1 + \frac{8}{9}x + \frac{35}{54}x^2 + \dots$</p>
<p>Alternative Methods for part (a)</p>		
<p>Alternative method 1: Candidates can apply an alternative form of the binomial expansion.</p>		
$\left\{\frac{1}{\sqrt{(9-10x)}}\right\} = (9-10x)^{-\frac{1}{2}} = (9)^{-\frac{1}{2}} + \left(-\frac{1}{2}\right)(9)^{-\frac{3}{2}}(-10x) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2!}(9)^{-\frac{5}{2}}(-10x)^2$		
	B1	<p>Writes down $(9-10x)^{-\frac{1}{2}}$ or uses power of $-\frac{1}{2}$.</p>
	B1	<p>$9^{-\frac{1}{2}}$ or $\frac{1}{3}$</p>
	M1	<p>Any two of three (un-simplified or simplified) terms correct.</p>
	A1	<p>$\frac{1}{3} + \frac{5}{27}x$</p>
	A1	<p>$\frac{25}{162}x^2$</p>
	Note	<p>The terms in C need to be evaluated, so $^{-\frac{1}{2}}C_0(9)^{-\frac{1}{2}} + ^{-\frac{1}{2}}C_1(9)^{-\frac{3}{2}}(-10x) + ^{-\frac{1}{2}}C_2(9)^{-\frac{5}{2}}(-10x)^2$ without further working is B1B0M0A0A0.</p>

<p>1. (a)</p>	<p><u>Alternative Method 2: Maclaurin Expansion</u></p> <p>Let $f(x) = \frac{1}{\sqrt{(9-10x)}}$</p> <p>$\{f(x) = (9-10x)^{-\frac{1}{2}}$</p> <p>$f''(x) = 75(9-10x)^{-\frac{5}{2}}$</p> <p>$f'(x) = (-\frac{1}{2})(9-10x)^{-\frac{3}{2}}(-10)$</p> <p>$\left\{ \therefore f(0) = \frac{1}{3}, f'(0) = \frac{5}{27} \text{ and } f''(0) = \frac{75}{243} = \frac{25}{81} \right\}$</p> <p>$f(x) = \frac{1}{3} + \frac{5}{27}x + \frac{25}{162}x^2 + \dots$</p>	<p>$(9-10x)^{-\frac{1}{2}}$ B1</p> <p>Correct $f''(x)$ B1 oe</p> <p>$\pm a(9-10x)^{-\frac{3}{2}}; a \neq \pm 1$ M1</p> <p>A1; A1</p>
---------------	---	---

2.




Figure 1

Figure 1 shows a sketch of part of the curve with equation

$$y = (2 - x)e^{2x}, \quad x \in \mathbb{R}$$

The finite region R , shown shaded in Figure 1, is bounded by the curve, the x -axis and the y -axis.

The table below shows corresponding values of x and y for $y = (2 - x)e^{2x}$

x	0	0.5	1	1.5	2
y	2	4.077	7.389	10.043	0

- (a) Use the trapezium rule with all the values of y in the table, to obtain an approximation for the area of R , giving your answer to 2 decimal places. (3)
- (b) Explain how the trapezium rule can be used to give a more accurate approximation for the area of R . (1)
- (c) Use calculus, showing each step in your working, to obtain an exact value for the area of R . Give your answer in its simplest form. (5)



Question Number	Scheme		Marks
2. (a)	$\text{Area} \approx \frac{1}{2} \times 0.5 \times [2 + 2(4.077 + 7.389 + 10.043) + 0]$ $= \frac{1}{4} \times 45.018 = 11.2545 = 11.25 \text{ (2 dp)}$		B1; M1 11.25 A1 cao [3]
(b)	Any one of <ul style="list-style-type: none"> • Increase the number of strips • Use more trapezia • Make h smaller • Increase the number of x and/or y values used • Shorter /smaller intervals for x • More values of y. • More intervals of x • Increase n 		B1 [1]
(c)	$\left\{ \int (2-x)e^{2x} dx \right\}, \left\{ \begin{array}{l} u = 2-x \Rightarrow \frac{du}{dx} = -1 \\ \frac{dv}{dx} = e^{2x} \Rightarrow v = \frac{1}{2}e^{2x} \end{array} \right\}$		
$= \frac{1}{2}(2-x)e^{2x} - \int -\frac{1}{2}e^{2x} \{dx\}$			Either $(2-x)e^{2x} \rightarrow \pm \lambda(2-x)e^{2x} \pm \int \mu e^{2x} \{dx\}$ or $\pm x e^{2x} \rightarrow \pm \lambda x e^{2x} \pm \int \mu e^{2x} \{dx\}$ M1
$= \frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x}$			$(2-x)e^{2x} \rightarrow \frac{1}{2}(2-x)e^{2x} - \int -\frac{1}{2}e^{2x} \{dx\}$ A1
$\text{Area} = \left\{ \left[\frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x} \right]_0^2 \right\}$			$\frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x}$ A1 oe
$= \left(0 + \frac{1}{4}e^4 \right) - \left(\frac{1}{2}(2)e^0 + \frac{1}{4}e^0 \right)$			Applies limits of 2 and 0 <i>to all terms</i> and subtracts the correct way round. dM1
$= \frac{1}{4}e^4 - \frac{5}{4}$			$\frac{1}{4}e^4 - \frac{5}{4} \text{ or } \frac{e^4 - 5}{4} \text{ cao}$ A1 oe
Question 2 Notes			
(a)	B1 M1 Note A1 Note Note	Outside brackets $\frac{1}{2} \times 0.5$ or $\frac{0.5}{2}$ or 0.25 or $\frac{1}{4}$. For structure of trapezium rule [.....]. Condone missing 0. No errors are allowed [eg. an omission of a y-ordinate or an extra y-ordinate or a repeated y ordinate]. 11.25 cao Working must be seen to demonstrate the use of the trapezium rule. The actual area is 12.39953751... Award B1M1A1 for $\frac{0.5}{2}(2+0) + \frac{1}{2}(4.077 + 7.389 + 10.043) = 11.25$	[5] 9

2. (a) contd	<p>Bracketing mistake: Unless the final answer implies that the calculation has been done correctly. Award B1M0A0 for $\frac{1}{2} \times 0.5 + 2 + 2(4.077 + 7.389 + 10.043) + 0$ (nb: answer of 45.268).</p>
	<p>Alternative method for part (a): Adding individual trapezia</p> $\text{Area} \approx 0.5 \times \left[\frac{2+4.077}{2} + \frac{4.077+7.389}{2} + \frac{7.389+10.043}{2} + \frac{10.043+0}{2} \right] = 11.2545 = 11.25 \text{ (2 dp) cao}$ <p>B1 0.5 and a divisor of 2 on all terms inside brackets. M1 First and last ordinates once and the middle ordinates twice inside brackets ignoring the 2. A1 11.25 cao</p>
(b)	<p>B0 Give B0 for</p> <ul style="list-style-type: none"> • smaller values of x and/or y. • use more decimal places
(c)	<p>M1 Either $(2-x)e^{2x} \rightarrow \pm \lambda(2-x)e^{2x} \pm \int \mu e^{2x} \{dx\}$ or $\pm xe^{2x} \rightarrow \pm \lambda xe^{2x} \pm \int \mu e^{2x} \{dx\}$</p> <p>A1 $(2-x)e^{2x} \rightarrow \frac{1}{2}(2-x)e^{2x} - \int -\frac{1}{2}e^{2x} \{dx\}$ either un-simplified or simplified.</p> <p>A1 Correct expression, i.e. $\frac{1}{2}(2-x)e^{2x} + \frac{1}{4}e^{2x}$ or $\frac{5}{4}e^{2x} - xe^{2x}$ (or equivalent)</p> <p>dM1 which is dependent on the 1st M1 mark being awarded. Complete method of applying limits of 2 and 0 to all terms and subtracting the correct way round.</p> <p>Note Evidence of a proper consideration of the limit of 0 is needed for M1. So, just subtracting zero is M0.</p> <p>A1 $\frac{1}{4}e^4 - \frac{5}{4}$ or $\frac{e^4 - 5}{4}$. Do not allow $\frac{1}{4}e^4 - \frac{5}{4}e^0$ unless simplified to give $\frac{1}{4}e^4 - \frac{5}{4}$</p> <p>Note 12.39953751... without seeing $\frac{1}{4}e^4 - \frac{5}{4}$ is A0.</p> <p>Note 12.39953751... from NO working is M0A0A0M0A0.</p>

Question Number	Scheme	Marks
<p>3.</p> <p>(a)</p> <p>(b)</p>	$x^2 + y^2 + 10x + 2y - 4xy = 10$ $\left\{ \begin{array}{l} \cancel{dx} \\ \cancel{dx} \end{array} \right\} \times \frac{2x + 2y \frac{dy}{dx} + 10 + 2 \frac{dy}{dx} - \left(4y + 4x \frac{dy}{dx} \right)}{dx} = \frac{0}{dx}$ $2x + 10 - 4y + (2y + 2 - 4x) \frac{dy}{dx} = 0$ $\frac{dy}{dx} = \frac{2x + 10 - 4y}{4x - 2y - 2}$ <p>Simplifying gives $\frac{dy}{dx} = \frac{x + 5 - 2y}{2x - y - 1} \left\{ = \frac{-x - 5 + 2y}{-2x + y + 1} \right\}$</p> $\left\{ \frac{dy}{dx} = 0 \Rightarrow \right\} x + 5 - 2y = 0$ <p>So $x = 2y - 5$,</p> $(2y - 5)^2 + y^2 + 10(2y - 5) + 2y - 4(2y - 5)y = 10$ $4y^2 - 20y + 25 + y^2 + 20y - 50 + 2y - 8y^2 + 20y = 10$ <p>gives $-3y^2 + 22y - 35 = 0$ or $3y^2 - 22y + 35 = 0$</p> $(3y - 7)(y - 5) = 0 \text{ and } y = \dots$ $y = \frac{7}{3}, 5$	<p>See notes</p> <p>M1 A1 M1</p> <p>Dependent on the first M1 mark.</p> <p>dM1</p> <p>A1 cso oe</p> <p>[5]</p> <p>M1</p> <p>M1</p> <p>A1 oe</p> <p>see notes</p> <p>Method mark for solving a quadratic equation.</p> <p>ddM1</p> <p>A1 cao</p> <p>[5]</p>
<p>(b)</p>	<p>Alternative method for part (b)</p> $\left\{ \frac{dy}{dx} = 0 \Rightarrow \right\} x + 5 - 2y = 0$ <p>So $y = \frac{x + 5}{2}$,</p> $x^2 + \left(\frac{x + 5}{2} \right)^2 + 10x + 2 \left(\frac{x + 5}{2} \right) - 4x \left(\frac{x + 5}{2} \right) = 10$ $x^2 + \frac{x^2 + 10x + 25}{4} + 10x + x + 5 - 2x^2 - 10x = 10$ $4x^2 + x^2 + 10x + 25 + 40x + 4x + 20 - 8x^2 - 40x = 40$ <p>gives $-3x^2 + 14x + 5 = 0$ or $3x^2 - 14x - 5 = 0$</p> $(3x + 1)(x - 5) = 0, x = \dots$ $y = \frac{-\frac{1}{3} + 5}{2}, \frac{5 + 5}{2}$ $y = \frac{7}{3}, 5$	<p>M1</p> <p>M1</p> <p>A1 oe</p> <p>see notes</p> <p>Solves a quadratic and finds at least one value for y.</p> <p>ddM1</p> <p>A1 cao</p> <p>[5]</p>
		10

Question 3 Notes			
3. (a)	M1	Differentiates implicitly to include either $\pm 4x \frac{dy}{dx}$ or $y^2 \rightarrow 2y \frac{dy}{dx}$ or $2y \rightarrow 2 \frac{dy}{dx}$. (Ignore $\left(\frac{dy}{dx} = \right)$).	
	A1	$x^2 + y^2 + 10x + 2y \rightarrow 2x + 2y \frac{dy}{dx} + 10 + 2 \frac{dy}{dx}$ and $10 \rightarrow 0$	
	M1	$-4xy \rightarrow \pm 4y \pm 4x \frac{dy}{dx}$	
	Note	If an extra term appears then award 1 st A0.	
	Note	$2x + 2y \frac{dy}{dx} + 10 + 2 \frac{dy}{dx} - 4y - 4x \frac{dy}{dx} \rightarrow 2x + 10 - 4y = -2y \frac{dy}{dx} - 2 \frac{dy}{dx} + 4x \frac{dy}{dx}$ will get 1 st A1 (implied) as the "= 0" can be implied by rearrangement of their equation.	
	dM1	dependent on the first method mark being awarded. An attempt to factorise out all the terms in $\frac{dy}{dx}$ as long as there are at least two terms in $\frac{dy}{dx}$.	
	A1	$\frac{x + 5 - 2y}{2x - y - 1}$ or $\frac{-x - 5 + 2y}{-2x + y + 1}$ (must be simplified).	
	cs0:	If the candidate's solution is not completely correct, then do not give this mark.	
	(b)	M1	Sets the numerator of their $\frac{dy}{dx}$ equal to zero (or the denominator of their $\frac{dx}{dy}$ equal to zero) oe.
		NOTE	If the numerator involves one variable only then <i>only</i> the 1st M1 mark is possible in part (b).
M1		Substitutes their x or their y into the printed equation to give an equation in one variable only.	
A1		For obtaining either $-3y^2 + 22y - 35 \{= 0\}$ or $3y^2 - 22y + 35 \{= 0\}$	
Note		This mark can also awarded for a correct three term equation, eg. either $-3y^2 + 22y = 35$ $3y^2 - 22y = -35$ or $3y^2 + 35 = 22y$ are all fine for A1.	
ddM1		Dependent on the previous 2 M marks. See notes at the beginning of the mark scheme: Method mark for solving a 3 term quadratic	
		<ul style="list-style-type: none"> • $(3y - 7)(y - 5) = 0 \Rightarrow y = \dots$ • $y = \frac{22 \pm \sqrt{(-22)^2 - 4(3)(35)}}{2(3)}$ • $y^2 - \frac{22}{3}y - \frac{35}{3} = 0 \Rightarrow \left(y - \frac{11}{3}\right)^2 - \frac{121}{9} + \frac{35}{3} = 0 \Rightarrow y = \dots$ • Or writes down at least one correct y-root from their quadratic equation. This is usually found from their calculator. 	
Note		If a candidate applies the alternative method then they also need to use their $y = \frac{x + 5}{2}$ in order to find at least one value for y in order to gain the final M1.	
A1		$y = \frac{7}{3}, 5$. cao. (2.33 or 2.3 without reference to $\frac{7}{3}$ or $2\frac{1}{3}$ is not allowed for this mark.)	
Note		It is possible for a candidate who does not achieve full marks in part (a), (but has a correct numerator for $\frac{dy}{dx}$) to gain all 5 marks in part (b).	

Question Number	Scheme	Marks
4. (a)	$\frac{25}{x^2(2x+1)} \equiv \frac{A}{x} + \frac{B}{x^2} + \frac{C}{(2x+1)}$ $B = 25, C = 100$	B1 B1 cso See notes.
	$25 \equiv Ax(2x+1) + B(2x+1) + Cx^2$ $x=0, \quad 25 = B$ $x = -\frac{1}{2}, \quad 25 = \frac{1}{4}C \Rightarrow C = 100$ $x^2 \text{ terms: } 0 = 2A + C$ $0 = 2A + 100 \Rightarrow A = -50$ $x^2: 0 = 2A + C, \quad x: 0 = A + 2B,$ $\text{constant: } 25 = B$	Writes down a correct identity and attempts to find the value of either one of "A", "B" or "C". M1
	leading to $A = -50$ $\left\{ \frac{25}{x^2(2x+1)} \equiv -\frac{50}{x} + \frac{25}{x^2} + \frac{100}{(2x+1)} \right\}$	Correct value for "A" which is found using a correct identity and follows from their partial fraction decomposition. A1 [4]
(b)	$V = \pi \int_1^4 \left(\frac{5}{x\sqrt{(2x+1)}} \right)^2 dx$	For $\pi \int \left(\frac{5}{x\sqrt{(2x+1)}} \right)^2$ Ignore limits and dx. Can be implied. B1
	<p>For their partial fraction</p> $\left\{ \int \frac{25}{x^2(2x+1)} dx = \int -\frac{50}{x} + \frac{25}{x^2} + \frac{100}{(2x+1)} dx \right\}$ $= -50 \ln x + \frac{25x^{-1}}{(-1)} + \frac{100}{2} \ln(2x+1) \{+ c\}$	Either $\pm \frac{A}{x} \rightarrow \pm a \ln x$ or $\pm a \ln kx$ or $\pm \frac{B}{x^2} \rightarrow \pm b x^{-1}$ or $\frac{C}{(2x+1)} \rightarrow \pm c \ln(2x+1)$ At least two terms correctly integrated All three terms correctly integrated. M1 A1ft A1ft
	$\left\{ \int_1^4 \frac{25}{x^2(2x+1)} dx = \left[-50 \ln x - \frac{25}{x} + 50 \ln(2x+1) \right]_1^4 \right\}$ $= \left(-50 \ln 4 - \frac{25}{4} + 50 \ln 9 \right) - \left(0 - 25 + 50 \ln 3 \right)$ $= 50 \ln 9 - 50 \ln 4 - 50 \ln 3 - \frac{25}{4} + 25$ $= 50 \ln \left(\frac{3}{4} \right) + \frac{75}{4}$ <p>So, $V = \frac{75}{4} \pi + 50 \pi \ln \left(\frac{3}{4} \right)$ or allow $\pi \left(\frac{75}{4} + 50 \ln \left(\frac{3}{4} \right) \right)$</p>	Applies limits of 4 and 1 and subtracts the correct way round. dM1 A1 oe [6] 10

Question 4 Notes	
4. (a)	<p>BE CAREFUL! Candidates will assign <i>their own</i> "A, B and C" for this question.</p> <p>B1 At least one of "B" or "C" are correct.</p> <p>B1 Breaks up their partial fraction correctly into three terms and both "B" = 25 and "C" = 100.</p> <p>Note If a candidate does not give partial fraction decomposition then: <ul style="list-style-type: none"> the 2nd B1 mark can follow from a correct identity. </p> <p>M1 Writes down a correct identity (although this can be implied) and attempts to find the value of either one of "A" or "B" or "C". This can be achieved by either substituting values into their identity or comparing coefficients and solving the resulting equations simultaneously.</p> <p>A1 Correct value for "A" which is found using a correct identity and follows from their partial fraction decomposition.</p> <p>Note If a candidate does not give partial fraction decomposition then the final A1 mark can be awarded for a correct "A" if a candidate writes out their partial fractions at the end.</p> <p>Note The correct partial fraction from no working scores B1B1M1A1.</p> <p>Note A number of candidates will start this problem by writing out the correct identity and then attempt to find "A" or "B" or "C". Therefore the B1 marks can be awarded from this method.</p> <p>Note Award SC B1B0M0A0 for $\frac{25}{x^2(2x+1)} \equiv \frac{B}{x^2} + \frac{C}{(2x+1)}$ leading to "B" = 25 or "C" = 100</p>
(b)	<p>B1 For a correct statement of $\pi \int \left(\frac{5}{x\sqrt{2x+1}} \right)^2$ or $\pi \int \frac{25}{x^2(2x+1)}$. Ignore limits and dx. Can be implied.</p> <p>Note The π can only be recovered later from a correct expression.</p>
	<p>For their partial fraction, (not $\sqrt{\text{their partial fraction}}$), where A, B, C are "their" part (a) constants</p> <p>M1 Either $\pm \frac{A}{x} \rightarrow \pm a \ln x$ or $\pm \frac{B}{x^2} \rightarrow \pm b x^{-1}$ or $\frac{C}{(2x+1)} \rightarrow \pm c \ln(2x+1)$.</p> <p>Note $\sqrt{\frac{B}{x^2}} \rightarrow \frac{\sqrt{B}}{x}$ which integrates to $\sqrt{B} \ln x$ is not worthy of M1.</p> <p>A1ft At least two terms from any of $\pm \frac{A}{x}$ or $\pm \frac{B}{x^2}$ or $\frac{C}{(2x+1)}$ correctly integrated. Can be un-simplified.</p> <p>A1ft All 3 terms from $\pm \frac{A}{x}$, $\pm \frac{B}{x^2}$ and $\frac{C}{(2x+1)}$ correctly integrated. Can be un-simplified.</p> <p>Note The 1st A1 and 2nd A1 marks in part (b) are both follow through accuracy marks.</p>
	<p>dM1 Dependent on the previous M mark. Applies limits of 4 and 1 and subtracts the correct way round.</p> <p>A1 Final correct exact answer in the form $a + b \ln c$. i.e. either $\frac{75}{4}\pi + 50\pi \ln\left(\frac{3}{4}\right)$ or $50\pi \ln\left(\frac{3}{4}\right) + \frac{75}{4}\pi$ or $50\pi \ln\left(\frac{9}{12}\right) + \frac{75}{4}\pi$ or $\frac{75}{4}\pi - 50\pi \ln\left(\frac{4}{3}\right)$ or $\frac{75}{4}\pi + 25\pi \ln\left(\frac{9}{16}\right)$ etc. Also allow $\pi \left(\frac{75}{4} + 50 \ln\left(\frac{3}{4}\right) \right)$ or equivalent.</p> <p>Note A candidate who achieves full marks in (a), but then mixes up the correct constants when writing their partial fraction can only achieve a maximum of B1M1A1A0M1A0 in part (b).</p> <p>Note The π in the volume formula is only required for the B1 mark and the final A1 mark.</p>

<p>4. (b)</p>	<p>Alternative method of integration</p> $V = \pi \int_1^4 \left(\frac{5}{x\sqrt{(2x+1)}} \right)^2 dx$ $\int \frac{25}{x^2(2x+1)} dx ; u = \frac{1}{x} \Rightarrow \frac{du}{dx} = -\frac{1}{x^2}$ $= \int \frac{-25}{\left(\frac{2}{u}+1\right)} du = \int \frac{-25}{\left(\frac{2+u}{u}\right)} du = \int \frac{-25u}{(2+u)} du = -25 \int \frac{2+u-2}{(2+u)} du$ $= -25 \int 1 - \frac{2}{(2+u)} du = -25(u - 2\ln(2+u))$ $\left\{ \int_1^4 \frac{25}{x^2(2x+1)} dx = [-25u + 50\ln(2+u)]_1^4 \right\}$ $= \left(-\frac{25}{4} + 50\ln\left(\frac{9}{4}\right) \right) - (-25 + 50\ln 3)$ $= 50\ln\left(\frac{9}{4}\right) - 50\ln 3 - \frac{25}{4} + 25$ $= 50\ln\left(\frac{3}{4}\right) + \frac{75}{4}$ <p>So, $V = \frac{75}{4}\pi + 50\pi\ln\left(\frac{3}{4}\right)$</p>	<table border="1"> <tr> <td data-bbox="810 190 901 376">B1</td> <td data-bbox="901 190 1549 376">For $\pi \int \left(\frac{5}{x\sqrt{(2x+1)}} \right)^2$ Ignore limits and dx. Can be implied.</td> </tr> <tr> <td data-bbox="810 376 901 555">M1</td> <td data-bbox="901 376 1549 555">Achieves $\pm \alpha \pm \frac{\beta}{(k+u)}$ and integrates to give either $\pm \alpha u$ or $\pm \beta \ln(k+u)$</td> </tr> <tr> <td data-bbox="810 555 901 689">A1</td> <td data-bbox="901 555 1549 689">Dependent on the M mark. Either $-25u$ or $50\ln(2+u)$</td> </tr> <tr> <td data-bbox="810 689 901 801">A1</td> <td data-bbox="901 689 1549 801">$-25(u - 2\ln(2+u))$</td> </tr> <tr> <td data-bbox="810 801 901 1292">dM1</td> <td data-bbox="901 801 1549 1292">Applies limits of $\frac{1}{4}$ and 1 in u or 4 and 1 in x in their integrated function and subtracts the correct way round.</td> </tr> <tr> <td data-bbox="810 1292 901 2175">A1</td> <td data-bbox="901 1292 1549 2175">$\frac{75}{4}\pi + 50\pi\ln\left(\frac{3}{4}\right)$ or allow $\pi\left(\frac{75}{4} + 50\ln\left(\frac{3}{4}\right)\right)$</td> </tr> </table>	B1	For $\pi \int \left(\frac{5}{x\sqrt{(2x+1)}} \right)^2$ Ignore limits and dx. Can be implied.	M1	Achieves $\pm \alpha \pm \frac{\beta}{(k+u)}$ and integrates to give either $\pm \alpha u$ or $\pm \beta \ln(k+u)$	A1	Dependent on the M mark. Either $-25u$ or $50\ln(2+u)$	A1	$-25(u - 2\ln(2+u))$	dM1	Applies limits of $\frac{1}{4}$ and 1 in u or 4 and 1 in x in their integrated function and subtracts the correct way round.	A1	$\frac{75}{4}\pi + 50\pi\ln\left(\frac{3}{4}\right)$ or allow $\pi\left(\frac{75}{4} + 50\ln\left(\frac{3}{4}\right)\right)$
B1	For $\pi \int \left(\frac{5}{x\sqrt{(2x+1)}} \right)^2$ Ignore limits and dx. Can be implied.													
M1	Achieves $\pm \alpha \pm \frac{\beta}{(k+u)}$ and integrates to give either $\pm \alpha u$ or $\pm \beta \ln(k+u)$													
A1	Dependent on the M mark. Either $-25u$ or $50\ln(2+u)$													
A1	$-25(u - 2\ln(2+u))$													
dM1	Applies limits of $\frac{1}{4}$ and 1 in u or 4 and 1 in x in their integrated function and subtracts the correct way round.													
A1	$\frac{75}{4}\pi + 50\pi\ln\left(\frac{3}{4}\right)$ or allow $\pi\left(\frac{75}{4} + 50\ln\left(\frac{3}{4}\right)\right)$													

Question Number	Scheme	Marks
5. (a)	From question, $V = \frac{4}{3}\pi r^3$, $S = 4\pi r^2$, $\frac{dV}{dt} = 3$ $\left\{ V = \frac{4}{3}\pi r^3 \Rightarrow \right\} \frac{dV}{dr} = 4\pi r^2$ $\frac{dV}{dr} = 4\pi r^2$ (Can be implied)	B1 oe
	$\left\{ \frac{dV}{dr} \times \frac{dr}{dt} = \frac{dV}{dt} \Rightarrow \right\} (4\pi r^2) \frac{dr}{dt} = 3$ $\left(\text{Candidate's } \frac{dV}{dr} \right) \times \frac{dr}{dt} = 3$ $\left\{ \frac{dr}{dt} = \frac{dV}{dt} \div \frac{dV}{dr} \Rightarrow \right\} \frac{dr}{dt} = (3) \frac{1}{4\pi r^2}; \left\{ = \frac{3}{4\pi r^2} \right\}$ or $3 \div \text{Candidate's } \frac{dV}{dr}$;	M1 oe
	When $r = 4\text{cm}$, $\frac{dr}{dt} = \frac{3}{4\pi(4)^2} \left\{ = \frac{3}{64\pi} \right\}$ dependent on previous M1. see notes	dM1
	Hence, $\frac{dr}{dt} = 0.01492077591\dots(\text{cm}^2 \text{ s}^{-1})$ anything that rounds to 0.0149	A1 [4]
(b)	$\left\{ \frac{dS}{dt} = \frac{dS}{dr} \times \frac{dr}{dt} = \right\} \Rightarrow \frac{dS}{dt} = 8\pi r \times \frac{3}{4\pi r^2} \left\{ \text{or } \frac{6}{r} \text{ or } 8\pi r \times 0.0149\dots \right\}$ $8\pi r \times \text{Candidate's } \frac{dr}{dt}$	M1; oe
	When $r = 4\text{cm}$, $\frac{dr}{dt} = 8\pi(4) \times \frac{3}{4\pi(4)^2}$ or $\frac{6}{4}$ or $8\pi(4) \times 0.0149\dots$ Hence, $\frac{dS}{dt} = 1.5 (\text{cm}^2 \text{ s}^{-1})$ anything that rounds to 1.5	A1 cs [2] 6

Question 5 Notes

(a)	B1 $\frac{dV}{dr} = 4\pi r^2$ Can be implied by later working.
	M1 $\left(\text{Candidate's } \frac{dV}{dr} \right) \times \frac{dr}{dt} = 3$ or $3 \div \text{Candidate's } \frac{dV}{dr}$
	dM1 (dependent on the previous method mark) Substitutes $r = 4$ into an expression which is a result of a quotient of “3” and their $\frac{dV}{dr}$.
	A1 anything that rounds to 0.0149 (units are not required)
(b)	M1 $8\pi r \times \text{Candidate's } \frac{dr}{dt}$
	A1 anything that rounds to 1.5 (units are not required). Correct solution only.
	Note Using $\frac{dr}{dt} = 0.0149$ gives $\frac{dS}{dt} = 1.4979\dots$ which is fine for A1.

Question Number	Scheme	Marks
<p>6.</p> <p>(a)</p> <p>(b)</p>	<p>$l_1: \mathbf{r} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} + \lambda \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}, \quad l_2: \mathbf{r} = \begin{pmatrix} 7 \\ 0 \\ 7 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix} \quad \overline{OA} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, \overline{OB} = \begin{pmatrix} 4 \\ p \\ 3 \end{pmatrix}$</p> <p>$\{B \text{ lies on } l_2 \Rightarrow \mu = -1 \Rightarrow\} \quad p = 5$</p> <p>$\{l_1 = l_2 \Rightarrow\} \begin{cases} \mathbf{i}: & 1 = 7 + 3\mu \\ \mathbf{j}: & 2 + 2\lambda = -5\mu \\ \mathbf{k}: & 3 - \lambda = 7 + 4\mu \end{cases}$</p> <p>e.g. $\mathbf{i}: 7 + 3\mu = 1$</p> <p>So, $\mu = -2$</p> <p>Point of intersection is $\overline{OC} = \mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p> <p>Finds $\lambda = 4$ and either</p> <ul style="list-style-type: none"> checks $\lambda = 4$ and $\mu = -2$ is true for the third component. substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ <p>and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p>	<p>A lies on l_1 and B lies on l_2</p> <p>$p = 5$</p> <p>B1</p> <p>[1]</p> <p>Writes down an equation involving only one parameter. $\mu = -2$</p> <p>M1 A1 B1</p> <p>B1</p> <p>[4]</p>
<p>(b)</p>	<p>Alternative Method: Solving \mathbf{j} and \mathbf{k} simultaneously gives</p> <p>$8 = 14 + 3\mu$ or $23 + 3\lambda = 35$</p> <p>So, $\mu = -2$ or $\lambda = 4$</p> <p>Point of intersection is $\overline{OC} = \mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p> <p>Finds $\lambda = 4$ and either</p> <ul style="list-style-type: none"> checks $\mu = -2$ is true for the \mathbf{i} component. substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ <p>and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p>	<p>Writes down an equation involving only one parameter. Either $\mu = -2$ or $\lambda = 4$</p> <p>$\mathbf{i} + 10\mathbf{j} - \mathbf{k}$</p> <p>M1 A1 B1</p> <p>B1</p> <p>[4]</p>
<p>(c)</p> <p>(d)</p>	<p>$\overline{AC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix}$ and $\overline{BC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix}$</p> <p>$\pm \left(\begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \cdot \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix} \right)$</p> <p>$\cos ACB = \frac{\overline{AC} \cdot \overline{BC}}{ \overline{AC} \cdot \overline{BC} } = \frac{\pm \left(\begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \cdot \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix} \right)}{\sqrt{(0)^2 + (8)^2 + (-4)^2} \cdot \sqrt{(-3)^2 + (5)^2 + (-4)^2}}$</p> <p>$\left\{ \cos ACB = \frac{0 + 40 + 16}{\sqrt{80} \cdot \sqrt{50}} = \frac{56}{\sqrt{4000}} \Rightarrow \right\} ACB = 27.69446... = 27.7$ (3 sf)</p> <p>Area $ACB = \frac{1}{2}(\sqrt{80})(\sqrt{50})\sin 27.69446...^\circ = 14.696888...$</p>	<p>An attempt to find both the vectors $(\overline{AC}$ or $\overline{CA})$ and $(\overline{BC}$ or $\overline{CB})$.</p> <p>Applies dot product formula between their $(\overline{AC}$ or $\overline{CA})$ and their $(\overline{BC}$ or $\overline{CB})$.</p> <p>Anything that rounds to 27.7</p> <p>See notes</p> <p>Anything that rounds to 14.7</p> <p>M1 M1 A1</p> <p>A1</p> <p>[3]</p> <p>M1 A1</p> <p>[2] 10</p>

Question 6: Alternative Methods for Part (c)

<p>6. (c)</p>	<p>Alternative Method 1: Using the direction vectors of Line 1 and Line 2</p> $\mathbf{d}_1 = \begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix}, \quad \mathbf{d}_2 = \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix}$ $\cos \theta = \frac{\mathbf{d}_1 \cdot \mathbf{d}_2}{ \mathbf{d}_1 \mathbf{d}_2 } = \frac{\begin{pmatrix} 0 \\ 2 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -5 \\ 4 \end{pmatrix}}{\sqrt{(0)^2 + (2)^2 + (-1)^2} \cdot \sqrt{(3)^2 + (-5)^2 + (4)^2}}$ $\left\{ \cos \theta = \frac{0 - 10 - 4}{\sqrt{5} \cdot \sqrt{50}} = \frac{-7\sqrt{10}}{25} \Rightarrow \right\} \theta = 152.3054385\dots$ <p>Angle $ACB = 180 - 152.3054385\dots = 27.69446145\dots = 27.7$ (3 sf)</p>	<p>Applies dot product formula between their \mathbf{d}_1 and \mathbf{d}_2</p> <p>M2</p> <p>Anything that rounds to 27.7</p> <p>A1</p> <p>[3]</p>
	<p>Alternative Method 2: The Cosine Rule</p> $\overrightarrow{AC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \quad \text{and} \quad \overrightarrow{BC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix}$ <p>Also $\overrightarrow{AB} = \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \\ 0 \end{pmatrix}$</p> <p>Note $\overrightarrow{AC} = \sqrt{80}$, $\overrightarrow{BC} = \sqrt{50}$ and $\overrightarrow{AB} = \sqrt{18}$</p> $(\sqrt{18})^2 = (\sqrt{80})^2 + (\sqrt{50})^2 - 2(\sqrt{80})(\sqrt{50})\cos \theta$ $\left\{ \cos \theta = \frac{7\sqrt{10}}{25} \right\} \Rightarrow \theta = 27.69446145\dots = 27.7$ (3 sf)	<p>An attempt to find both the vectors (\overrightarrow{AC} or \overrightarrow{CA}) and (\overrightarrow{BC} or \overrightarrow{CB}).</p> <p>M1</p> <p>Applies the cosine rule the correct way round.</p> <p>Anything that rounds to 27.7</p> <p>M1 oe</p> <p>A1</p> <p>[3]</p>
	<p>Alternative Method 3: Vector Cross Product</p> <p>Only apply this scheme if it is clear that a candidate is applying a vector cross product method.</p> $\overrightarrow{AC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \quad \text{and} \quad \overrightarrow{BC} = \begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix}$ $\overrightarrow{AC} \times \overrightarrow{BC} = \begin{pmatrix} 0 \\ 8 \\ -4 \end{pmatrix} \times \begin{pmatrix} -3 \\ 5 \\ -4 \end{pmatrix} = \left\{ \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 8 & -4 \\ -3 & 5 & -4 \end{vmatrix} = 24\mathbf{i} + 12\mathbf{j} + 24\mathbf{k} \right\}$ $\sin ACB = \frac{\sqrt{(24)^2 + (12)^2 + (12)^2}}{\sqrt{(0)^2 + (8)^2 + (-4)^2} \cdot \sqrt{(-3)^2 + (5)^2 + (-4)^2}}$ $\left\{ \sin ACB = \frac{\sqrt{864}}{\sqrt{80} \cdot \sqrt{50}} = \frac{3\sqrt{15}}{25} \Rightarrow \right\} \theta = 27.69446145\dots = 27.7$ (3 sf)	<p>An attempt to find both the vectors (\overrightarrow{AC} or \overrightarrow{CA}) and (\overrightarrow{BC} or \overrightarrow{CB}).</p> <p>M1</p> <p>Full method for applying the vector cross product formula between their (\overrightarrow{AC} or \overrightarrow{CA}) and their (\overrightarrow{BC} or \overrightarrow{CB}).</p> <p>M1</p> <p>Anything that rounds to 27.7</p> <p>A1</p> <p>[3]</p>

Question 6 Notes		
6. (a)	B1	$p = 5$ (Ignore working.)
(b)		Method 1
	M1	Writes down an equation involving only one parameter. This equation will usually be $7 + 3\mu = 1$ which is found from equating the i components of l_1 and l_2 .
	A1	Finds $\mu = -2$
	B1	Point of intersection of $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$. Allow $(1, 10, -1)$ or $\begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix}$.
	B1	Finds $\lambda = 4$ and either <ul style="list-style-type: none"> • checks $\lambda = 4$ and $\mu = -2$ is true for the third component. • substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$
(b)		Alternative Method
	M1	Writes down an equation involving only one parameter. Solving the j and k components simultaneously will usually give either $8 = 14 + 3\mu$ or $23 + 3\lambda = 35$
	A1	Finds either $\mu = -2$ or $\lambda = 4$
	B1	Point of intersection of $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$. Allow $(1, 10, -1)$ or $\begin{pmatrix} 1 \\ 10 \\ -1 \end{pmatrix}$.
	B1	Finds $\lambda = 4$ and either <ul style="list-style-type: none"> • checks $\mu = -2$ is true for the i component. • substitutes $\mu = -2$ into l_1 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$ and substitutes $\lambda = 4$ into l_2 to give $\mathbf{i} + 10\mathbf{j} - \mathbf{k}$
(c)	M1	An attempt to find both the vectors $(\overline{AC}$ or $\overline{CA})$ and $(\overline{BC}$ or $\overline{CB})$ by subtracting.
	M1	Applies dot product formula between their $(\overline{AC}$ or $\overline{CA})$ and their $(\overline{BC}$ or $\overline{CB})$.
	A1	anything that rounds to 27.7
	Note	An answer of 0.48336... in radians without the correct answer in degrees is A0.
	Note	Some candidates will apply the dot product formula between vectors which are the wrong way round and achieve 152.3054385...°. If they give the acute equivalent of awrt 27.7 then award A1.
(d)	M1	$\frac{1}{2}(\text{their length } AC)(\text{their length } BC)\sin(\text{their } 27.7^\circ \text{ from part (c)})$
	A1	anything that rounds to 14.7. Also allow $6\sqrt{6}$.
	Note	Area $ACB = \frac{1}{2}(\sqrt{80})(\sqrt{50})\sin(152.3054385...^\circ) = \text{awrt } 14.7$ is M1A1.

Question Number	Scheme	Marks												
7. (a)	$\frac{dN}{dt} = \frac{(kt - 1)(5000 - N)}{t}, \quad t > 0, \quad 0 < N < 5000$ $\int \frac{1}{5000 - N} dN = \int \frac{(kt - 1)}{t} dt \quad \left\{ \text{or} = \int \left(k - \frac{1}{t} \right) dt \right\}$ $-\ln(5000 - N) = kt - \ln t; + c$ <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%;"><i>then eg either...</i></td> <td style="width: 33%;"><i>or...</i></td> <td style="width: 33%;"><i>or...</i></td> </tr> <tr> <td>$-kt + c = \ln(5000 - N) - \ln t$</td> <td>$kt + c = \ln t - \ln(5000 - N)$</td> <td>$\ln(5000 - N) = -kt + \ln t + c$</td> </tr> <tr> <td>$-kt + c = \ln\left(\frac{5000 - N}{t}\right)$</td> <td>$kt + c = \ln\left(\frac{t}{5000 - N}\right)$</td> <td>$5000 - N = e^{-kt + \ln t + c}$</td> </tr> <tr> <td>$e^{-kt + c} = \frac{5000 - N}{t}$</td> <td>$e^{kt + c} = \frac{t}{5000 - N}$</td> <td>$5000 - N = te^{-kt + c}$</td> </tr> </table>	<i>then eg either...</i>	<i>or...</i>	<i>or...</i>	$-kt + c = \ln(5000 - N) - \ln t$	$kt + c = \ln t - \ln(5000 - N)$	$\ln(5000 - N) = -kt + \ln t + c$	$-kt + c = \ln\left(\frac{5000 - N}{t}\right)$	$kt + c = \ln\left(\frac{t}{5000 - N}\right)$	$5000 - N = e^{-kt + \ln t + c}$	$e^{-kt + c} = \frac{5000 - N}{t}$	$e^{kt + c} = \frac{t}{5000 - N}$	$5000 - N = te^{-kt + c}$	<p>See notes B1</p> <p>See notes M1 A1; A1</p>
<i>then eg either...</i>	<i>or...</i>	<i>or...</i>												
$-kt + c = \ln(5000 - N) - \ln t$	$kt + c = \ln t - \ln(5000 - N)$	$\ln(5000 - N) = -kt + \ln t + c$												
$-kt + c = \ln\left(\frac{5000 - N}{t}\right)$	$kt + c = \ln\left(\frac{t}{5000 - N}\right)$	$5000 - N = e^{-kt + \ln t + c}$												
$e^{-kt + c} = \frac{5000 - N}{t}$	$e^{kt + c} = \frac{t}{5000 - N}$	$5000 - N = te^{-kt + c}$												
leading to $N = 5000 - Ate^{-kt}$ with no incorrect working/statements. See notes		A1 * cs0												
(b)	$\{t = 1, N = 1200 \Rightarrow\} \quad 1200 = 5000 - Ae^{-k}$ $\{t = 2, N = 1800 \Rightarrow\} \quad 1800 = 5000 - 2Ae^{-2k}$ <p>So $Ae^{-k} = 3800$</p> <p>and $2Ae^{-2k} = 3200$ or $Ae^{-2k} = 1600$</p> <p>Eg. $\frac{e^{-k}}{2e^{-2k}} = \frac{3800}{3200}$ or $\frac{2e^{-2k}}{e^{-k}} = \frac{3200}{3800}$</p> <p>So $\frac{1}{2}e^k = \frac{3800}{3200}$ or $2e^{-k} = \frac{3200}{3800}$</p> $k = \ln\left(\frac{7600}{3200}\right) \text{ or equivalent } \left\{ \text{eg } k = \ln\left(\frac{19}{8}\right) \right\}$ $\left\{ A = 3800(e^k) = 3800\left(\frac{19}{8}\right) \Rightarrow \right\} A = 9025$	<p>At least one correct statement written down using the boundary conditions B1</p> <p>An attempt to eliminate A by producing an equation in only k. M1</p> <p>At least one of $A = 9025$ cao or $k = \ln\left(\frac{7600}{3200}\right)$ or exact equivalent A1</p> <p>Both $A = 9025$ cao or $k = \ln\left(\frac{7600}{3200}\right)$ or exact equivalent A1</p> <p style="text-align: right;">[4]</p>												
<u>Alternative Method for the M1 mark in (b)</u>														
$e^{-k} = \frac{3800}{A}$ $2A\left(\frac{3800}{A}\right)^2 = 3200$		<p>An attempt to eliminate k by producing an equation in only A M1</p>												
(c)	$\left\{ t = 5, N = 5000 - 9025(5)e^{-5\ln\left(\frac{19}{8}\right)} \right\}$ <p>$N = 4402.828401... = 4400$ (fish) (nearest 100)</p>	<p>anything that rounds to 4400 B1</p> <p style="text-align: right;">[1] 10</p>												

Question 7 Notes	
7. (a)	<p>B1 Separates variables as shown. dN and dt should be in the correct positions, though this mark can be implied by later working. Ignore the integral signs.</p> <p>M1 Either $\pm \lambda \ln(5000 - N)$ or $\pm \lambda \ln(N - 5000)$ or $kt - \ln t$ where $\lambda \neq 0$ is a constant.</p> <p>A1 For $-\ln(5000 - N) = kt - \ln t$ or $\ln(5000 - N) = -kt + \ln t$ or $-\frac{1}{k} \ln(5000 - N) = t - \frac{1}{k} \ln t$ or</p> <p>A1 which is dependent on the 1st M1 mark being awarded.</p> <p>For applying a constant of integration, eg. $+c$ or $+\ln e^c$ or $+\ln c$ or A to their integrated equation $+c$ can be on either side of their equation for the 2nd A1 mark.</p> <p>Note</p> <p>A1 Uses a constant of integration eg. "c" or "$\ln e^c$" "$\ln c$" or and applies a fully correct method to prove the result $N = 5000 - Ate^{-kt}$ with no incorrect working seen. (Correct solution only.)</p> <p>NOTE IMPORTANT</p> <p>There needs to be an intermediate stage of justifying the A and the e^{-kt} in Ate^{-kt} by for example</p> <ul style="list-style-type: none"> • either $5000 - N = e^{\ln t - kt + c}$ • or $5000 - N = t e^{-kt + c}$ • or $5000 - N = t e^{-kt} e^c$ <p>or equivalent needs to be stated before achieving $N = 5000 - Ate^{-kt}$</p>
(b)	<p>B1 At least one of either $1200 = 5000 - Ae^{-k}$ (or equivalent) or $1800 = 5000 - 2Ae^{-2k}$ (or equivalent)</p> <p>M1</p> <ul style="list-style-type: none"> • Either an attempt to eliminate A by producing an equation in only k. • or an attempt to eliminate k by producing an equation in only A <p>A1 At least one of $A = 9025$ or $k = \ln\left(\frac{7600}{3200}\right)$ or equivalent</p> <p>A1 Both $A = 9025$ or $k = \ln\left(\frac{7600}{3200}\right)$ or equivalent</p> <p>Note Alternative correct values for k are $k = \ln\left(\frac{19}{8}\right)$ or $k = -\ln\left(\frac{8}{19}\right)$ or $k = \ln 7600 - \ln 3200$</p> <p>or $k = -\ln\left(\frac{3800}{9025}\right)$ or equivalent.</p> <p>Note $k = 0.8649...$ without a correct exact equivalent is A0.</p>
(c)	<p>B1 anything that rounds to 4400</p>

8.




Figure 3

The curve shown in Figure 3 has parametric equations

$$x = t - 4 \sin t, \quad y = 1 - 2 \cos t, \quad -\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$$

The point A , with coordinates $(k, 1)$, lies on the curve.

Given that $k > 0$

(a) find the exact value of k , (2)

(b) find the gradient of the curve at the point A . (4)

There is one point on the curve where the gradient is equal to $-\frac{1}{2}$

(c) Find the value of t at this point, showing each step in your working and giving your answer to 4 decimal places.

[Solutions based entirely on graphical or numerical methods are not acceptable.] (6)



Question Number	Scheme	Marks
<p>8.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>$x = t - 4\sin t, \quad y = 1 - 2\cos t, \quad -\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$ $A(k, 1)$ lies on the curve, $k > 0$</p> <p>{When $y = 1$,} $1 = 1 - 2\cos t \Rightarrow t = -\frac{\pi}{2}, \frac{\pi}{2}$ Sets $y = 1$ to find t and uses their t to find x.</p> <p>k (or x) $= \frac{\pi}{2} - 4\sin\left(\frac{\pi}{2}\right)$ or $x = -\frac{\pi}{2} - 4\sin\left(-\frac{\pi}{2}\right)$</p> <p>{When $t = -\frac{\pi}{2}, k > 0$,} so $k = 4 - \frac{\pi}{2}$ or $\frac{8 - \pi}{2}$ x or $k = 4 - \frac{\pi}{2}$</p> <hr/> <p>$\frac{dx}{dt} = 1 - 4\cos t, \quad \frac{dy}{dt} = 2\sin t$ At least one of $\frac{dx}{dt}$ or $\frac{dy}{dt}$ correct.</p> <p>Both $\frac{dx}{dt}$ and $\frac{dy}{dt}$ are correct.</p> <hr/> <p>So, $\frac{dy}{dx} = \frac{2\sin t}{1 - 4\cos t}$ Applies their $\frac{dy}{dt}$ divided by their $\frac{dx}{dt}$ and substitutes their t into their $\frac{dy}{dx}$.</p> <p>At $t = -\frac{\pi}{2}, \frac{dy}{dx} = \frac{2\sin\left(-\frac{\pi}{2}\right)}{1 - 4\cos\left(-\frac{\pi}{2}\right)}; = -2$ Correct value for $\frac{dy}{dx}$ of -2</p> <hr/> <p>$\frac{2\sin t}{1 - 4\cos t} = -\frac{1}{2}$ Sets their $\frac{dy}{dx} = -\frac{1}{2}$</p> <p>gives $4\sin t - 4\cos t = -1$ See notes</p> <p>So $4\sqrt{2}\sin\left(t - \frac{\pi}{4}\right); = -1$ or $-4\sqrt{2}\cos\left(t + \frac{\pi}{4}\right); = -1$ See notes</p> <p>$t = \sin^{-1}\left(\frac{-1}{4\sqrt{2}}\right) + \frac{\pi}{4}$ or $t = \cos^{-1}\left(\frac{1}{4\sqrt{2}}\right) - \frac{\pi}{4}$ See notes</p> <p>$t = 0.6076875626... = 0.6077$ (4 dp) anything that rounds to 0.6077</p>	<p>M1</p> <p>A1</p> <p>[2]</p> <p>B1</p> <p>B1</p> <p>M1;</p> <p>A1</p> <p>cao cso</p> <p>[4]</p> <p>M1</p> <p>A1</p> <p>M1; A1</p> <p>dM1</p> <p>A1</p> <p>[6]</p> <p>12</p>
Question 8 Notes		
<p>(c)</p>	<p>NOTE <u>VERY IMPORTANT NOTE FOR PART (c)</u> Candidates who state $t = 0.6077$ with no intermediate working from $4\sin t - 4\cos t = -1$ will get 2nd M0, 2nd A0, 3rd M0, 3rd A0.</p> <p>They will not express $4\sin t - 4\cos t$ as either $4\sqrt{2}\sin\left(t - \frac{\pi}{4}\right)$ or $-4\sqrt{2}\cos\left(t + \frac{\pi}{4}\right)$.</p> <p>OR use any acceptable alternative method to achieve $t = 0.6077$</p> <p>NOTE Alternative methods for part (c) are given on the next page.</p>	

Question 8: Alternative Methods for Part (c)	
8. (c)	<p>Alternative Method 1:</p> $\frac{2 \sin t}{1 - 4 \cos t} = -\frac{1}{2}$ <p style="text-align: right;">Sets their $\frac{dy}{dx} = -\frac{1}{2}$ M1</p> <p>eg. $\left(\frac{2 \sin t}{1 - 4 \cos t}\right)^2 = \frac{1}{4}$ or $(4 \sin t)^2 = (4 \cos t - 1)^2$ Squaring to give a correct equation. A1 This mark can be implied by a "squared" correct equation.</p> <p style="text-align: center;">Note: You can also give 1st A1 in this method for $4 \sin t - 4 \cos t = -1$ as in the main scheme.</p> <p style="text-align: center;">Squares their equation, applies $\sin^2 t + \cos^2 t = 1$ and achieves a three term quadratic equation of the form $\pm a \cos^2 t \pm b \cos t \pm c = 0$ M1 or $\pm a \sin^2 t \pm b \sin t \pm c = 0$ or eg. $\pm a \cos^2 t \pm b \cos t = \pm c$ where $a \neq 0, b \neq 0$ and $c \neq 0$.</p> <ul style="list-style-type: none"> • Either $32 \cos^2 t - 8 \cos t - 15 = 0$ • or $32 \sin^2 t + 8 \sin t - 15 = 0$ For a correct three term quadratic equation. A1 • Either $\cos t = \frac{8 \pm \sqrt{1984}}{64} = \frac{1 + \sqrt{31}}{8} \Rightarrow t = \cos^{-1}(\dots)$ which is dependent on the 2nd M1 mark. dM1 Uses correct algebraic processes to give $t = \dots$ • or $\sin t = \frac{-8 \pm \sqrt{1984}}{64} = \frac{-1 \pm \sqrt{31}}{8} \Rightarrow t = \sin^{-1}(\dots)$ anything that rounds to 0.6077 A1 $t = 0.6076875626\dots = 0.6077$ (4 dp) <p style="text-align: right;">[6]</p>
8. (c)	<p>Alternative Method 2:</p> $\frac{2 \sin t}{1 - 4 \cos t} = -\frac{1}{2}$ <p style="text-align: right;">Sets their $\frac{dy}{dx} = -\frac{1}{2}$ M1</p> <p>eg. $(4 \sin t - 4 \cos t)^2 = (-1)^2$ Squaring to give a correct equation. A1 This mark can be implied by a correct equation. Note: You can also give 1st A1 in this method for $4 \sin t - 4 \cos t = -1$ as in the main scheme.</p> <p>So $16 \sin^2 t - 32 \sin t \cos t + 16 \cos^2 t = 1$</p> <p style="text-align: center;">Squares their equation, applies both $\sin^2 t + \cos^2 t = 1$ and $\sin 2t = 2 \sin t \cos t$ and then achieves an equation of the form $\pm a \pm b \sin 2t = \pm c$ M1</p> <p>leading to $16 - 16 \sin 2t = 1$ $16 - 16 \sin 2t = 1$ or equivalent. A1</p> <p>$\left\{ \sin 2t = \frac{15}{16} \Rightarrow \right\} t = \frac{\sin^{-1}(\dots)}{2}$ which is dependent on the 2nd M1 mark. dM1 Uses correct algebraic processes to give $t = \dots$ anything that rounds to 0.6077 A1</p> <p style="text-align: right;">[6]</p>

Question 8 Notes		
8. (a)	M1	Sets $y = 1$ to find t and uses their t to find x .
	Note	M1 can be implied by either x or $k = 4 - \frac{\pi}{2}$ or 2.429... or $\frac{\pi}{2} - 4$ or $-2.429...$
	A1	x or $k = 4 - \frac{\pi}{2}$ or $\frac{8 - \pi}{2}$
	Note	A decimal answer of 2.429... (without a correct exact answer) is A0.
	Note	Allow A1 for a candidate using $t = \frac{\pi}{2}$ to find $x = \frac{\pi}{2} - 4$ and then stating that k must be $4 - \frac{\pi}{2}$ o.e.
(b)	B1	At least one of $\frac{dx}{dt}$ or $\frac{dy}{dt}$ correct. Note: that this mark can be implied from their working.
	B1	Both $\frac{dx}{dt}$ and $\frac{dy}{dt}$ are correct. Note: that this mark can be implied from their working.
	M1	Applies their $\frac{dy}{dt}$ divided by their $\frac{dx}{dt}$ and attempts to substitute their t into their expression for $\frac{dy}{dx}$.
	Note	This mark may be implied by their final answer. i.e. $\frac{dy}{dx} = \frac{2\sin t}{1 - 4\cos t}$ followed by an answer of -2 (from $t = -\frac{\pi}{2}$) or 2 (from $t = \frac{\pi}{2}$)
	Note	Applying $\frac{dx}{dt}$ divided by their $\frac{dy}{dt}$ is M0, even if they state $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$.
	A1	Using $t = -\frac{\pi}{2}$ (and not $t = \frac{3\pi}{2}$) to find a correct $\frac{dy}{dx}$ of -2 by correct solution only .
(c)	NOTE	If a candidate uses an incorrect $\frac{dy}{dx}$ expression in part (c) then the accuracy marks are not obtainable.
	1st M1	Sets their $\frac{dy}{dx} = -\frac{1}{2}$
	1st A1	Rearranges to give the correct equation with $\sin t$ and $\cos t$ on the same side . eg. $4\sin t - 4\cos t = -1$ or $4\cos t - 4\sin t = 1$ or $\sin t - \cos t = -\frac{1}{4}$ or $\cos t - \sin t = \frac{1}{4}$ or $4\sin t - 4\cos t + 1 = 0$ or $4\cos t - 4\sin t - 1 = 0$ or $\sin t - \cos t + \frac{1}{4} = 0$ etc. are fine for A1.
	2nd M1	Rewrites $\pm\lambda\sin t \pm \mu\cos t$ in the form of either $R\cos(t \pm \alpha)$ or $R\sin(t \pm \alpha)$ where $R \neq 1$ or 0 and $\alpha \neq 0$
	2nd A1	Correct equation. Eg. $4\sqrt{2}\sin\left(t - \frac{\pi}{4}\right) = -1$ or $-4\sqrt{2}\cos\left(t + \frac{\pi}{4}\right) = -1$ or $\sqrt{2}\sin\left(t - \frac{\pi}{4}\right) = -\frac{1}{4}$ or $\sqrt{2}\cos\left(t + \frac{\pi}{4}\right) = \frac{1}{4}$, etc.
	Note	Unless recovered, give A0 for $4\sqrt{2}\sin(t - 45^\circ) = -1$ or $-4\sqrt{2}\cos(t + 45^\circ) = -1$, etc.
	3rd M1	which is dependent on the 2nd M1 mark. Uses correct algebraic processes to give $t = \dots$
	4th A1	anything that rounds to 0.6077
	Note	Do not give the final A1 mark in (c) if there any extra solutions given in the range $-\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$.
	Note	You can give the final A1 mark in (c) if extra solutions are given outside of $-\frac{2\pi}{3} \leq t \leq \frac{2\pi}{3}$.