

Write your name here

Surname

Other names

**Pearson Edexcel**  
**International**  
**Advanced Level**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

# Mechanics M1

## Advanced/Advanced Subsidiary

Tuesday 20 January 2015 – Morning

**Time: 1 hour 30 minutes**

Paper Reference

**WME01/01****You must have:**

Mathematical Formulae and Statistical Tables (Blue)

Total Marks

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$ , and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

### Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P45061A

©2015 Pearson Education Ltd.

1/5/5/5/



PEARSON

Leave  
blank


1. A railway truck  $A$  of mass  $m$  and a second railway truck  $B$  of mass  $4m$  are moving in opposite directions on a smooth straight horizontal track when they collide directly. Immediately before the collision the speed of truck  $A$  is  $3u$  and the speed of truck  $B$  is  $2u$ . In the collision the trucks join together. Modelling the trucks as particles, find
- (a) the speed of  $A$  immediately after the collision, (3)
- (b) the direction of motion of  $A$  immediately after the collision, (1)
- (c) the magnitude of the impulse exerted by  $A$  on  $B$  in the collision. (3)



Jan 2015 (IAL) Post QPEC  
6677 Mechanics M1  
Mark Scheme

Question Number	Scheme	Marks
1.(a)	$8mu - 3mu = 5mv$ $v = u$	M1 A1 A1 (3)
(b)	Original direction of motion of B o.e.	B1 (1)
(c)	For A: $I = m(u - -3u)$ OR    For B: $I = 4m(-u - -2u)$  $= 4mu$ $= 4mu$	M1 A1 A1 (3) <b>7</b>
<p style="text-align: center;"><b><u>NOTES</u></b></p> <p><b><u>Question 1(a)</u></b> M1 for attempt at CLM equation, with correct no. of terms, dimensionally correct. Allow consistent extra g's and cancelled m's and sign errors. (M1 if they find the impulse on each particle <i>and eliminate</i> the impulse to give an equation – then use above criteria for their equation) First A1 for a correct equation. (<math>3mu - 8mu = 5mv</math> or <math>-5mv</math> oe) Second A1 for <math>u</math> (-<math>u</math> A0) N.B. Allow <math>u</math>'s to be dropped or omitted in the equation if <math>u</math> is inserted in answer at the end. (Full marks can be scored). However, if <math>u</math> is not inserted then M0.</p> <p><b><u>Question 1(b)</u></b> B1 for (original) direction of B or opposite to original direction (of A) oe. (B0 for 'left' or direction changed). N.B. Must follow from <math>v = u</math> or <math>-u</math> obtained in (a).</p> <p><b><u>Question 1(c)</u></b> M1 for attempt at impulse = difference in momenta, for either particle, (must be considering <i>one</i> particle) (M0 if g's are included or if <math>m</math> omitted or if mass doesn't match velocities used) A1 for <math>\pm m(-u - 3u)</math> or <math>\pm 4m(-u - (-2u))</math> A1 for <math>4mu</math> cao ( <math>-4mu</math> is A0) Allow change of sign at end to obtain magnitude.</p>		

Leave  
blank



A diagram showing a block of mass  $50\text{ kg}$  on an inclined plane. The angle of the incline with the horizontal is  $\alpha$ . A vertical line is drawn from the top of the block, perpendicular to the incline.

### Figure 1

A block of mass 50 kg lies on a rough plane which is inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{7}{24}$ . The block is held at rest by a vertical rope, as shown in

Figure 1, and is on the point of sliding down the plane. The block is modelled as a particle and the rope is modelled as a light inextensible string. Given that the friction force acting on the block has magnitude 65.8 N, find

- (a) the tension in the rope, (4)
- (b) the coefficient of friction between the block and the plane. (4)



<p><b>2. (a)</b></p>    <p><b>(b)</b></p>	$T \sin \alpha + 65.8 = 50g \sin \alpha$ $T = 255 \text{ N or } 260 \text{ N}$  $65.8 \cos \alpha = R \sin \alpha$ $\mu = 65.8/R = \tan \alpha = 7/24, 0.29 \text{ or better}$	<p>M1 A1</p> <p><b>DM1A1</b> (4)</p> <hr/> <p>M1 A1</p> <p>M1 A1 (4)</p> <p style="text-align: right;"><b>8</b></p>
<div style="text-align: center;"> <p><b><u>NOTES</u></b></p> <p><b><u>Question 2(a)</u></b>  First M1 for resolving parallel to the plane (or an equation in <math>T</math> only)  First A1 for a correct equation.  Second M1 dependent for producing a value for <math>T</math>.  Second A1 for 255 (N) or 260 (N).</p> <p><b><u>Question 2(b)</u></b>  First M1for <b>any</b> equation containing <math>R</math>.  First A1 for a correct equation. (If equation includes a <math>T</math> term, they must be using a correct value of <math>T</math> to score this mark)  Second M1 for (65.8/their <math>R</math>).  Second A1 for 7/24, 0.29 or better.</p> </div>		

This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire width, providing a guide for handwriting or typing. The paper itself is a clean, off-white color.

3.(a)	$\tan \alpha = 1/3 \Rightarrow \alpha = 18.4^\circ$	M1 A1
(b)	Bearing is $288^\circ$ (nearest degree)	A1 (3)
(c)	$\mathbf{r} = (21\mathbf{i} + 5\mathbf{j}) + t(-6\mathbf{i} + 2\mathbf{j})$	B1 (1)
	$21 - 6t = -(5 + 2t)$	M1 A1
	$t = 6.5$	A1 (3)
		<b>7</b>

**NOTES****Question 3(a)**

First M1 for  $\arctan\left(\frac{\pm 2}{\pm 6}\right)$

First A1 for a correct value from their expression, usually  $18.4^\circ$  or  $71.6^\circ$

Second A1 for  $288$  (nearest degree)

**Question 3(b)**

B1 for  $(21\mathbf{i} + 5\mathbf{j}) + t(-6\mathbf{i} + 2\mathbf{j})$

**Question 3(c)**

M1 for equating the negative of their  $\mathbf{i}$ -component to their  $\mathbf{j}$ -component  
oe

Allow equating the components for the M mark.

First A1 for a correct equation.

Second A1 for  $t = 6.5$

4. The points  $P$  and  $Q$  are at the same height  $h$  metres above horizontal ground. A small stone is dropped from rest from  $P$ . Half a second later a second small stone is thrown vertically downwards from  $Q$  with speed  $7.35 \text{ m s}^{-1}$ . Given that the stones hit the ground at the same time, find the value of  $h$ .

(7)





[illegible]

5.




Figure 2

A particle  $P$  of mass  $2\text{ kg}$  is pushed up a line of greatest slope of a rough plane by a horizontal force of magnitude  $X$  newtons, as shown in Figure 2. The force acts in the vertical plane which contains  $P$  and a line of greatest slope of the plane. The plane is

inclined to the horizontal at an angle  $\alpha$ , where  $\tan \alpha = \frac{3}{4}$

The coefficient of friction between  $P$  and the plane is  $0.5$

Given that the acceleration of  $P$  is  $1.45\text{ m s}^{-2}$ , find the value of  $X$ .

**(10)**

5.	$R = X \sin \alpha + 2g \cos \alpha$ $X \cos \alpha - F - 2g \sin \alpha = 2 \times 1.45$ $F = 0.5R$ <p>Eliminating <math>R</math>: solving for <math>X</math></p> $X = 45$	M1 A2 M1 A2 B1 DM1;DM1 A1 <b>10</b>
	<p style="text-align: center;"><b><u>NOTES</u></b></p> <p><b><u>Question 5</u></b>  First M1 for resolving perp to the plane..  First A2 for a correct equation; -1 each error.  Second M1 for resolving parallel to the plane.  Second A2 for a correct equation; -1 each error. (Allow <math>F</math> at this stage)  B1 for <math>F = \frac{1}{2} R</math>  Third M1 dependent on previous two M's for eliminating <math>R</math>.  Fourth M1 dependent on previous M for solving for <math>X</math>  Third A1 for <math>X = 45</math>.</p>	

6. A uniform rod  $AC$ , of weight  $W$  and length  $3l$ , rests horizontally on two supports, one at  $A$  and one at  $B$ , where  $AB = 2l$ . A particle of weight  $2W$  is placed on the rod at a distance  $x$  from  $A$ . The rod remains horizontal and in equilibrium.

(5)

(5)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


<p><b>6. (a)</b></p>	<p><math>x</math> is greatest when rod is about to tip about <math>B</math> <i>i.e.</i> <math>R_A = 0</math> (can be implied)</p> $M(B), 2W(x - 2l) = W \frac{1}{2}l$ $x = 2.25l$	<p>B1</p> <p>M1 A1</p> <p><b>DM1 A1 (5)</b></p>
<p><b>(b)</b></p>	<p>Use of <math>R_A = 2W</math> in an equation</p> $M(B), 2W(2l - x) + W \frac{1}{2}l = 2W \cdot 2l$ $x = 0.25l$	<p>M1</p> <p>M1 A1 A1</p> <p>A1 (5)</p> <p><b>10</b></p>
	<p style="text-align: center;"><b><u>NOTES</u></b></p> <p><b><u>Question 6(a)</u></b>            B1 for <math>x</math> greatest when <math>R_A = 0</math> (usually implied in moments equation) or correct use of <math>R_A \geq 0</math>.            First M1 for an equation in <math>x</math> and <math>l</math> ONLY (usually moments about <math>B</math> but could come from two equations). Allow if there is <math>W</math> (uncancelled) in each term. (M0 if <math>R_A</math> term included unless it subsequently becomes zero)            First A1 for a correct equation –again allow even if <math>W</math> has not been cancelled.            Second M1, dependent on previous M, for solving for <math>x</math> in terms of <math>l</math>.            Second A1 for <math>x = 2.25l</math>.  <b>N.B.</b> If '<math>l</math>' omitted consistently and then inserted at end award full marks.            If not inserted then can score max B1M1A0M1A0</p> <p><b><u>Question 6(b)</u> Scheme change</b>            First M1 for use of <math>R_A = 2W</math> in any equation (vertical resolution or moments) or for correct use of <math>R_A \leq 2W</math>.            Second M1 for an equation in <math>x</math> and <math>l</math> ONLY (usually moments about <math>B</math> but could come from two equations). Allow if there is <math>W</math> (uncancelled) in each term.            A2 for the equation, again allow even if <math>W</math> has not been cancelled, -1 each error.            Third A1 for <math>x = 0.25l</math>.  <b>N.B.</b> If '<math>l</math>' omitted consistently and then inserted at end award full marks.            If not inserted then can score max M1M1A0A0A0.</p>	

7. A train travels along a straight horizontal track between two stations  $A$  and  $B$ . The train starts from rest at  $A$  and moves with constant acceleration until it reaches its maximum speed of  $108 \text{ km h}^{-1}$ . The train then travels at this speed before it moves with constant deceleration coming to rest at  $B$ . The journey from  $A$  to  $B$  takes 8 minutes.

- (2)


- (2)

(6)

<p>7 (a)</p> <p>(b)</p> <p>(c)</p>	<div style="text-align: right;"><math>108 \times 1000 / 3600 = 30 \text{ m s}^{-1}</math></div>  <div style="text-align: center;"><math>12000 = \frac{1}{2} \times 30(480 + 480 - 4T)</math></div> <div style="text-align: center;"><math>T = 40</math></div> <div style="text-align: center;"><math>a = 30/40 = 0.75 \text{ m s}^{-2}</math></div>	<div>M1 A1 (2)</div> <div>B1 shape</div> <div>DB1 ft figs (2)</div> <div>M1 A2</div> <div>A1</div> <div>M1 A1 (6)</div> <div>10</div>
	<p style="text-align: center;"><b><u>NOTES</u></b></p> <p><b><u>Question 7(a)</u></b> M1 for <math>108 \times 1000 / 3600</math> oe A1 for 30</p> <p><b><u>Question 7(b)</u></b> First B1 for trapezium (B0 for triangle), from the origin, finishing on the <math>t</math>-axis. Second dependent B1 ft on their '30' and 480 or 108 and (8/60 oe).</p> <p><b><u>Question 7(c)</u></b> First M1 for clear attempt at equating <i>total</i> area under a trapezium to distance travelled oe (equation must include at least one '1/2') to give equation in ONE unknown. A2 for a correct equation, -1 each error. <b>N.B.</b> Repeated use of an incorrect <math>v</math> from part (a) is ONE error. Third A1 for <math>T = 40</math> (or 120) <b>N.B.</b> (First M1 <b>only</b> for <math>\frac{1}{2}(480 + x).30 = 12000</math> First A1 for <math>480 - x = 160</math>; Second A1 if they divide 160 in ratio 1:3) (First M0 if they use <math>s</math> = the full distance in any single <i>suvat</i> equation) Second M1 (independent) for a complete method to find <math>a</math>. Fourth A1 for 0.75</p>	

Leave  
blank

**8.**



### Figure 3

A particle  $A$  of mass  $3m$  is held at rest on a rough horizontal table. The particle is attached to one end of a light inextensible string. The string passes over a small smooth pulley  $P$  which is fixed at the edge of the table. The other end of the string is attached to a particle  $B$  of mass  $2m$ , which hangs freely, vertically below  $P$ . The system is released from rest, with the string taut, when  $A$  is  $1.3\text{ m}$  from  $P$  and  $B$  is  $1\text{ m}$  above the horizontal floor, as shown in Figure 3.

Given that  $B$  hits the floor 2 s after release and does not rebound,

- (a) find the acceleration of  $A$  during the first two seconds, (2)
- (b) find the coefficient of friction between  $A$  and the table, (8)
- (c) determine whether  $A$  reaches the pulley. (6)





8 (a)	For B: $1 = \frac{1}{2} a \cdot 2^2 \Rightarrow a = \frac{1}{2} \text{ m s}^{-2}$	M1 A1 (2)
(b)	$R = 3mg; \quad F = \mu R$  $T - F = 3m \times 0.5$  $2mg - T = 2m \times 0.5$  Solving for $\mu$  $\mu = 0.58 \text{ or } 0.582$	B1 ; B1  M1 A1ft  M1 A1 ft  DM1  A1 (8)
		B1 ft  M1  M1  A1  DM1A1 cso (6)  <b>16</b>
	<p style="text-align: center;"><b><u>NOTES</u></b></p> <p><b><u>Question 8(a)</u></b>  First M1 for a complete method to find <math>a</math>. M0 if <math>s = 1.3</math> is used  First A1 for <math>a = 0.5</math></p> <p><b><u>Question 8(b)</u></b>  First B1 for <math>R = 3mg</math>  Second B1 for <math>F = \mu R</math> seen (could be on diagram)  First M1 for resolving horizontally for A (this M mark can be scored if they just use <math>m</math> for mass but M0 if no mass used)  First A1ft on their <math>a</math>, for correct equation. (allow <math>F</math>)  Second M1 for resolving vertically for B (this M mark can be scored if they just use <math>m</math> for mass but M0 if no mass used)  Second A1ft on their <math>a</math>, for correct equation.  (Allow M2A2 for 'whole system' equation but M0 if not using <math>5m</math>)  Third M1 dependent on both previous M marks for solving for <math>\mu</math>  N.B. If <math>m</math> omitted consistently throughout (b), can score max B0B1M1A0M1A0M1A0  <b><u>Question 8(c)</u></b>  B1 ft for (their <math>a \times 2</math>) oe to find <math>v</math>  First M1 for resolving horizontally for A with <math>T = 0</math>  Second M1 for a complete method (must have found a new '<math>a</math>') to find distance moved by A.  First A1 for 0.09 or better (0.087719..)  Third M1, dependent on first and second M marks, for comparison with 0.3 or 1.3 (Must explicitly refer to either 0.3 or 1.3 or an appropriate equivalent)  Second A1 cso for does not reach pulley.</p>	