This resource	was created	and owned b	by Pearson	Edexcel

Surname	Other n	ames
Pearson Edexcel nternational Advanced Level	Centre Number	Candidate Number
Mochanic	с M1	
Advanced/Advance	S IVI I d Subsidiary	
Advanced/Advance Wednesday 25 October 201 Time: 1 hour 30 minutes	S IVI I d Subsidiary 17 – Morning	Paper Reference

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take g = 9.8 m s⁻², and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets – use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over 🕨



P50804A ©2017 Pearson Education Ltd. 1/1/1/



CS M1 WME01

Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

umn Paper	2017 www.mystudybro.com r This resource was created and owned by Pearson Edexcel	Mathemati
1.	A suitcase of mass 40 kg is being dragged in a straight line along a ro floor at constant speed using a thin strap. The strap is inclined at 20° above	ugh horizontal the horizontal.
	The coefficient of friction between the suitcase and the floor is $\frac{3}{2}$. The st	rap is modelled
	4 as a light inextensible string and the suitcase is modelled as a particle. Fin the strap	d the tension in
	the strup.	(7)



General Principles for Mechanics Marking

Question Number	Scheme	Marks	8
1	$T\cos 70^\circ + R = 40g$	M1A1	
	$T\cos 20^{\circ} = F$	M1A1	
	$F = \frac{3}{4}R$	B1	
	Eliminate <i>R</i> and solve for <i>T</i>	DM 1	
	T = 250 N or 246 N	A1	
			7
	Notos		
1	First M1 for resolving vertically with usual rules (must be using either		
1	First M1 for resolving vertically with usual rules (must be using either 20° or 70°) First A1 for a correct equation Second M1 for resolving horizontally with usual rules (must be using either 20° or 70°) Second A1 for a correct equation B1 for $F = \frac{3}{R}$ seen (could be on a diagram)		
	4		
	Third DM1 dependent on previous two M marks		
	Third AT for either 250 (N) or 246 (N)		
2a	$M(D)$, $(1080 \times 1) - (400 \times 2) = R_c \times 3.5$	M1 A1	
	$R_{c} = 80 \text{ (N)}$	A1	
	$M(C)$, $(1080 \times 2.5) + (400 \times 5.5) = R_D \times 3.5$	M1A1	
	$R_{D} = 1400 \text{ (N)}$	A1	(6)
	OR (\uparrow) $R_c + R_p = 1480$	M1A1	
2b	$R_{c} + (R_{c} + 520) = 1480$ OR $R_{p} + (R_{p} - 520) = 1480$	M1 A1	
	$M(D), (1080 \times 1) - 400(x-4) = R_c \times 3.5$	M1 A1	
	x = 2.5	A1	(5)
			11
20	Notes		
2 a	First W11 for a moments equation of a vertical resolution First A1 for a correct equation $(R_C \text{ and/or } R_D do NOT need to be substituted but if one is, it can be their value found from a previous equation)$		

DO NOT WRITE IN THIS ARE

DO NOT WRITE IN THIS ARE



The boy now stands at a point *E* on the girder, where AE = x metres, and the girder remains horizontal and in equilibrium. Given that the magnitude of the reaction on the girder at Dis now 520N greater than the magnitude of the reaction on the girder at C,

(b) find the value of x.



General Principles for Mechanics Marking

Question Number	Scheme	Marks	8
1	$T\cos 70^\circ + R = 40g$	M1A1	
	$T\cos 20^{\circ} = F$	M1A1	
	$F = \frac{3}{4}R$	B1	
	Eliminate <i>R</i> and solve for <i>T</i>	DM 1	
	T = 250 N or 246 N	A1	
			7
	Notos		
1	First M1 for resolving vertically with usual rules (must be using either		
1	First M1 for resolving vertically with usual rules (must be using either 20° or 70°) First A1 for a correct equation Second M1 for resolving horizontally with usual rules (must be using either 20° or 70°) Second A1 for a correct equation B1 for $F = \frac{3}{R}$ seen (could be on a diagram)		
	4		
	Third DM1 dependent on previous two M marks		
	Third AT for either 250 (N) or 246 (N)		
2a	$M(D)$, $(1080 \times 1) - (400 \times 2) = R_c \times 3.5$	M1 A1	
	$R_{c} = 80 \text{ (N)}$	A1	
	$M(C)$, $(1080 \times 2.5) + (400 \times 5.5) = R_D \times 3.5$	M1A1	
	$R_{D} = 1400 \text{ (N)}$	A1	(6)
	OR (\uparrow) $R_c + R_p = 1480$	M1A1	
2b	$R_{c} + (R_{c} + 520) = 1480$ OR $R_{p} + (R_{p} - 520) = 1480$	M1 A1	
	$M(D), (1080 \times 1) - 400(x-4) = R_c \times 3.5$	M1 A1	
	x = 2.5	A1	(5)
			11
20	Notes		
2 a	First W11 for a moments equation of a vertical resolution First A1 for a correct equation $(R_C \text{ and/or } R_D do NOT need to be substituted but if one is, it can be their value found from a previous equation)$		

Question Number	Scheme	Marks
	Second A1 for $R_{a} = 80$ (N)	
	Second M1 for a moments equation or a vertical resolution	
	Third A1 for a correct equation (R_{c} and/or R_{b} do NOT need to be	
	substituted but if one is, it can be their value found from a previous	
	equation)	
	Equation) Fourth A1 for $D = 1400$ (ND)	
	Fourth A1 for $R_D = 1400$ (IN)	
	Enter marks for equations on ePEN, in the order they appear	
	First M1 for a moments equation or a vertical resolution	
2b	First A1 for a correct equation (R_C and/or R_D do NOT need to be	
	substituted but if one is, it can be their value found from a previous	
	equation)	
	Second M1 for a moments equation or a vertical resolution	
	Second A1 for a correct equation (R_C and/or R_D do NOT need to be	
	substituted but if one is, it can be their value found from a previous	
	equation)	
	Third A1 for $x = 2.5$	
	Enter marks for equations on ePEN, in the order they appear	
	N.B. Equations may contain any or all of R_C , R_D or x for M marks but	
	must contain only one of R_C or R_D to earn the A mark.	
	N.B. If they assume that $R_D = 520$, they lose all the marks for part (b).	
	N.B If they start with $2R = 1480$ and then add or subtract (or both) 520	
	to their R value, M0.	
	N.B. If brackets are omitted in a moments equation e.g. $(520 + R_c).4$ is	
	written as $520 + R_c.4$, the M mark can be scored	
3	8 mu - 1 mu - 5 mu	M1A1
Ũ	$\frac{3}{2} \frac{3}{2} \frac{3}$	A1
	$\frac{V - 0.0u}{F_{0} r P} = -I - Am(0.8u - 2u)$	M1 A1
	$\frac{1011.}{1-\pi m(0.0u-2u)}$	
	I = 4.8 mu	AI
	$OD = (0, 0, \dots, 4)$	2.54.4.4
	OR For Q : $I = m(0.8u + 4u)$	MIAI
	I = 4.8 mu	A1
		6
	Notes	
	First M1 for CLM with correct no. of terms, all dimensionally correct, to give	
2	an equation in <i>m</i> , <i>u</i> and their <i>v</i> only. Condone consistent <i>g</i> 's or cancelled <i>m</i> 's	
3	and sign errors.	
	(N.B. The CLM equation could be obtained by equating the magnitudes of the	
	Impulses on each particle)	
	First A1 for a correct equation (they may have $-5mv$)	
	Second A1 for $0.8u$ or $-0.8u$ (as appropriate)	
	Second M1 for using Impulse = Change in Momentum for either P or Q	
	(MU if <i>clearly</i> adding momenta or if g is included or if different mass in the	
	two momentum terms) but condone sign errors.	

Autumn 2017

Р

Pape	This resource was created and owned by Pearson Edexcel	WME01
5.	Two particles P and Q have masses $4m$ and m respectively. They are moving in opposite directions towards each other along the same straight line on a smooth horizontal plane and collide directly. Immediately before the collision the speed of P is $2u$ and the speed of Q is $4u$. In the collision, the particles join together to form a single particle.	Leave blank
	Find, in terms of m and u , the magnitude of the impulse exerted by P on Q in the collision. (6)	
8		
0		

Question Number	Scheme	Marks
	Second A1 for $R_{a} = 80$ (N)	
	Second M1 for a moments equation or a vertical resolution	
	Third A1 for a correct equation (R_{c} and/or R_{b} do NOT need to be	
	substituted but if one is, it can be their value found from a previous	
	equation)	
	Equation) Fourth A1 for $D = 1400$ (ND)	
	Fourth A1 for $R_D = 1400$ (IN)	
	Enter marks for equations on ePEN, in the order they appear	
	First M1 for a moments equation or a vertical resolution	
2b	First A1 for a correct equation (R_C and/or R_D do NOT need to be	
	substituted but if one is, it can be their value found from a previous	
	equation)	
	Second M1 for a moments equation or a vertical resolution	
	Second A1 for a correct equation (R_C and/or R_D do NOT need to be	
	substituted but if one is, it can be their value found from a previous	
	equation)	
	Third A1 for $x = 2.5$	
	Enter marks for equations on ePEN, in the order they appear	
	N.B. Equations may contain any or all of R_C , R_D or x for M marks but	
	must contain only one of R_C or R_D to earn the A mark.	
	N.B. If they assume that $R_D = 520$, they lose all the marks for part (b).	
	N.B If they start with $2R = 1480$ and then add or subtract (or both) 520	
	to their R value, M0.	
	N.B. If brackets are omitted in a moments equation e.g. $(520 + R_c).4$ is	
	written as $520 + R_c.4$, the M mark can be scored	
3	8 mu - 1 mu - 5 mu	M1A1
Ũ	$\frac{3}{2} \frac{3}{2} \frac{3}$	A1
	$\frac{V - 0.0u}{F_{0} r P} = -I - Am(0.8u - 2u)$	M1 A1
	$\frac{1011.}{1-\pi m(0.0u-2u)}$	
	I = 4.8 mu	AI
	$OD = (0, 0, \dots, 4)$	2.54.4.4
	OR For Q : $I = m(0.8u + 4u)$	MIAI
	I = 4.8 mu	A1
		6
	Notes	
	First M1 for CLM with correct no. of terms, all dimensionally correct, to give	
2	an equation in <i>m</i> , <i>u</i> and their <i>v</i> only. Condone consistent <i>g</i> 's or cancelled <i>m</i> 's	
3	and sign errors.	
	(N.B. The CLM equation could be obtained by equating the magnitudes of the	
	Impulses on each particle)	
	First A1 for a correct equation (they may have $-5mv$)	
	Second A1 for $0.8u$ or $-0.8u$ (as appropriate)	
	Second M1 for using Impulse = Change in Momentum for either P or Q	
	(MU if <i>clearly</i> adding momenta or if g is included or if different mass in the	
	two momentum terms) but condone sign errors.	

Question Number	Scheme	Marks
	Third A1 for $4m(0.8u-2u)$ or $-4m(0.8u-2u)$ OR for $m(0.8u+4u)$ or $-m(0.8u+4u)$ Fourth A1 for 4.8mu (must be positive since magnitude)	
4(i)	$ \mathbf{F}_{2} ^{2} = 8^{2} + 14^{2} - 2 \times 8 \times 14 \cos 30$	M1 A1
	Solve for $ \mathbf{F}_2 = 8.1$ (N) or better	M1 A1 (4)
	OR: $\frac{ \mathbf{F}_2 \cos\alpha = 14\cos 30 - 8}{ \mathbf{F}_2 \sin\alpha = 14\sin 30}$	M1 A1
	Solve for $ \mathbf{F}_2 = 8.1$ (N) or better	M1 A1 (4)
4(ii)	$\frac{\sin\theta}{8} = \frac{\sin 30}{8.12467} \text{ or } \frac{\sin\phi}{14} = \frac{\sin 30}{8.12467}$	M1 A1
	Solve: $\theta = 29.49^{\circ}$ or $\phi = 120.51^{\circ}$	M1 A1
	Bearing is 149° (nearest degree)	A1 (5)
	OR: $\frac{ \mathbf{F}_2 \cos\alpha = 14\cos 30 - 8 = 4.124(355.)}{ \mathbf{F}_2 \sin\alpha = 14\sin 30}$	M1 A1
	Solve: $\alpha = 59.49^{\circ}$ Bearing is 149° (nearest degree)	$\begin{array}{c c} M1 & A1 \\ \hline A1 & (5) \end{array}$
		AI (5)
	Notos	
4(i)	First M1 for use of cos rule with 30° First A1 for a correct equation OR: First M1 for 'resolving' in 2 directions with $30^{\circ}/60^{\circ}$ (N.B. M0 here if cos/sin confused) First A1 for TWO correct equations Second M1 for solving for $ \mathbf{F}_2 $, <u>independent</u> but must be solving a 'correct cosine formula but with wrong angle' if using method 1 OR for eliminating α from two equations, <u>independent</u> but equations must have the correct structure if using method 2	
	Second A1 for 8.1 (N) or better	
4(ii)	First M1 for use of sin rule with 30° First A1 for a correct equation (allow 8.12 or better) OR: First M1 for 'resolving' in 2 directions with 30°/60°	

Two forces \mathbf{F}_1 and \mathbf{F}_2 act on a particle. The force \mathbf{F}_1 has magnitude 8N and acts due The resultant of \mathbf{F}_1 and \mathbf{F}_2 is a force of magnitude 14N acting in a direction whose be is 120°.	e east. earing	blank	
Find			
(i) the magnitude of \mathbf{F}_2 ,	(4)		
(ii) the direction of \mathbf{F}_2 , giving your answer as a bearing to the nearest degree.	(5)		

Question Number	Scheme	Marks
	Third A1 for $4m(0.8u-2u)$ or $-4m(0.8u-2u)$ OR for $m(0.8u+4u)$ or $-m(0.8u+4u)$ Fourth A1 for 4.8mu (must be positive since magnitude)	
4(i)	$ \mathbf{F}_{2} ^{2} = 8^{2} + 14^{2} - 2 \times 8 \times 14 \cos 30$	M1 A1
	Solve for $ \mathbf{F}_2 = 8.1$ (N) or better	M1 A1 (4)
	OR: $\frac{ \mathbf{F}_2 \cos\alpha = 14\cos 30 - 8}{ \mathbf{F}_2 \sin\alpha = 14\sin 30}$	M1 A1
	Solve for $ \mathbf{F}_2 = 8.1$ (N) or better	M1 A1 (4)
4(ii)	$\frac{\sin\theta}{8} = \frac{\sin 30}{8.12467} \text{ or } \frac{\sin\phi}{14} = \frac{\sin 30}{8.12467}$	M1 A1
	Solve: $\theta = 29.49^{\circ}$ or $\phi = 120.51^{\circ}$	M1 A1
	Bearing is 149° (nearest degree)	A1 (5)
	OR: $\frac{ \mathbf{F}_2 \cos\alpha = 14\cos 30 - 8 = 4.124(355.)}{ \mathbf{F}_2 \sin\alpha = 14\sin 30}$	M1 A1
	Solve: $\alpha = 59.49^{\circ}$ Bearing is 149° (nearest degree)	$\begin{array}{c c} M1 & A1 \\ \hline A1 & (5) \end{array}$
		AI (5)
	Notos	
4(i)	First M1 for use of cos rule with 30° First A1 for a correct equation OR: First M1 for 'resolving' in 2 directions with $30^{\circ}/60^{\circ}$ (N.B. M0 here if cos/sin confused) First A1 for TWO correct equations Second M1 for solving for $ \mathbf{F}_2 $, <u>independent</u> but must be solving a 'correct cosine formula but with wrong angle' if using method 1 OR for eliminating α from two equations, <u>independent</u> but equations must have the correct structure if using method 2	
	Second A1 for 8.1 (N) or better	
4(ii)	First M1 for use of sin rule with 30° First A1 for a correct equation (allow 8.12 or better) OR: First M1 for 'resolving' in 2 directions with 30°/60°	

Question Number	Scheme	Mark	s
	First A1 for TWO correct equations (allow 4.12 or better) Second M1, independent, for solving a 'correct sine formula' for θ or ϕ		
	OR <u>independent</u> for solving two equations, with correct structure, for α		
	Second A1 for $\theta = AWRT 29^{\circ}$ or $\phi = AWRT 121^{\circ}$		
	OR $\alpha = \text{AWRT 59}^{o}$		
	Third A1 for Bearing is 149° (nearest degree)		
	N.B. First M1A1 Could use cos rule to find an angle		
	N.B. If the resolving method is used and there are no (i) or (ii) labels, only award M1A1 in both cases when an answer is reached.		
5-		M1 A 1	
5a	$\frac{0 = 14.7^2 - 2 \times 9.8h}{1 = 11.025}$	MIAI	
	n = 11.025 may ht = 13.5 or 14 (m)		(4)
	maxin = 15.5 or 14 (m)	AI	(4)
5b	$-1.5 = 14.7t - 4.9t^2$	M1A1	
	$49t^2 - 147t - 15 = 0$		
	$\frac{147 \pm \sqrt{147^2 \pm 6 \times 49}}{147 \pm 100}$		
	$t = \frac{110 - 9100 + 600000}{9.8}$	DM 1	
	t = 3.1 or 3.10 (s)	A1	(4)
			~ /
5c	$v^2 = 14.7^2 + 2 \times (-9.8) \times (-2.5)$	M1 A1	
	$v = 16.3 \text{ or } 16 \text{ (m s}^{-1}\text{)}$	A1	(3)
			11
	Notes		
5a	N.B. If they use $g = 9.81$, lose first A mark (once for whole question)		
	but all other A marks can be scored.		
	suvat equations) condone sign errors		
	First A1 for a correct equation (or equations)		
	Second A1 for $h = 11$ (may be unsimplified) or better (For other		
	methods, give this A1 for any correct (may be unsimplified)		
	intermediate answer)		
5h	Third A1 for 13.5 or 14 (m)		
20	the time up (1.5 s) and then add on the time down. Condone sign errors		
	First A1 for a correct equation or equations		
	Second DM1, dependent, for solving to find required time		
	Second A1 for 3.1 or 3.10 (s)		

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

	,,, _,	Leave
5.	A small ball is projected vertically upwards from a point O with speed 14.7 m s ⁻¹ . point O is 2.5 m above the ground. The motion of the ball is modelled as that of a par moving freely under gravity.	The ticle
	Find	
	(a) the maximum height above the ground reached by the ball,	
		(4)
	(b) the time taken for the ball to first reach a height of 1 m above the ground,	(4)
	(c) the speed of the ball at the instant before it strikes the ground for the first time.	(3)
16	$\begin{array}{ $	

Question Number	Scheme	Mark	s
	First A1 for TWO correct equations (allow 4.12 or better) Second M1, independent, for solving a 'correct sine formula' for θ or ϕ		
	OR <u>independent</u> for solving two equations, with correct structure, for α		
	Second A1 for $\theta = AWRT 29^{\circ}$ or $\phi = AWRT 121^{\circ}$		
	OR $\alpha = \text{AWRT 59}^{o}$		
	Third A1 for Bearing is 149° (nearest degree)		
	N.B. First M1A1 Could use cos rule to find an angle		
	N.B. If the resolving method is used and there are no (i) or (ii) labels, only award M1A1 in both cases when an answer is reached.		
5-		N#1 A 1	
5a	$\frac{0 = 14.7^2 - 2 \times 9.8h}{1 = 11.025}$	MIAI	
	n = 11.025 may ht = 13.5 or 14 (m)		(4)
	maxin = 15.5 or 14 (m)	AI	(4)
5b	$-1.5 = 14.7t - 4.9t^2$	M1A1	
	$49t^2 - 147t - 15 = 0$		
	$\frac{147 \pm \sqrt{147^2 \pm 6 \times 49}}{147 \pm 100}$		
	$t = \frac{110 - 9100 + 600000}{9.8}$	DM 1	
	t = 3.1 or 3.10 (s)	A1	(4)
			~ /
5c	$v^2 = 14.7^2 + 2 \times (-9.8) \times (-2.5)$	M1 A1	
	$v = 16.3 \text{ or } 16 \text{ (m s}^{-1}\text{)}$	A1	(3)
			11
	Notes		
5a	N.B. If they use $g = 9.81$, lose first A mark (once for whole question)		
	but all other A marks can be scored.		
	suvat equations) condone sign errors		
	First A1 for a correct equation (or equations)		
	Second A1 for $h = 11$ (may be unsimplified) or better (For other		
	methods, give this A1 for any correct (may be unsimplified)		
	intermediate answer)		
5h	Third A1 for 13.5 or 14 (m)		
20	the time up (1.5 s) and then add on the time down. Condone sign errors		
	First A1 for a correct equation or equations		
	Second DM1, dependent, for solving to find required time		
	Second A1 for 3.1 or 3.10 (s)		

Question Number	Scheme	Marks
5c	First M1 for a complete method to find the speed / velocity(Could involve two <i>suvat</i> equations) Condone sign errors but must have correct numbers in their equation(s) First A1 for a correct equation (or equations) Second A1 for 16 or 16.3 (m s ⁻¹) Must be <i>positive (speed</i>)	
6а	V 0 270	B1 shape B1 270, V (2)
6b	$\frac{V}{0.6} = \frac{5V}{3}$ Given answer	M1A1 (2)
6с	Time decelerating is 5V	B1
	$\frac{1}{2}V\frac{5V}{3} + (270 - 5V - \frac{5V}{3})V + \frac{1}{2}V.5V = 1500$ OR: $\frac{1}{2}(270 + 270 - 5V - \frac{5V}{3})V = 1500$	M1 A2 DM1A1
	V - 81V + 430 = 0 Given answer	(6)
6d	$V^{2} - 81V + 450 = 0$ (V-6)(V-75) = 0 or $V = \frac{81 \pm \sqrt{81^{2} - 4 \times 450}}{2}$	M1 solving
	V = 6 or 75 $V = 6 \text{ since } (5 \times 75) > 270 \text{ or } V = 75 \text{ unrealistic}$	A1 A1 B1 (4)
		14
	Notes	
<u>6a</u>	First B1 for a trapezium with line starting at the origin Second B1 for 270 and V correctly marked	
6b	M1 for $(t =) \frac{V}{0.6}$; N.B. M1A0 for V=0.6t then answer Must see division or intermediate step from V=0.6t e.g. Changing 0.6 into 3/5. A1 for $t = \frac{5V}{3}$ Given answer	

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

6	An athlata good for a run along a straight harizantal read. Starting from rest sha	Le bl
υ.	An athlete goes for a full along a straight horizontal road. Starting from rest, she accelerates at $0.6 \mathrm{ms^{-2}}$ up to a speed of $V \mathrm{ms^{-1}}$. She then maintains this constant speed of $V \mathrm{ms^{-1}}$ before finally decelerating at $0.2 \mathrm{ms^{-2}}$ back to rest. She covers a total distance of 1500 m in 270 s.	
	(a) Sketch a speed-time graph to represent the athlete's run. (2)	
	(b) Show that she accelerates for $\frac{5V}{-}$ seconds.	
	(2)	
	(c) Show that $V^2 - kV + 450 = 0$, where k is a constant to be found. (6)	
	(d) Find the value of V, justifying your answer. (4)	
20		

Question Number	Scheme	Marks
5c	First M1 for a complete method to find the speed / velocity(Could involve two <i>suvat</i> equations) Condone sign errors but must have correct numbers in their equation(s) First A1 for a correct equation (or equations) Second A1 for 16 or 16.3 (m s ⁻¹) Must be <i>positive (speed</i>)	
6а	V 0 270	B1 shape B1 270, V (2)
6b	$\frac{V}{0.6} = \frac{5V}{3}$ Given answer	M1A1 (2)
6с	Time decelerating is 5V	B1
	$\frac{1}{2}V\frac{5V}{3} + (270 - 5V - \frac{5V}{3})V + \frac{1}{2}V.5V = 1500$ OR: $\frac{1}{2}(270 + 270 - 5V - \frac{5V}{3})V = 1500$	M1 A2 DM1A1
	V - 81V + 430 = 0 Given answer	(6)
6d	$V^{2} - 81V + 450 = 0$ (V-6)(V-75) = 0 or $V = \frac{81 \pm \sqrt{81^{2} - 4 \times 450}}{2}$	M1 solving
	V = 6 or 75 $V = 6 \text{ since } (5 \times 75) > 270 \text{ or } V = 75 \text{ unrealistic}$	A1 A1 B1 (4)
		14
	Notes	
<u>6a</u>	First B1 for a trapezium with line starting at the origin Second B1 for 270 and V correctly marked	
6b	M1 for $(t =) \frac{V}{0.6}$; N.B. M1A0 for V=0.6t then answer Must see division or intermediate step from V=0.6t e.g. Changing 0.6 into 3/5. A1 for $t = \frac{5V}{3}$ Given answer	

Question Number	Scheme	Marks	5
бс	B1 for 5V identified appropriately First M1 for clear attempt to equate the <i>total</i> area under graph to 1500.		
ŬĊ.	(Must include all 3 parts (if not using the trapezium rule) with $\frac{1}{2}$ seen at		
	least once to give equation in V only; may use (1 triangle + 1 trapezium)		
	or (rectangle - trapezium) (May use <i>suvat</i> for one or more, parts of the area)		
	A2 for a correct equation, -1 e.e.o.o.		
	Second DM 1 dependent on first M1 for multiplying out and collecting terms and putting into appropriate form		
	Third A1 for correct equation. Given answer		
6d	N.B. This M1 can be implied by two correct roots but if either answer incorrect		
	then an explicit method must be shown for this M mark. First A1 for $V = 6$		
	Second A1 for $V = 75$		
	B1 on ePEN but treat as DM 1, dependent on both previous A marks, for either reason		
		N 1 A 1	
7 a	$\frac{1 - 3mg\sin\alpha - F}{4m\sigma - T} = 4m\sigma$	MIAI M1A1	(A)
	11112 1 - 11114		(+)
7b	$F = \frac{1}{4}R; R = 3mg\cos\alpha$	B1; M1/	A1
	T-2.4mg=3ma	M1	
	4mg - T = 4ma	A 1	(5)
	$a = \frac{8g}{25}$ Given answer	AI	(5)
7c	Particles have same acceleration	B1	(1)
7d	$v^2 = 2 \times \frac{8g}{35} \times 1.75 (= 0.8g)$	M1 A1	
	$-3mg\sin\alpha - F = 3ma'$	M1	
	a' = -0.8g	A1	
	$0 = 0.8g + 2 \times (-0.8g)s$	M1 A1	(7)
	1 otal distance = 0.5 + 1.75 = 2.25 (m) Accept 2.3 (m)	AI	(/) 17
			1
	Notes		
7a	First M1 for equation of motion for A with usual rules First A1 for a correct equation		
	Second M1 for equation of motion for B with usual rules		
	Second A1 for a correct equation NB If using different tension in second equation M0 for that equation		
	14.D. It using unterent tension in second equation, who for that equation		

12



DO NOT WRITE IN THIS ARE

DO NOT WRITE IN THIS ARE



Figure 2

Figure 2 shows two particles A and B, of masses 3m and 4m respectively, attached to the ends of a light inextensible string. Initially A is held at rest on the surface of a fixed rough inclined plane. The plane is inclined to the horizontal at an angle α where $\tan \alpha = \frac{3}{4}$. The coefficient of friction between A and the plane is $\frac{1}{4}$. The string passes over a small smooth light pulley P which is fixed at the top of the plane. The part of the string from A to P is parallel to a line of greatest slope of the plane. The particle B hangs freely and is vertically below P. The system is released from rest with the string taut and with B at a height of 1.75 m above the ground. In the subsequent motion, A does not hit the pulley.

For the period before B hits the ground,

(a) write down an equation of motion for each particle.

(b) Hence show that the acceleration of *B* is $\frac{8}{35}g$.

(c) Explain how you have used the fact that the string is inextensible in your calculation.

(1)

(5)

(4)

When B hits the ground, B does not rebound and comes immediately to rest.

(d) Find the distance travelled by *A* from the instant when the system is released to the instant when *A* first comes to rest.

(7)

I A MARTINA TANA MANANA MAN

Question Number	Scheme	Marks	5
бс	B1 for 5V identified appropriately First M1 for clear attempt to equate the <i>total</i> area under graph to 1500.		
ŬĊ	(Must include all 3 parts (if not using the trapezium rule) with $\frac{1}{2}$ seen at		
	least once to give equation in V only; may use (1 triangle + 1 trapezium)		
	or (rectangle - trapezium) (May use <i>suvat</i> for one or more, parts of the area)		
	A2 for a correct equation, -1 e.e.o.o.		
	Second DM 1 dependent on first M1 for multiplying out and collecting terms and putting into appropriate form		
	Third A1 for correct equation. Given answer		
6d	N.B. This M1 can be implied by two correct roots but if either answer incorrect		
	then an explicit method must be shown for this M mark. First A1 for $V = 6$		
	Second A1 for $V = 75$		
	B1 on ePEN but treat as DM 1, dependent on both previous A marks, for either reason		
		N 1 A 1	
7 a	$\frac{1 - 3mg\sin\alpha - F}{4m\sigma - T} = 4m\sigma$	MIAI M1A1	(A)
	11112 1 - 11114		(+)
7b	$F = \frac{1}{4}R; R = 3mg\cos\alpha$	B1; M1/	A1
	T-2.4mg=3ma	M1	
	4mg - T = 4ma	A 1	(5)
	$a = \frac{8g}{25}$ Given answer	AI	(5)
7c	Particles have same acceleration	B1	(1)
7d	$v^2 = 2 \times \frac{8g}{35} \times 1.75 (= 0.8g)$	M1 A1	
	$-3mg\sin\alpha - F = 3ma'$	M1	
	a' = -0.8g	A1	
	$0 = 0.8g + 2 \times (-0.8g)s$	M1 A1	(7)
	1 otal distance = 0.5 + 1.75 = 2.25 (m) Accept 2.3 (m)	AI	(/) 17
			1
	Notes		
7a	First M1 for equation of motion for A with usual rules First A1 for a correct equation		
	Second M1 for equation of motion for B with usual rules		
	Second A1 for a correct equation NB If using different tension in second equation M0 for that equation		
	14.D. It using unterent tension in second equation, who for that equation		

12

Question Number	Scheme	Marks
7b	B1 for $F = \frac{1}{4}R$ seen e.g. on diagram First M1 for resolving for A perp to the plane First A1 for correct equation N.B. These first 3 marks can be earned in (a). Second M1 (Hence) for substituting for R and F and trig. and solving for a (must be some evidence of this) their equations of motion from <u>part (a)</u> Second A1 for given answer (Not available if not using exact values for trig ratios)	
7c	B1 for particles have same acceleration (B0 for same velocity or if incorrect extras given)	
7d	First M1 for attempt to find speed (or speed ²) when <i>B</i> hits the ground (M0 if uses <i>g</i>) First A1 for a correct expression Second M1 for attempt to find deceleration of <i>A</i> Second A1 for correct deceleration Third M1 for using deceleration (must have found a deceleration) with $v = 0$ to find distance (M0 if uses <i>g</i>) Third A1 for a correct equation Fourth A1 for 2.25 (m)	