

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	7	/	0	1	Signature	

**Paper Reference(s)**

**6677/01**

# Edexcel GCE

# Mechanics M1

## Advanced/Advanced Subsidiary

Friday 12 January 2007 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--



ND003850623

[illegible]

### Materials required for examination

---

Mathematical Formulae (Green)

### Items included with question papers

Nil

**Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.**

### Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature.

Check that you have the correct question paper.

You must write your answer for each question in the space following the question.

Whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$ .

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

## Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 16 pages in this question paper. Any blank pages are indicated.

## Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.  
©2007 Edexcel Limited.

Printer's Log. No.

N23560A

W850/R6677/57570 3/3/3/3/3/14.400




N 2 3 5 6 0 A 0 1 1 6

*Turn over*

**edexcel**   
advancing learning, changing lives

1.

Figure 1



A particle of weight  $24\text{ N}$  is held in equilibrium by two light inextensible strings. One string is horizontal. The other string is inclined at an angle of  $30^\circ$  to the horizontal, as shown in Figure 1. The tension in the horizontal string is  $Q$  newtons and the tension in the other string is  $P$  newtons. Find

(a) the value of  $P$ ,

(3)

(b) the value of  $Q$ .

(3)




January 2007  
6677 Mechanics M1  
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) <math>P \sin 30^\circ = 24</math> <math>P = 48</math></p> <p>(b) <math>Q = P \cos 30^\circ</math> <math>\approx 41.6</math> accept <math>24\sqrt{3}</math>, awrt 42</p>	<p>M1 A1 A1 <u>3</u></p> <p>M1 A1 A1 <u>3</u> <b>6</b></p>
2.	<p>(a) <math>M(C) \quad 80 \times x = 120 \times 0.5</math> <math>x = 0.75</math> * cso</p> <p>(b) Using reaction at <math>C = 0</math> <math>M(D) \quad 120 \times 0.25 = W \times 1.25</math> ft their <math>x</math> <math>W = 24</math> (N)</p> <p>(c) i <math>X = 24 + 120 = 144</math> (N) ft their <math>W</math></p> <p>(d) The weight of the rock acts precisely at <math>B</math>.</p>	<p>M1 A1 A1 <u>3</u></p> <p>B1 M1 A1 A1 <u>4</u></p> <p>M1 A1ft <u>2</u></p> <p>B1 <u>1</u> <b>10</b></p>
3.	<p>(a) <math>\mathbf{a} = \frac{(15\mathbf{i} - 4\mathbf{j}) - (3\mathbf{i} + 2\mathbf{j})}{4} = 3\mathbf{i} - 1.5\mathbf{j}</math></p> <p>(b) N2L <math>\mathbf{F} = m\mathbf{a} = 6\mathbf{i} - 3\mathbf{j}</math> ft their <math>\mathbf{a}</math> <math> \mathbf{F}  = \sqrt{(6^2 + 3^2)} \approx 6.71</math> (N) accept <math>\sqrt{45}</math>, awrt 6.7</p> <p>(c) <math>\mathbf{v}_6 = (3\mathbf{i} + 2\mathbf{j}) + (3\mathbf{i} - 1.5\mathbf{j})6</math> ft their <math>\mathbf{a}</math> <math>= 21\mathbf{i} - 7\mathbf{j}</math> (<math>\text{ms}^{-1}</math>)</p>	<p>M1 A1 <u>2</u></p> <p>M1 A1 M1 A1 <u>4</u></p> <p>M1 A1ft A1 <u>1</u> <b>9</b></p>

2.

Figure 2



A uniform plank  $AB$  has weight  $120\text{ N}$  and length  $3\text{ m}$ . The plank rests horizontally in equilibrium on two smooth supports  $C$  and  $D$ , where  $AC = 1\text{ m}$  and  $CD = x\text{ m}$ , as shown in Figure 2. The reaction of the support on the plank at  $D$  has magnitude  $80\text{ N}$ . Modelling the plank as a rod,

(a) show that  $x = 0.75$

(3)

A rock is now placed at  $B$  and the plank is on the point of tilting about  $D$ . Modelling the rock as a particle, find

(b) the weight of the rock,

(4)

(c) the magnitude of the reaction of the support on the plank at  $D$ .

(2)

(d) State how you have used the model of the rock as a particle.

(1)



January 2007  
6677 Mechanics M1  
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) <math>P \sin 30^\circ = 24</math> <math>P = 48</math></p> <p>(b) <math>Q = P \cos 30^\circ</math> <math>\approx 41.6</math> accept <math>24\sqrt{3}</math>, awrt 42</p>	<p>M1 A1 A1 <u>3</u></p> <p>M1 A1 A1 <u>3</u> <b>6</b></p>
2.	<p>(a) M(C) <math>80 \times x = 120 \times 0.5</math> <math>x = 0.75</math> * cso</p> <p>(b) Using reaction at C = 0 M(D) <math>120 \times 0.25 = W \times 1.25</math> ft their x <math>W = 24</math> (N)</p> <p>(c) i <math>X = 24 + 120 = 144</math> (N) ft their W</p> <p>(d) The weight of the rock acts precisely at B.</p>	<p>M1 A1 A1 <u>3</u></p> <p>B1 M1 A1 A1 <u>4</u></p> <p>M1 A1ft <u>2</u></p> <p>B1 <u>1</u> <b>10</b></p>
3.	<p>(a) <math>\mathbf{a} = \frac{(15\mathbf{i} - 4\mathbf{j}) - (3\mathbf{i} + 2\mathbf{j})}{4} = 3\mathbf{i} - 1.5\mathbf{j}</math></p> <p>(b) N2L <math>\mathbf{F} = m\mathbf{a} = 6\mathbf{i} - 3\mathbf{j}</math> ft their <math>\mathbf{a}</math> <math> \mathbf{F}  = \sqrt{6^2 + 3^2} \approx 6.71</math> (N) accept <math>\sqrt{45}</math>, awrt 6.7</p> <p>(c) <math>\mathbf{v}_6 = (3\mathbf{i} + 2\mathbf{j}) + (3\mathbf{i} - 1.5\mathbf{j})6</math> ft their <math>\mathbf{a}</math> <math>= 21\mathbf{i} - 7\mathbf{j}</math> (<math>\text{ms}^{-1}</math>)</p>	<p>M1 A1 <u>2</u></p> <p>M1 A1 M1 A1 <u>4</u></p> <p>M1 A1ft A1 <u>1</u> <b>9</b></p>

Leave  
blank

3. A particle  $P$  of mass  $2\text{ kg}$  is moving under the action of a constant force  $\mathbf{F}$  newtons. When  $t = 0$ ,  $P$  has velocity  $(3\mathbf{i} + 2\mathbf{j})\text{ m s}^{-1}$  and at time  $t = 4\text{ s}$ ,  $P$  has velocity  $(15\mathbf{i} - 4\mathbf{j})\text{ m s}^{-1}$ . Find
- (a) the acceleration of  $P$  in terms of  $\mathbf{i}$  and  $\mathbf{j}$ , (2)
- (b) the magnitude of  $\mathbf{F}$ , (4)
- (c) the velocity of  $P$  at time  $t = 6\text{ s}$ . (3)



January 2007  
6677 Mechanics M1  
Mark Scheme

Question Number	Scheme	Marks
1.	<p>(a) <math>P \sin 30^\circ = 24</math> <math>P = 48</math></p> <p>(b) <math>Q = P \cos 30^\circ</math> <math>\approx 41.6</math> accept <math>24\sqrt{3}</math>, awrt 42</p>	<p>M1 A1 A1 <u>3</u></p> <p>M1 A1 A1 <u>3</u> <b>6</b></p>
2.	<p>(a) <math>M(C) \quad 80 \times x = 120 \times 0.5</math> <math>x = 0.75</math> * cso</p> <p>(b) Using reaction at <math>C = 0</math> <math>M(D) \quad 120 \times 0.25 = W \times 1.25</math> ft their <math>x</math> <math>W = 24</math> (N)</p> <p>(c) i <math>X = 24 + 120 = 144</math> (N) ft their <math>W</math></p> <p>(d) The weight of the rock acts precisely at <math>B</math>.</p>	<p>M1 A1 A1 <u>3</u></p> <p>B1 M1 A1 A1 <u>4</u></p> <p>M1 A1ft <u>2</u></p> <p>B1 <u>1</u> <b>10</b></p>
3.	<p>(a) <math>\mathbf{a} = \frac{(15\mathbf{i} - 4\mathbf{j}) - (3\mathbf{i} + 2\mathbf{j})}{4} = 3\mathbf{i} - 1.5\mathbf{j}</math></p> <p>(b) N2L <math>\mathbf{F} = m\mathbf{a} = 6\mathbf{i} - 3\mathbf{j}</math> ft their <math>\mathbf{a}</math> <math> \mathbf{F}  = \sqrt{(6^2 + 3^2)} \approx 6.71</math> (N) accept <math>\sqrt{45}</math>, awrt 6.7</p> <p>(c) <math>\mathbf{v}_6 = (3\mathbf{i} + 2\mathbf{j}) + (3\mathbf{i} - 1.5\mathbf{j})6</math> ft their <math>\mathbf{a}</math> <math>= 21\mathbf{i} - 7\mathbf{j} \text{ (ms}^{-1}\text{)}</math></p>	<p>M1 A1 <u>2</u></p> <p>M1 A1 M1 A1 <u>4</u></p> <p>M1 A1ft A1 <u>1</u> <b>9</b></p>

4. A particle  $P$  of mass  $0.3 \text{ kg}$  is moving with speed  $u \text{ m s}^{-1}$  in a straight line on a smooth horizontal table. The particle  $P$  collides directly with a particle  $Q$  of mass  $0.6 \text{ kg}$ , which is at rest on the table. Immediately after the particles collide,  $P$  has speed  $2 \text{ m s}^{-1}$  and  $Q$  has speed  $5 \text{ m s}^{-1}$ . The direction of motion of  $P$  is reversed by the collision. Find

- (4)

- (2)

(4)





Question Number	Scheme	Marks
4.	<p>(a) CLM <math>0.3u = 0.3 \times (-2) + 0.6 \times 5</math> <math>u = 8</math></p> <p>(b) <math>I = 0.6 \times 5 = 3</math> (Ns)</p> <p>(c) <math>v = u + at \Rightarrow 5 = a \times 1.5</math> (<math>a = \frac{10}{3}</math>) N2L <math>R = 0.6 \times \frac{10}{3} = 2</math></p>	<p>M1 A1 M1 A1 <u>4</u></p> <p>M1 A1 <u>2</u></p> <p>M1 A1 M1 A1 <u>4</u> <b>10</b></p>
5.	<p>(a) <math>v^2 = u^2 + 2as \Rightarrow 0^2 = 21^2 - 2 \times 9.8 \times h</math> <math>h = 22.5</math> (m)</p> <p>(b) <math>v^2 = u^2 + 2as \Rightarrow v^2 = 0^2 + 2 \times 9.8 \times 24</math> or equivalent (= 470.4) <math>v \approx 22</math> (<math>\text{ms}^{-1}</math>) accept 21.7</p> <p>(c) <math>v = u + at \Rightarrow -\sqrt{470.4} = 21 - 9.8t</math> or equivalent – 1 each error <math>t \approx 4.4</math> (s) accept 4.36</p>	<p>M1 A1 A1 <u>3</u></p> <p>M1 A1 A1 <u>3</u></p> <p>M1 A2 (1, 0) A1 <u>4</u> <b>10</b></p>

Leave  
blank


5. A ball is projected vertically upwards with speed  $21 \text{ m s}^{-1}$  from a point  $A$ , which is  $1.5 \text{ m}$  above the ground. After projection, the ball moves freely under gravity until it reaches the ground. Modelling the ball as a particle, find
- (a) the greatest height above  $A$  reached by the ball, (3)
  - (b) the speed of the ball as it reaches the ground, (3)
  - (c) the time between the instant when the ball is projected from  $A$  and the instant when the ball reaches the ground. (4)



Question Number	Scheme	Marks
4.	<p>(a) CLM <math>0.3u = 0.3 \times (-2) + 0.6 \times 5</math> <math>u = 8</math></p> <p>(b) <math>I = 0.6 \times 5 = 3</math> (Ns)</p> <p>(c) <math>v = u + at \Rightarrow 5 = a \times 1.5</math> (<math>a = \frac{10}{3}</math>) N2L <math>R = 0.6 \times \frac{10}{3} = 2</math></p>	<p>M1 A1 M1 A1 <u>4</u></p> <p>M1 A1 <u>2</u></p> <p>M1 A1 M1 A1 <u>4</u> <b>10</b></p>
5.	<p>(a) <math>v^2 = u^2 + 2as \Rightarrow 0^2 = 21^2 - 2 \times 9.8 \times h</math> <math>h = 22.5</math> (m)</p> <p>(b) <math>v^2 = u^2 + 2as \Rightarrow v^2 = 0^2 + 2 \times 9.8 \times 24</math> or equivalent (= 470.4) <math>v \approx 22</math> (<math>\text{ms}^{-1}</math>) accept 21.7</p> <p>(c) <math>v = u + at \Rightarrow -\sqrt{470.4} = 21 - 9.8t</math> or equivalent – 1 each error <math>t \approx 4.4</math> (s) accept 4.36</p>	<p>M1 A1 A1 <u>3</u></p> <p>M1 A1 A1 <u>3</u></p> <p>M1 A2 (1, 0) A1 <u>4</u> <b>10</b></p>

6.

### Figure 3



- (a) Find the value of  $P$ .


(8)

The tension in the rope is now increased to 150 N.

- (b) Find the acceleration of the box.

(6)



Question Number	Scheme	Marks
6.	<p>(a)</p>  <p>Use of <math>F = \mu R</math></p> <p><math>P \cos 20^\circ = \mu R</math></p> <p>i <math>R + P \sin 20^\circ = 30g</math></p> <p><math>P \cos 20^\circ = \mu(30g - P \sin 20^\circ)</math></p> <p><math>P = \frac{0.4 \times 30g}{\cos 20^\circ + 0.4 \sin 20^\circ}</math></p> <p><math>\approx 110 \text{ (N)}</math> accept 109</p> <p>(b) i <math>R + 150 \sin 20^\circ = 30g</math></p> <p><math>(R \approx 242.7)</math></p> <p>N2L <math>\quad \quad \quad 150 \cos 20^\circ - \mu R = 30a</math></p> <p><math>a \approx \frac{150 \cos 20^\circ - 0.4 \times 242.7}{30}</math></p> <p><math>= 1.5 \text{ (ms}^{-2}\text{)}</math> accept 1.46</p>	<p>B1</p> <p>M1 A1</p> <p>M1 A1</p> <p>M1</p> <p>M1</p> <p>A1 <u>8</u></p> <p>M1 A1</p> <p>M1 A1</p> <p>M1</p> <p>A1 <u>6</u> <b>14</b></p>

Leave  
blank

7.

Figure 4




Figure 4 shows two particles  $P$  and  $Q$ , of mass  $3\text{ kg}$  and  $2\text{ kg}$  respectively, connected by a light inextensible string. Initially  $P$  is held at rest on a fixed smooth plane inclined at  $30^\circ$  to the horizontal. The string passes over a small smooth light pulley  $A$  fixed at the top of the plane. The part of the string from  $P$  to  $A$  is parallel to a line of greatest slope of the plane. The particle  $Q$  hangs freely below  $A$ . The system is released from rest with the string taut.

(a) Write down an equation of motion for  $P$  and an equation of motion for  $Q$ . (4)

(b) Hence show that the acceleration of  $Q$  is  $0.98\text{ m s}^{-2}$ . (2)

(c) Find the tension in the string. (2)

(d) State where in your calculations you have used the information that the string is inextensible. (1)

On release,  $Q$  is at a height of  $0.8\text{ m}$  above the ground. When  $Q$  reaches the ground, it is brought to rest immediately by the impact with the ground and does not rebound. The initial distance of  $P$  from  $A$  is such that in the subsequent motion  $P$  does not reach  $A$ . Find

(e) the speed of  $Q$  as it reaches the ground, (2)

(f) the time between the instant when  $Q$  reaches the ground and the instant when the string becomes taut again. (5)

Question Number	Scheme	Marks
7.	<p>(a) N2L <math>Q</math> <math>2g - T = 2a</math>  N2L <math>P</math> <math>T - 3g \sin 30^\circ = 3a</math></p> <p>(b) <math>2g - 3g \sin 30^\circ = 5a</math>  <math>a = 0.98 \text{ (ms}^{-2}\text{)} \star</math> cso</p> <p>(c) <math>T = 2(g - a)</math> or equivalent  <math>\approx 18 \text{ (N)}</math> accept 17.6</p> <p>(d) The (magnitudes of the) accelerations of <math>P</math> and <math>Q</math> are equal</p> <p>(e) <math>v^2 = u^2 + 2as \Rightarrow v^2 = 2 \times 0.98 \times 0.8 \text{ (= 1.568)}</math>  <math>v \approx 1.3 \text{ (ms}^{-1}\text{)}</math> accept 1.25</p> <p>(f) N2L for <math>P</math> <math>-3g \sin 30^\circ = 3a</math>  <math>a = (-)\frac{1}{2}g</math>  <math>s = ut + \frac{1}{2}at^2 \Rightarrow 0 = \sqrt{1.568}t - \frac{1}{2}4.9t^2</math> or equivalent  <math>t = 0.51 \text{ (s)}</math> accept 0.511</p> <p><i>A maximum of one mark can be lost for giving too great accuracy.</i></p>	<p>M1 A1  M1 A1 <u>4</u></p> <p>M1  A1 <u>2</u></p> <p>M1  A1 <u>2</u></p> <p>B1 <u>1</u></p> <p>M1  A1 <u>2</u></p> <p>M1 A1  M1 A1  A1 <u>5</u> <b>16</b></p>