

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	7	/	0	1	Signature	

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Monday 24 May 2010 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with
Edexcel Limited copyright policy.
©2010 Edexcel Limited

Printer's Log. No. _____

Printer's Log. No.
N35390A

W850/R6677/57570 4/5/3



Turn over


edexcel 
advancing learning, changing lives

1. A particle P is moving with constant velocity $(-3\mathbf{i} + 2\mathbf{j}) \text{ m s}^{-1}$. At time $t = 6 \text{ s}$ P is at the point with position vector $(-4\mathbf{i} - 7\mathbf{j}) \text{ m}$. Find the distance of P from the origin at time $t = 2 \text{ s}$.

(5)



Summer 2010
Mechanics M1 6677
Mark Scheme

Question Number	Scheme	Marks
Q1	$(-4\mathbf{i} - 7\mathbf{j}) = \mathbf{r} + 4(-3\mathbf{i} + 2\mathbf{j})$ $\mathbf{r} = (8\mathbf{i} - 15\mathbf{j})$ $ \mathbf{r} = \sqrt{8^2 + (-15)^2} = 17 \text{ m}$	M1 A1 A1 M1 A1 ft [5]
Q2 (a)	 $4mu - 3mku = -2mu + 3mk \frac{u}{2}$ $k = \frac{4}{3}$	M1 A1 M1 A1cso (4)
(b)	For P, $I = m(2u - 4u) = 6mu$ OR For Q, $I = 3m(\frac{ku}{2} - ku)$	M1 A1 A1 (3) (M1A1) [7]
Q3	$(\rightarrow) 100\cos 30 = F$ $F = 0.5 R \text{ seen}$ $(\downarrow) mg + 100\cos 60 = R$ $m = 13 \text{ kg or } 12.6 \text{ kg}$	M1 A1 A1 (B1) M1 A1 DM1 A1 [7]

2. Particle P has mass m kg and particle Q has mass $3m$ kg. The particles are moving in opposite directions along a smooth horizontal plane when they collide directly. Immediately before the collision P has speed $4u$ m s⁻¹ and Q has speed ku m s⁻¹, where k is a constant. As a result of the collision the direction of motion of each particle is reversed and the speed of each particle is halved.

(a) Find the value of k .


(4)

(b) Find, in terms of m and u , the magnitude of the impulse exerted on P by Q .

(3)

This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Summer 2010
Mechanics M1 6677
Mark Scheme

Question Number	Scheme	Marks
Q1	$(-4\mathbf{i} - 7\mathbf{j}) = \mathbf{r} + 4(-3\mathbf{i} + 2\mathbf{j})$ $\mathbf{r} = (8\mathbf{i} - 15\mathbf{j})$ $ \mathbf{r} = \sqrt{8^2 + (-15)^2} = 17 \text{ m}$	M1 A1 A1 M1 A1 ft [5]
Q2 (a)	 $4mu - 3mku = -2mu + 3mk \frac{u}{2}$ $k = \frac{4}{3}$	M1 A1 M1 A1cso (4)
(b)	For P, $I = m(2u - 4u) = 6mu$ OR For Q, $I = 3m(\frac{ku}{2} - ku)$	M1 A1 A1 (3) (M1A1) [7]
Q3	$(\rightarrow) 100\cos 30 = F$ $F = 0.5 R \text{ seen}$ $(\downarrow) mg + 100\cos 60 = R$ $m = 13 \text{ kg or } 12.6 \text{ kg}$	M1 A1 A1 (B1) M1 A1 DM1 A1 [7]

Leave
blank



A diagram showing a block on a horizontal surface. A force of 100 N is applied to the block at an angle of 30° below the horizontal.

Figure 1


A small box is pushed along a floor. The floor is modelled as a rough horizontal plane and the box is modelled as a particle. The coefficient of friction between the box and the floor is $\frac{1}{2}$. The box is pushed by a force of magnitude 100 N which acts at an angle of 30° with the floor, as shown in Figure 1.

Given that the box moves with constant speed, find the mass of the box.

(7)





Summer 2010
Mechanics M1 6677
Mark Scheme

Question Number	Scheme	Marks
Q1	$(-4\mathbf{i} - 7\mathbf{j}) = \mathbf{r} + 4(-3\mathbf{i} + 2\mathbf{j})$ $\mathbf{r} = (8\mathbf{i} - 15\mathbf{j})$ $ \mathbf{r} = \sqrt{8^2 + (-15)^2} = 17 \text{ m}$	M1 A1 A1 M1 A1 ft [5]
Q2 (a)	 $4mu - 3mku = -2mu + 3mk \frac{u}{2}$ $k = \frac{4}{3}$	M1 A1 M1 A1cso (4)
(b)	For P, $I = m(2u - 4u) = 6mu$ OR For Q, $I = 3m(\frac{ku}{2} - ku)$	M1 A1 A1 (3) (M1A1) [7]
Q3	$(\rightarrow) 100\cos 30 = F$ $F = 0.5 R \text{ seen}$ $(\downarrow) mg + 100\cos 60 = R$ $m = 13 \text{ kg or } 12.6 \text{ kg}$	M1 A1 A1 (B1) M1 A1 DM1 A1 [7]

4. A beam AB has length 6 m and weight 200 N. The beam rests in a horizontal position on two supports at the points C and D , where $AC = 1$ m and $DB = 1$ m. Two children, Sophie and Tom, each of weight 500 N, stand on the beam with Sophie standing twice as far from the end B as Tom. The beam remains horizontal and in equilibrium and the magnitude of the reaction at D is three times the magnitude of the reaction at C . By modelling the beam as a uniform rod and the two children as particles, find how far Tom is standing from the end B .

(7)





Question Number	Scheme	Marks
Q4	 <p> $M(B),$ $500x + 500.2x + 200x3 = Rx5 + Sx1$ (or any valid moments equation) $(\downarrow) R + S = 500 + 500 + 200 = 1200$ (or a moments equation) solving for $x; x = 1.2$ m </p>	<p>M1 A1 A1</p> <p>M1 A1</p> <p>M1 A1 cso</p> <p>[7]</p>
Q5 (a)	 <p>Shape (both) Cross Meet on t-axis Figures 25,20,T,25</p>	<p>B1 B1 B1 B1</p> <p>(4)</p>
(b)	<p>For Q: $20\left(\frac{t+25}{2}\right) = 800$ $t = 55$</p> <p>For P: $25\left(\frac{T+55}{2}\right) = 800$ solving for $T: T = 9$</p>	<p>M1 A1</p> <p>DM1 A1</p> <p>M1 A1</p> <p>DM1 A1 (8)</p> <p>[12]</p>

Leave
blank

-
- This image shows a full page of blank, lined paper. It features approximately 20 evenly spaced horizontal grey lines across its entire width, providing a template for handwriting practice or general note-taking. The margins are consistent on all sides.



Question Number	Scheme	Marks
Q4	 <p> $M(B),$ $500x + 500.2x + 200x3 = Rx5 + Sx1$ (or any valid moments equation) $(\downarrow) R + S = 500 + 500 + 200 = 1200$ (or a moments equation) solving for $x; x = 1.2$ m </p>	<p>M1 A1 A1</p> <p>M1 A1</p> <p>M1 A1 cso</p> <p>[7]</p>
Q5 (a)	 <p>Shape (both) Cross Meet on t-axis Figures 25,20,T,25</p>	<p>B1 B1 B1 B1</p> <p>(4)</p>
(b)	<p>For $Q: 20\left(\frac{t+25}{2}\right) = 800$ $t = 55$</p> <p>For $P: 25\left(\frac{T+55}{2}\right) = 800$ solving for $T: T = 9$</p>	<p>M1 A1</p> <p>DM1 A1</p> <p>M1 A1</p> <p>DM1 A1 (8)</p> <p>[12]</p>

6. A ball is projected vertically upwards with a speed of 14.7 m s^{-1} from a point which is 49 m above horizontal ground. Modelling the ball as a particle moving freely under gravity, find

- (a) the greatest height, above the ground, reached by the ball, (4)
- (b) the speed with which the ball first strikes the ground, (3)
- (c) the total time from when the ball is projected to when it first strikes the ground. (3)

[illegible]

Question Number	Scheme	Marks
Q6 (a)	$(\uparrow)v^2 = u^2 + 2as$ $0 = 14.7^2 - 2 \times 9.8 \times s$ $s = 11.025$ (or 11 or 11.0 or 11.03) m Height is 60 m or 60.0 m ft	M1A1 A1 A1ft (4)
(b)	$(\downarrow)v^2 = u^2 + 2as$ $v^2 = (-14.7)^2 + 2 \times 9.8 \times 49$ $v = 34.3$ or 34 m s^{-1}	M1 A1 A1 (3)
(c)	$(\downarrow)v = u + at$ OR $(\downarrow)s = ut + \frac{1}{2}at^2$ $34.3 = -14.7 + 9.8t$ $49 = -14.7t + 4.9t^2$ $t = 5$ $t = 5$	M1 A1 A1 (3) [10]
Q7 (a)	$F = \frac{1}{3}R$ $(\uparrow) R \cos \alpha - F \sin \alpha = 0.4g$ $R = \frac{2}{3}g = 6.53$ or 6.5	B1 M1 A1 M1 A1 (5)
(b)	$(\rightarrow)P - F \cos \alpha - R \sin \alpha = 0$ $P = \frac{26}{45}g = 5.66$ or 5.7	M1 A2 M1 A1 (5) [10]

7.




Figure 2

The coefficient of friction between the particle and the plane is $\frac{1}{3}$.

Given that the particle is on the point of sliding up the plane, find


- (a) the magnitude of the normal reaction between the particle and the plane, (5)

- (b) the value of P .
- (5)

[illegible]

Question Number	Scheme	Marks
Q6 (a)	$(\uparrow)v^2 = u^2 + 2as$ $0 = 14.7^2 - 2 \times 9.8 \times s$ $s = 11.025$ (or 11 or 11.0 or 11.03) m Height is 60 m or 60.0 m ft	M1A1 A1 A1ft (4)
(b)	$(\downarrow)v^2 = u^2 + 2as$ $v^2 = (-14.7)^2 + 2 \times 9.8 \times 49$ $v = 34.3$ or 34 m s^{-1}	M1 A1 A1 (3)
(c)	$(\downarrow)v = u + at$ OR $(\downarrow)s = ut + \frac{1}{2}at^2$ $34.3 = -14.7 + 9.8t$ $49 = -14.7t + 4.9t^2$ $t = 5$ $t = 5$	M1 A1 A1 (3) [10]
Q7 (a)	$F = \frac{1}{3}R$ $(\uparrow) R \cos \alpha - F \sin \alpha = 0.4g$ $R = \frac{2}{3}g = 6.53$ or 6.5	B1 M1 A1 M1 A1 (5)
(b)	$(\rightarrow)P - F \cos \alpha - R \sin \alpha = 0$ $P = \frac{26}{45}g = 5.66$ or 5.7	M1 A2 M1 A1 (5) [10]

8.

**Figure 3**

Two particles A and B have mass 0.4 kg and 0.3 kg respectively. The particles are attached to the ends of a light inextensible string. The string passes over a small smooth pulley which is fixed above a horizontal floor. Both particles are held, with the string taut, at a height of 1 m above the floor, as shown in Figure 3. The particles are released from rest and in the subsequent motion B does not reach the pulley.

(a) Find the tension in the string immediately after the particles are released. (6)

(b) Find the acceleration of A immediately after the particles are released. (2)

When the particles have been moving for 0.5 s , the string breaks.

(c) Find the further time that elapses until B hits the floor. (9)



Question Number	Scheme	Marks
Q8 (a) Mark together	$(\downarrow)0.4g - T = 0.4a$ $(\uparrow)T - 0.3g = 0.3a$ solving for T $T = 3.36 \text{ or } 3.4 \text{ or } 12g/35 \text{ (N)}$	M1 A1 M1 A1 DM1 A1 (6)
(b)	$0.4g - 0.3g = 0.7a$ $a = 1.4 \text{ m s}^{-2}, g/7$	DM1 A1 (2)
(c)	$(\uparrow)v = u + at$ $v = 0.5 \times 1.4$ $= 0.7$ $(\uparrow)s = ut + \frac{1}{2}at^2$ $s = 0.5 \times 1.4 \times 0.5^2$ $= 0.175$ $(\downarrow)s = ut + \frac{1}{2}at^2$ $1.175 = -0.7t + 4.9t^2$ $4.9t^2 - 0.7t - 1.175 = 0$ $t = \frac{0.7 \pm \sqrt{0.7^2 + 19.6 \times 1.175}}{9.8}$ $= 0.5663 \text{..or } - \dots$ Ans 0.57 or 0.566 s	M1 A1 ft on a M1 A1 ft on a DM1 A1 ft DM1 A1 cao A1 cao (9) [17]