

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	7	/	0	1	Signature	

Paper Reference(s)

6677/01

Edexcel GCE

Mechanics M1

Advanced/Advanced Subsidiary

Monday 13 May 2013 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2013 Pearson Education Ltd

Printer's Log. No. _____

Printer's Log. No.
P41828A

W850/R6677/57570 5/5/5/5/



Turn over

PEARSON

Leave
blank



Question Number	Scheme	Marks
1.		
(a)	For P , $-I = 3(1 - 4)$	M1 A1
	$I = 9 \text{ Ns}$	A1
		(3)
(b)	For Q , $9 = m(1.5 - -3)$	M1 A1
	$m = 2$	A1
	OR	
	$12 - 3m = 3 + 1.5m$	M1 A1
	$m = 2$	A1
		(3)
		[6]
Notes for Question 1		
Q1(a)	<p>M1 for attempt at Impulse = difference in momenta for particle P, (must be considering <i>one</i> particle i.e. have <i>same mass</i> in both terms) (M0 if g is included or if mass omitted).</p> <p>First A1 for $\pm 3(1 - 4)$</p> <p>Second A1 for 9 (Must be positive). Allow change of sign at end to obtain magnitude.</p> <p>N.B. For M1 they may use CLM to find a value for m first and then use it when considering the change in momentum of Q to find the impulse.</p>	
Q1(b)	<p>EITHER</p> <p>M1 for attempt at: their Impulse from (a) = difference in momenta for particle Q, (must be considering <i>one</i> particle) (M0 if g is included or if mass omitted).</p> <p>First A1 for $9 = m(1.5 - -3)$ oe.</p> <p>Second A1 for $m = 2$.</p> <p>OR</p> <p>M1 for attempt at CLM equation, with correct no. of terms, dimensionally correct. Allow consistent extra g's and sign errors.</p> <p>First A1 for a correct equation i.e. $12 - 3m = 3 + 1.5m$ oe.</p> <p>Second A1 for $m = 2$.</p>	

Leave
blank

(3)



Question Number	Scheme	Marks
2.		
(a)	For system, $(\uparrow), T - 950g - 50g = 1000 \times -2$	M1 A1
	$T = 7800 \text{ N}$	A1
		(3)
(b)	For woman, $(\uparrow), R - 50g = 50 \times -2$	M1 A1
	$R = 390 \text{ N}$	A1
		(3)
		[6]
Notes for Question 2		
Q2(a)	(In both parts, use the <i>mass</i> to decide which part of the system is being considered and M marks can only be scored if an equation contains only forces acting on that part of the system) M1 is for a complete method for finding T i.e. for an equation in T only, dimensionally correct, with the correct number of terms. First A1 for a correct equation. Second A1 for 7800 (N).	
Q2(b)	M1 is for a complete method for finding R i.e. for an equation in R only, dimensionally correct, with the correct number of terms. First A1 for a correct equation. Second A1 for 390 (N). N.B. Equation for lift <i>only</i> is: $T - 950g - R = 950 \times (-2)$	

3.




Figure 1

A box of mass 2 kg is held in equilibrium on a fixed rough inclined plane by a rope. The rope lies in a vertical plane containing a line of greatest slope of the inclined plane. The rope is inclined to the plane at an angle α , where $\tan \alpha = \frac{3}{4}$, and the plane is at an angle of 30° to the horizontal, as shown in Figure 1. The coefficient of friction between the box and the inclined plane is $\frac{1}{3}$ and the box is on the point of slipping up the plane. By modelling the box as a particle and the rope as a light inextensible string, find the tension in the rope.

(8)



Question Number	Scheme	Marks
3.	$T \cos \alpha - F = 2g \cos 60^\circ$	M1 A1
	$T \sin \alpha + R = 2g \cos 30^\circ$	M1 A1
	$F = \frac{1}{3} R$	B1
	eliminating F and R	DM1
	$T = g(1 + \frac{1}{\sqrt{3}})$, 1.6g (or better), 15.5, 15 (N)	DM1 A1
		(8)
		[8]
Notes for Question 3		
Q3	<p>First M1 for resolving parallel to the plane with correct no. of terms and both T and $2g$ terms resolved.</p> <p>First A1 for a correct equation. (use of α instead of 30° or 60° or vice versa is an A error not M error; similarly if they use $\sin(3/5)$ or $\cos(4/5)$ when resolving, this can score M1A0)</p> <p>Second M1 for resolving perpendicular to the plane with correct no. of terms and both T and $2g$ terms resolved.</p> <p>Second A1 for a correct equation (use of α instead of 30° or 60° or vice versa is an A error not M error; similarly if they use $\sin(3/5)$ or $\cos(4/5)$ when resolving, this can score M1A0)</p> <p>B1 for $F = 1/3 R$ seen or implied.</p> <p>Third M1, dependent on first two M marks and appropriate angles used when resolving in <i>both</i> equations, for eliminating F and R.</p> <p>Fourth M1 dependent on third M1, for solving for T</p> <p>Third A1 for 15(N) or 15.5 (N).</p> <p>N.B. The first two M marks can be for two resolutions in any directions. Use of $\tan \alpha = 4/3$ leads to an answer of 17.83...and can score max 7/8.</p>	

Leave
blank

4. A lorry is moving along a straight horizontal road with constant acceleration. The lorry passes a point A with speed $u \text{ m s}^{-1}$, ($u < 34$), and 10 seconds later passes a point B with speed 34 m s^{-1} . Given that $AB = 240 \text{ m}$, find

(a) the value of u ,

(3)

(b) the time taken for the lorry to move from A to the mid-point of AB .

(6)



Question Number	Scheme	Marks
4.		
(a)	$240 = \frac{1}{2}(u + 34)10$	M1 A1
	$u = 14$	A1
		(3)
(b)	$34 = 14 + 10a \Rightarrow a = 2$	M1 A1
	$120 = 14t + \frac{1}{2} \times 2 \times t^2$	M1 A1
	$t^2 + 14t - 120 = 0$	
	Solving, $t = -20$ or 6	DM1
	$t = 6$	A1
	OR	
	$34 = 14 + 10a \Rightarrow a = 2$	M1 A1
	$v^2 = 14^2 + 2 \times 2 \times 120 \Rightarrow v = 26$	
	AND $26 = 14 + 2t$	M1 A1
	$t = 6$	DM1 A1
		(6)
		[9]
Notes for Question 4		
Q4(a)	First M1 for a complete method to produce an equation in u only. First A1 for a correct equation. ($u^2 - 48u + 476 = 0$ oe is possible). Second A1 for $u = 14$.	
Q4(b)	EITHER First M1 for an equation in a only. (M0 if $v = 34$ when $s = 120$ is used) First A1 for $a = 2$. (This may have been found in part (a)) Second M1 for a 3-term quadratic equation in t only, allow sign errors (must have found a value of a . (M0 if $v = 34$ when $s = 120$ is used) Second A1 for a correct equation. Third M1 dependent on previous M1 for solving for t . Third A1 for $t = 6$ OR First M1 for an equation in a only. First A1 for $a = 2$. (This may have been found in part (a)) Second M1 for a complete method to obtain an equation in t only, allow sign errors. (must have found a value of a) Second A1 for a correct equation. Third M1 dependent on previous M1 for solving for t . Third A1 for $t = 6$	

5. A car is travelling along a straight horizontal road. The car takes 120 s to travel between two sets of traffic lights which are 2145 m apart. The car starts from rest at the first set of traffic lights and moves with constant acceleration for 30 s until its speed is 22 m s^{-1} . The car maintains this speed for T seconds. The car then moves with constant deceleration, coming to rest at the second set of traffic lights.

- (a) Sketch, in the space below, a speed-time graph for the motion of the car between the two sets of traffic lights.

(2)

- (b) Find the value of T .

(3)

A motorcycle leaves the first set of traffic lights 10 s after the car has left the first set of traffic lights. The motorcycle moves from rest with constant acceleration, $a \text{ m s}^{-2}$, and passes the car at the point A which is 990 m from the first set of traffic lights. When the motorcycle passes the car, the car is moving with speed 22 m s^{-1} .


- (c) Find the time it takes for the motorcycle to move from the first set of traffic lights to the point A .

(4)

- (d) Find the value of a .

(2)



Question Number	Scheme	Marks
5.		
(a)	Speed 	Shape B1 Figures B1 (2)
(b)	$\frac{(120 + T)22}{2} = 2145$ $T = 75$	M1 A1 A1 (3)
(c)	$\frac{(t + t - 30)22}{2} = 990$ $t = 60$ Answer = $60 - 10 = 50$	M1 A1 A1 A1 (4)
(d)	$990 = 0.5a50^2$ $a = 0.79, 0.792, 99/125$ oe	M1 A1 (2)
		[11]

Notes for Question 5

Q5(a)	First B1 for a trapezium starting at the origin and ending on the t -axis. Second B1 for the figures marked (allow missing 0 and a delineator oe for T) (allow if they have used $T = 75$ correctly on their graph)	
Q5(b)	First M1 for producing an equation in their T only by equating the area of the trapezium to 2145, with the correct no. of terms. If using a single trapezium, we need to see evidence of using $\frac{1}{2}$ the sum of the two parallel sides or if using triangle(s), need to see $\frac{1}{2}$ base x height. Second A1 cao for a correct equation in T (This is not f.t. on their T) Third A1 for $T = 75$. N.B. Use of a single <i>suvat</i> equation for the whole motion of the car e.g. $s = t(u+v)/2$ is M0	
Q5(c)	First M1 for producing an equation in t only (they may use $(t - 30)$ oe as their variable) by equating the area of the trapezium to 990, with the correct no. of terms. If using a trapezium, we need to see evidence of using $\frac{1}{2}$ the sum of the two parallel sides or if using triangle(s), need to see $\frac{1}{2}$ base x height. First A1 for a correct equation. Second A1 for $t = 60$ (Allow $30 + 30$). Third A1 for answer of 50. N.B. Use of a single <i>suvat</i> equation for the whole motion of the car e.g. $s = t(u+v)/2$ is M0. Use of the motion of the motorcycle is M0 (insufficient information). Use of $v = 22$ for the motorcycle is M0.	
Q5(d)	First M1 for an equation in a only. First A1 for $a = 0.79, 0.792, 99/125$ oe N.B. Use of $v = 22$ for the motorcycle is M0.	

6. A beam AB has length 15 m. The beam rests horizontally in equilibrium on two smooth supports at the points P and Q , where $AP = 2$ m and $QB = 3$ m. When a child of mass 50 kg stands on the beam at A , the beam remains in equilibrium and is on the point of tilting about P . When the same child of mass 50 kg stands on the beam at B , the beam remains in equilibrium and is on the point of tilting about Q . The child is modelled as a particle and the beam is modelled as a non-uniform rod.

- (ii) Find the distance of the centre of mass of the beam from A .

When the child stands at the point X on the beam, it remains horizontal and in equilibrium. Given that the reactions at the two supports are equal in magnitude,

- (b) find AX .



Question Number	Scheme	Marks
6.		
(a)	<p>Diagram description: A horizontal beam AB of length 12 m. A vertical force P acts upwards at point A. A vertical force Mg acts downwards at a point x m from A. A vertical force Q acts upwards at point B, which is 3 m from the end B. The distance from A to the point of application of Mg is 2 m.</p>	
	$M(P), \quad 50g \times 2 = Mg \times (x - 2)$	M1 A1
	$M(Q), \quad 50g \times 3 = Mg \times (12 - x)$	M1 A1
(i)	$M = 25 \text{ (kg)}$	DM1 A1
(ii)	$x = 6 \text{ (m)}$	DM1 A1
		(8)
(b)	<p>Diagram description: A horizontal beam AB of length 12 m. A vertical force R acts upwards at point A. A vertical force $25g$ acts downwards at a point 6 m from A. A vertical force $50g$ acts downwards at a point X m from the $25g$ force. A vertical force R acts upwards at point B, which is 3 m from the end B. The distance from A to the point of application of $25g$ is 2 m.</p>	
	$(\uparrow)R + R = 25g + 50g$	M1 A1 ft
	$M(A), \quad 2R + 12R = 25g \times 6 + 50g \times AX$	M1 A1 ft
	$AX = 7.5 \text{ (m)}$	DM1 A1
		(6)
		[14]

Notes for Question 6		
Q6(a)	<p>First M1 for moments about P equation with usual rules (or moments about a different point AND vertical resolution and R then eliminated) (M0 if non-zero reaction at Q)</p> <p>Second M1 for moments about Q equation with usual rules (or moments about a different point AND vertical resolution) (M0 if non-zero reaction at P)</p> <p>Second A1 for a correct equation in M and same unknown.</p> <p>Third M1, dependent on first and second M marks, for solving for M</p> <p>Third A1 for 25 (kg)</p> <p>Fourth M1, dependent on first and second M marks, for solving for x</p> <p>Fourth A1 for 6 (m)</p> <p><u>N.B. No marks available if rod is assumed to be uniform but can score max 5/6 in part (b), provided they have found values for M and x to f.t. on.</u></p> <p>If they have just invented values for M and x in part (a), they can score the M marks in part (b) but <u>not</u> the A marks.</p>	
Q6(b)	<p>First M1 for vertical resolution or a moments equation, with usual rules.</p> <p>First A1 ft on their M and x from part (a), for a correct equation. (must have <i>equal reactions</i> in vertical resolution to earn this mark)</p> <p>Second M1 for a moments equation with usual rules.</p> <p>Second A1 ft on their M and x from part (a), for a correct equation in R and same unknown length.</p> <p>Third M1, dependent on first and second M marks, for solving for AX (<i>not their unknown length</i>) with $AX \leq 15$</p> <p>Third A1 for $AX = 7.5$ (m)</p> <p>N.B. If a single equation is used (see below), equating the sum of the moments of the child and the weight about P to the sum of the moments of the child and the weight about Q, this can score M2 A2 ft on their M and x from part (a), provided the equation is in one unknown. Any method error, loses both M marks.</p> <p>e.g. $25g.4 + 50g(x - 2) = 25g.6 + 50g(12 - x)$ oe.</p>	

7. [In this question, the horizontal unit vectors \mathbf{i} and \mathbf{j} are directed due east and due north respectively.]

$$\mathbf{v} = (1 - 2t)\mathbf{i} + (3t - 3)\mathbf{j}$$

- (a) Find the speed of P when $t = 0$ (3)
- (b) Find the bearing on which P is moving when $t = 2$ (2)
- (c) Find the value of t when P is moving
- (i) parallel to \mathbf{j} ,
- (ii) parallel to $(-\mathbf{i} - 3\mathbf{j})$. (6)



Question Number	Scheme	Marks
7.		
(a)	$t = 0$ gives $\mathbf{v} = \mathbf{i} - 3\mathbf{j}$	B1
	speed = $\sqrt{1^2 + (-3)^2}$	M1
	$= \sqrt{10} = 3.2$ or better	A1
		(3)
(b)	$t = 2$ gives $\mathbf{v} = (-3\mathbf{i} + 3\mathbf{j})$	M1
	Bearing is 315°	A1
		(2)
(c)(i)	$1 - 2t = 0 \Rightarrow t = 0.5$	M1 A1
(ii)	$-(3t - 3) = -3(1 - 2t)$	M1 A1
	Solving for t	DM1
	$t = 2/3, 0.67$ or better	A1
		(6)
		[11]
Notes for Question 7		
Q7(a)	B1 for $\mathbf{i} - 3\mathbf{j}$. M1 for $\sqrt{\text{(sum of squares of cpt.s)}}$ A1 for $\sqrt{10}, 3.2$ or better	
Q7(b)	M1 for clear attempt to sub $t = 2$ into given expression. A1 for 315 .	
Q7(c)	(i) First M1 for $1 - 2t = 0$. First A1 for $t = 0.5$. N.B. If they offer two solutions, by equating both the \mathbf{i} and \mathbf{j} components to zero, give M0. (ii) First M1 for $\frac{1-2t}{3t-3} = \pm\left(\frac{-1}{-3}\right)$ o.e. (Must be an equation in t only) First A1 for a correct equation (the + sign) Second M1, dependent on first M1, for solving for t . Second A1 for $2/3, 0.67$ or better.	

Leave
blank


 $A(2m)$ 

Figure 2

Two particles A and B have masses $2m$ and $3m$ respectively. The particles are attached to the ends of a light inextensible string. Particle A is held at rest on a smooth horizontal table. The string passes over a small smooth pulley which is fixed at the edge of the table. Particle B hangs at rest vertically below the pulley with the string taut, as shown in Figure 2. Particle A is released from rest. Assuming that A has not reached the pulley, find

- (a) the acceleration of B , (5)
- (b) the tension in the string, (1)
- (c) the magnitude and direction of the force exerted on the pulley by the string. (4)



Question Number	Scheme	Marks
8.		
(a)	For A , $T = 2ma$	B1
	For B , $3mg - T = 3ma$	M1 A1
	$3mg = 5ma$	DM1
	$\frac{3g}{5} = a$ (5.9 or 5.88 m s ⁻²)	A1
		(5)
(b)	$T = 6mg/5$; $12m$; $11.8m$	B1
		(1)
(c)	$F = \sqrt{T^2 + T^2}$	M1 A1 ft
	$F = \frac{6mg\sqrt{2}}{5}$; $1.7mg$ (or better); $16.6m$; $17m$	A1
	Direction clearly marked on a diagram, with an arrow, and 45° (oe) marked	B1
		(4)
		[10]
<u>Notes for Question 8</u>		
Q8(a)	<p>B1 for $T = 2ma$</p> <p>First M1 for resolving vertically (up or down) for B, with correct no. of terms. (allow omission of m, provided 3 is there)</p> <p>First A1 for a correct equation.</p> <p>Second M1, dependent on first M1, for eliminating T, to give an equation in a only.</p> <p>Second A1 for 0.6g, 5.88 or 5.9.</p> <p>N.B. 'Whole system' equation: $3mg = 5ma$ earns first 4 marks but any error loses all 4.</p>	
Q8(b)	B1 for $\frac{6mg}{5}$, $11.8m$, $12m$	
Q8(c)	<p>M1 $\sqrt{(T^2 + T^2)}$ or $\frac{T}{\sin 45^\circ}$ or $\frac{T}{\cos 45^\circ}$ or $2T\cos 45^\circ$ or $2T\sin 45^\circ$ (allow if m omitted)</p> <p>(M0 for $T \sin 45^\circ$)</p> <p>First A1 ft on their T.</p> <p>Second A1 cao for $\frac{6mg\sqrt{2}}{5}$ oe, $1.7mg$ (or better), $16.6m$, $17m$</p> <p>B1 for the direction clearly shown on a diagram with an arrow and 45° marked.</p>	