

Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	6	8	/	0	1	Signature	

Paper Reference(s)

6668/01

Edexcel GCE

Further Pure Mathematics FP2

Advanced/Advanced Subsidiary

Friday 6 June 2014 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

[illegible]

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.
©2014 Pearson Education Ltd

Printer's Log. No. _____

Printer's Log. No.
P44512A

W850/R6668/57570 5/5/1/



Turn over

PEARSON

Leave
blank

- (1)

- $$\sum_{r=1}^n \frac{2}{(r+2)(r+4)} = \frac{n(7n+25)}{12(n+3)(n+4)}$$

(5)



Question Number	Scheme		Marks
1.(a)	$\frac{2}{(r+2)(r+4)} = \frac{1}{r+2} - \frac{1}{r+4}$	Correct partial fractions. Can be seen in (b) – give B1 for that.	B1 (1)
(b)	$\sum_{r=1}^n \frac{2}{(r+2)(r+4)} = \sum_{r=1}^n \left(\frac{1}{r+2} - \frac{1}{r+4} \right)$		
	$= \frac{1}{3} - \frac{1}{5} + \frac{1}{4} - \frac{1}{6} + \dots + \frac{1}{n+1} - \frac{1}{n+3} + \frac{1}{n+2} - \frac{1}{n+4}$	Attempts at least the first 2 terms and at least the last 2 terms as shown. (May be implied by later work) Must start at 1 and end at n	M1
	$= \frac{1}{3} + \frac{1}{4} - \frac{1}{n+3} - \frac{1}{n+4}$	M1: Identifies their four fractions that do not cancel. If all terms are positive this mark is lost. A1: Correct four fractions	M1A1
	$= \frac{7}{12} - \frac{1}{n+3} - \frac{1}{n+4}$		
	$= \frac{7(n+3)(n+4) - 12(n+4) - 12(n+3)}{12(n+3)(n+4)}$ $= \frac{7n^2 + 49n + 84 - 12n - 48 - 12n - 36}{12(n+3)(n+4)}$	Attempt to combine at least 3 fractions, 2 of which have a function of n in the denominator and expands the numerator. As a minimum, the product of 2 linear factors must be expanded in the numerator.	M1
	$= \frac{n(7n+25)}{12(n+3)(n+4)} *$	cso Must be factorised. If worked with r instead of n throughout, deduct last mark only.	A1 (5)
			Total 6
(b) Way 2	$\frac{7}{12} - \left(\frac{1}{n+3} + \frac{1}{n+4} \right)$		
	$= \frac{7}{12} - \left(\frac{n+4+n+3}{(n+3)(n+4)} \right)$		
	$= \frac{7(n+3)(n+4) - 24n - 84}{12(n+3)(n+4)}$		
	$= \frac{7n^2 + 49n + 84 - 24n - 84}{12(n+3)(n+4)}$	Attempt to combine at least 3 fractions, 2 of which have a function of n in the denominator and expands the numerator. Min as above	M1
	$= \frac{n(7n+25)}{12(n+3)(n+4)} *$	cso	A1

Leave
blank

2. Use algebra to find the set of values of x for which

$$|3x^2 - 19x + 20| < 2x + 2$$

(6)

Question Number	Scheme		Marks
2.	$ 3x^2 - 19x + 20 < 2x + 2$		
	$3x^2 - 19x + 20 = 2x + 2$ $\Rightarrow 3x^2 - 21x + 18 = 0 \Rightarrow x = ..$	$3x^2 - 19x + 20 = 2x + 2$ and attempt to solve correctly May be solved as an inequality	M1
	$x = 1, \quad x = 6$	Both (ie critical values seen)	A1
	$-(3x^2 - 19x + 20) = 2x + 2$ $\Rightarrow 3x^2 - 17x + 22 = 0 \Rightarrow x = ..$	$-(3x^2 - 19x + 20) = 2x + 2$ and attempt to solve correctly May be solved as an inequality	M1
	$x = \frac{11}{3}, \quad x = 2$	Both (critical values seen) Accept awrt 3.67	A1
	$1 < x < 2, \quad \frac{11}{3} < x < 6$	Must be strict inequalities. Accept awrt 3.67 A1 either correct, A1 both correct. But give A1A0 if both correct apart from \leq seen somewhere in the final answers. Give A1A0 if both correct and extra intervals seen	A1, A1
			(6)
			Total 6

If *no algebra* seen (implies a calculator solution) no marks.

With algebra:

M1 Squaring and reaching a quartic = 0

M1 Attempt to factorise and obtain at least one solution for x . Coefficient of x^4 and constant term correct for their quartic.

A1 Any 2 correct values

A1 All 4 correct values

Final 2 A marks as above

Accept set notation for the final 2 A marks. $x \in (1, 2), \quad x \in (\frac{11}{3}, 6)$ not $[1, 2]$



P 4 4 5 1 2 A 0 6 2 8

Question Number	Scheme		Marks
3.	$y = \sqrt{8 + e^x}$		
	$y = (8 + e^x)^{\frac{1}{2}} \Rightarrow \frac{dy}{dx} = \frac{1}{2}(8 + e^x)^{-\frac{1}{2}} \times e^x$	M1: $\frac{dy}{dx} = k(8 + e^x)^{-\frac{1}{2}} \times e^x$ A1: Correct differentiation	M1A1
	$\frac{d^2y}{dx^2} = \frac{1}{2}(8 + e^x)^{-\frac{1}{2}} \times e^x - \frac{1}{4}(8 + e^x)^{-\frac{3}{2}} \times e^{2x}$	M1: Correct use of the product rule $\frac{dy}{dx} = k(8 + e^x)^{-\frac{1}{2}} \times e^x$ $\pm K(8 + e^x)^{-\frac{3}{2}} \times e^{(2)x}$ A1: Correct second derivative with $e^x \times e^x$ or e^{2x}	M1A1
	$f(0) = 3$	May only appear in the expansion	B1
	$f'(0) = \frac{1}{6}, f''(0) = \frac{17}{108}$	Attempt both $f'(0)$ and $f''(0)$ with their derivatives found above	M1
	$(y=)3 + \frac{1}{6}x + \frac{17}{216}x^2$	M1: Uses the correct Maclaurin series with their values. Accept 2 or 2! in x^2 term A1: Correct expression	M1 A1cso (8)
			Total 8
	Alternative Methods:		
	$e^x = 1 + x + \frac{x^2}{2!} + \dots$	2 or 2!	
	$y = \left(9 + x + \frac{x^2}{2} \dots\right)^{\frac{1}{2}}$	M1: Subst correct expansion	M1
	$= 3 \left(1 + \frac{x}{9} + \frac{x^2}{9 \times 2} + \dots\right)^{\frac{1}{2}}$	B1: for 3 A1: for bracket	B1 A1
	$= 3 \left(1 + \frac{1}{2} \left(\frac{x}{9} + \frac{x^2}{2 \times 9}\right) + \frac{\frac{1}{2} \times (-\frac{1}{2})}{2!} \left(\frac{x}{9} + \frac{x^2}{2 \times 9}\right)^2\right)^{\frac{1}{2}}$	M1: Binomial expansion up to at least the squared term, 2 or 2! With squared term A1: Correct expansion ie contents of bracket correct	M1A1
	$= 3 + \frac{x}{6} + \frac{x^2}{12} - \frac{3}{8} \times \frac{x^2}{81}$	M1 Remove all brackets	M1
	$(y=)3 + \frac{1}{6}x + \frac{17}{216}x^2$	M1: Combine x^2 terms and obtain a 3 term quadratic A1: Correct expression with or without $y = \dots$	M1A1

By implicit differentiation: For the first 4 marks (rest as first method)

$$y^2 = 8 + e^x$$

$$\text{M1A1 } 2y \frac{dy}{dx} = e^x \quad \text{M1A1 } 2 \frac{dy}{dx} + 2y \frac{d^2y}{dx^2} = e^x$$

Leave
blank

- $$\cos 6\theta = 32\cos^6\theta - 48\cos^4\theta + 18\cos^2\theta - 1 \quad (5)$$

- (b) Hence solve for $0 \leq \theta \leq \frac{\pi}{2}$

$$64\cos^6\theta - 96\cos^4\theta + 36\cos^2\theta - 3 = 0$$

giving your answers as exact multiples of π . (5)



Question Number	Scheme		Marks
4.(a)	$\cos 6\theta = \operatorname{Re}[(\cos \theta + i \sin \theta)^6]$	Ignore any imaginary parts included in their expansion	
	$(\cos \theta + i \sin \theta)^6 = c^6 + 6c^5 i s + 15c^4 i^2 s^2 + 20c^3 i^3 s^3 + 15c^2 i^4 s^4 + 6c i^5 s^5 + i^6 s^6$		M1
	Attempt to expand correctly or only show real terms (May be implied) Often seen with powers of i simplified. If $i s^n$ seen, but becomes $i^n s^n$ (oe) later, deduct the final A mark of (a) even if no further errors.		
	$\cos 6\theta = c^6 - 15c^4 s^2 + 15c^2 s^4 - s^6$	M1: Attempt to identify real parts. These 2 M marks may be awarded together	M1A1
		A1: Correct expression	
	$= c^6 - 15c^4(1 - c^2) + 15c^2(1 - c^2)^2 - (1 - c^2)^3$		M1
	Correct use of $s^2 = 1 - c^2$ in all their sine terms		
	$\cos 6\theta = c^6 - 15c^4 + 15c^6 + 15c^2(1 - 2c^2 + c^4) - (1 - 3c^2 + 3c^4 - c^6)$		
	$\cos 6\theta = 32 \cos^6 \theta - 48 \cos^4 \theta + 18 \cos^2 \theta - 1$ *	($\cos 6\theta$ must be seen somewhere)	A1cso
			(5)
(b)	$64 \cos^6 \theta - 96 \cos^4 \theta + 36 \cos^2 \theta - 3 = 0$ $\Rightarrow 2 \cos 6\theta - 1 = 0 \therefore \cos 6\theta = \frac{1}{2}$ or <u>0.5</u>	M1: Uses part (a) to obtain an equation in $\cos 6\theta$	M1A1
		A1: Correct underlined equation	
	$\cos 6\theta = \frac{1}{2} \Rightarrow (6\theta =) \frac{\pi}{3}, \frac{5\pi}{3}, \frac{7\pi}{3}$		
	$\theta = \frac{\pi}{18}, \frac{5\pi}{18}, \frac{7\pi}{18}$	M1: Valid attempt to solve $\cos 6\theta = k$, $-1 \leq k \leq 1$ leading to $\theta = \dots$ Can be degrees A1 2 correct answers A1 3 rd correct answer with no extras within the range, ignore extras outside the range. Must be radians Answers in degrees or decimal answers score A0A0	M1 A1A1
			(5)
			Total 10

Leave
blank

5. (a) Find the general solution of the differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + 10y = 27e^{-x} \quad (6)$$

- (b) Find the particular solution that satisfies $y = 0$ and $\frac{dy}{dx} = 0$ when $x = 0$

(6)



Question Number	Scheme		Marks
5.(a)	$\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + 10y = 27e^{-x}$		
	$m^2 + 2m + 10 (=0) \Rightarrow m = \dots$	Form and solve the aux equation	M1
	$m = -1 \pm 3i$		A1
	$(y =) e^{-x} (A \cos 3x + B \sin 3x)$ or $(y =) A e^{(-1+3i)x} + B e^{(-1-3i)x}$	y= not needed May be seen with θ instead of x	A1
	$y = k e^{-x}, y' = -k e^{-x}, y'' = k e^{-x}$	$y = k e^{-x}$ and attempt to differentiate twice	M1
	$e^{-x} (k - 2k + 10k) = 27e^{-x} \Rightarrow k = 3$		A1
	$y = e^{-x} (A \cos 3x + B \sin 3x + 3)$ or $y = A e^{(-1+3i)x} + B e^{(-1-3i)x} + 3e^{-x}$	Must be x and have $y = \dots$ Ignore any attempts to change the second form. (But see note at end about marking (b)) ft, so $y =$ their CF + their PI	B1ft (NB A1 on e-PEN)
			(6)
(b)	$x = 0, y = 0 \Rightarrow A = (-3)$	Uses $x = 0, y = 0$ in an attempt to find A	M1
	$y' = -e^{-x} (A \cos 3x + B \sin 3x + 3) + e^{-x} (3B \cos 3x - 3A \sin 3x)$	M1: Attempt to differentiate using the product rule, with A or their value of A A1: Correct derivative, with A or their value of A	M1A1
	$x = 0, y' = 0 \Rightarrow B = 0$	M1: Uses $x = 0, y' = 0$ and their value of A in an attempt to find B A1: $B = 0$	M1A1
	$y = e^{-x} (3 - 3 \cos 3x)$ oe	cao and cso	A1 (6)
			Total 12
	Alternative for (b) using	$y = A e^{(-1+3i)x} + B e^{(-1-3i)x} + 3e^{-x}$	
	$x = 0, y = 0$ to get an equation in A and B	May come from the real part of their derivative instead	M1
	$y' = (-1 + 3i) A e^{(-1+3i)x} + (-1 - 3i) B e^{(-1-3i)x} - 3e^{-x}$	M1: Attempt differentiation using chain rule A1: Correct differentiation	M1A1
	$x = 0, y' = 0 \Rightarrow -A - B - 3 = 0$ from real parts and $3A - 3B = 0$ from imaginary parts So $A = B = -\frac{3}{2}$	M1: Uses $x = 0, y' = 0$ and equates imaginary parts to obtain a second equation for A and B and attempts to solve their equations A1: $A = B = -\frac{3}{2}$	M1A1
	$y = -\frac{3}{2} e^{(-1+3i)x} - \frac{3}{2} e^{(-1-3i)x} + 3e^{-x}$	A1: Ignore any attempts to change.	A1

Some may change the second form in (a) before proceeding to (b). If their changed form is correct, all marks for (b) are available; if their changed form is incorrect only M marks are available.

6. The transformation T from the z -plane, where $z = x + iy$, to the w -plane, where $w = u + iv$, is given by

$$w = \frac{4(1-i)z - 8i}{2(-1+i)z - i}, \quad z \neq \frac{1}{4} - \frac{1}{4}i$$

(a) Show that

$$w = \frac{ax^2 + bxi + c}{16x^2 + 1}$$

where a , b and c are real constants to be found.

(6)

(b) Hence show that the circle C has equation

$$(u - 3)^2 + v^2 = k^2$$

where k is a constant to be found.

(4)



Question Number	Scheme		Marks
6.(a)	$w = \frac{4(1-i)z - 8i}{2(i-1)z - i}$		
	Method 1.....:Substituting $z = x + xi$	at the start	
	$w = \frac{4(1-i)(x + xi) - 8i}{2(i-1)(x + xi) - i}$	Substitutes for z	M1
	$w = \frac{4(x + xi - xi + x) - 8i}{2(xi - x - x - xi) - i}$	M1: Attempt to expand numerator and denominator	M1A1
		A1: Correct expression	
	$\frac{8i - 8x}{4x + i} \cdot \frac{4x - i}{4x - i}$	M1: Multiplies numerator and denominator by the conjugate of their denom. No expansion needed A1: Uses correct conjugate (not ft)	M1A1
	$= \frac{-32x^2 + 40xi + 8}{16x^2 + 1}$	cso Award only if final answer is correct and follows correct working	B1
	NB: The B mark appears first on e-PEN but will be awarded last		
			(6)
	Method 2: if they proceed without $y = x$ (substitution may happen anywhere in the working)		
	$w = \frac{(1-i)z - 8i}{2(-1+i) - i} = \frac{4(1-i)(x + iy) - 8i}{2(-1+i)(x + iy) - i}$	Substitutes for z	M1
	$= \frac{4(1-i)x + 4(1-i)iy - 8i}{2(-1+i)x + 2(-1+i)iy - i}$	Attempt to expand numerator and denominator	M1
	$= \frac{4x + 4y + (4y - 4x - 8)i}{-2x - 2y + (2x - 2y - 1)i}$	Correct expression	A1
	$= \frac{4x + 4y + (4y - 4x - 8)i}{-2x - 2y + (2x - 2y - 1)i} \times \frac{-2x - 2y - (2x - 2y - 1)i}{-2x - 2y - (2x - 2y - 1)i}$ M1: Multiplies numerator and denominator by the conjugate of their denom. No expansion needed. A1: Uses correct conjugate. (not ft)		M1A1
	$= \frac{-16x^2 - 16y^2 + 12y - 12x + 8 + (20x + 20y)i}{8x^2 + 8y^2 - 4x + 4y + 1}$		
	$= \frac{-32x^2 + 40xi + 8}{16x^2 + 1}$	cso Correct answer using $y = x$ Award only if final answer is correct and follows correct working	B1
	NB: The B mark appears first on e-PEN but will be awarded last		(6)

Question Number	Scheme		Marks
NB: The order of awarding the marks here has changed from the original mark scheme, but they must still be entered on e-PEN by their descriptors (M or A)			
6(b)	$u = \frac{-32x^2 + 8}{16x^2 + 1}, \quad v = \frac{40x}{16x^2 + 1}$	Identifies u and v (Real and imaginary parts) May be implied by their working and may be in terms of x and y .	M1 1st M mark on e-PEN
	$\left(\frac{8 - 32x^2}{16x^2 + 1} - 3\right)^2 + \left(\frac{40x}{16x^2 + 1}\right)^2$	Substitutes for their u and v in the given equation. May be in terms of x and y . May have a, b, c instead of their values (which may be chosen by the candidate if unable to do (a))	dM1 2nd M mark on e-PEN
	$= \left(\frac{8 - 32x^2 - 48x^2 - 3}{16x^2 + 1}\right)^2 + \left(\frac{40x}{16x^2 + 1}\right)^2$		
	$= \frac{(5 - 80x^2)^2}{(16x^2 + 1)^2} + \frac{1600x^2}{(16x^2 + 1)^2}$		
	$= \frac{6400x^4 + 800x^2 + 25}{(16x^2 + 1)^2}$	Combines to form a single correct fraction	A1 1 st A mark on e-PEN
	$= \frac{25(16x^2 + 1)^2}{(16x^2 + 1)^2} = 25$	$k = 5$ or $k^2 = 25$ may (but need not) be seen explicitly	A1 2 nd A mark on e-PEN
			(4)
			Total 10

Leave
blank

- $$x \frac{dy}{dx} + y = 2x^4 y^4 \quad (\text{I})$$

$$\frac{dv}{dx} - \frac{3v}{x} = -6x^3 \quad (\text{II})$$

(5)

- (6)



Question Number	Scheme		Marks
	Way 1		
7.(a)	$v = y^{-3} \Rightarrow \frac{dv}{dy} = -3y^{-4}$	Correct derivative	B1
	$\frac{dy}{dx} = \frac{dy}{dv} \frac{dv}{dx} = -\frac{y^4}{3} \frac{dv}{dx}$ Or $-3y^{-4} \frac{dy}{dx} x - 3y^{-3} = -6x^4$	M1: Correct use of the chain rule A1: Correct equation	M1A1
	$-\frac{y^4}{3} \frac{dv}{dx} x + y = 2x^4 y^4$		
	$-\frac{y^4}{3} \frac{dv}{dx} x + y = 2x^4 y^4 \Rightarrow \frac{dv}{dx} - \frac{3v}{x} = -6x^3$	dM1: Substitutes to obtain an equation in v and x . A1: Correct completion with no errors seen	dM1A1
	Way 2		
	$y = v^{-\frac{1}{3}} \Rightarrow \frac{dy}{dv} = -\frac{1}{3} v^{-\frac{4}{3}}$	Correct derivative	B1
	$\frac{dy}{dx} = \frac{dy}{dv} \frac{dv}{dx} = -\frac{1}{3} v^{-\frac{4}{3}} \frac{dv}{dx}$	M1: Correct use of the chain rule A1: Correct equation	M1A1
	$-\frac{v^{-\frac{4}{3}}}{3} \frac{dv}{dx} x + v^{-\frac{1}{3}} = 2x^4 v^{-\frac{4}{3}}$	dM1: Substitutes to obtain an equation in v and x .	dM1
	$-\frac{v^{-\frac{4}{3}}}{3} \frac{dv}{dx} x + v^{-\frac{1}{3}} = 2x^4 v^{-\frac{4}{3}} \Rightarrow \frac{dv}{dx} - \frac{3v}{x} = -6x^3$	A1: Correct completion with no errors seen	A1
	Way 3 (Working in reverse)		
	$v = y^{-3} \Rightarrow \frac{dv}{dy} = -3y^{-4}$	B1: Correct derivative	B1
	$\frac{dv}{dx} = \frac{dv}{dy} \frac{dy}{dx} = -3y^{-4} \frac{dy}{dx}$	M1: Correct use of chain rule A1: Correct expression for dv/dx	M1A1
	$-3y^{-4} \frac{dy}{dx} - \frac{3y^{-3}}{x} = -6x^3$	M1: Substitutes correctly for $\frac{dv}{dx}$ and v in equation (II) to obtain a D.E. in terms of x and y only. A1: Correct completion to obtain equation (I) with no errors seen	dM1A1

Question Number	Scheme		Marks
7(b)	$I = e^{\int \frac{-3}{x} dx} = e^{-3 \ln x} = \frac{1}{x^3}$	M1: $e^{\int \frac{-3}{x} dx}$ and attempt integration. If not correct, $\ln x$ must be seen.	M1A1
		A1: $\frac{1}{x^3}$	
	$\frac{v}{x^3} = \int -6 dx = -6x(+c)$	M1: $v \times \text{their } I = \int -6x^3 \times \text{their } I dx$	dM1A1
		A1: Correct equation with or without $+c$	
	$\frac{1}{y^3 x^3} = -6x + c \Rightarrow y^3 = \dots$	Include the constant, then substitute for y and attempt to rearrange to $y^3 = \dots$ or $y = \dots$ with the constant treated correctly	ddM1 dep on both M marks of (b)
	$y^3 = \frac{1}{cx^3 - 6x^4}$	Or equivalent	A1 (6) Total 11

8.




Figure 1

Figure 1 shows a sketch of part of the curve C with polar equation

$$r = 1 + \tan \theta, \quad 0 \leq \theta < \frac{\pi}{2}$$

The tangent to the curve C at the point P is perpendicular to the initial line.

(a) Find the polar coordinates of the point P .

(5)

The point Q lies on the curve C , where $\theta = \frac{\pi}{3}$

The shaded region R is bounded by OP , OQ and the curve C , as shown in Figure 1

(b) Find the exact area of R , giving your answer in the form

$$\frac{1}{2} (\ln p + \sqrt{q} + r)$$

where p , q and r are integers to be found.

(7)



Question Number	Scheme		Marks
	$r = 1 + \tan \theta$		
8.(a)	$x = r \cos \theta \Rightarrow x = (1 + \tan \theta) \cos \theta$	States or implies $x = r \cos \theta$	M1
	$x = \cos \theta + \sin \theta, \frac{dx}{d\theta} = \cos \theta - \sin \theta$	M1: Attempt to differentiate $x = r \cos \theta$ or $x = r \sin \theta$ A1: Correct derivative	M1A1
Alt for the 2 diff marks	$\frac{dx}{d\theta} = \sec^2 \theta \cos \theta + (1 + \tan \theta)(-\sin \theta)$	M1: Attempt to differentiate using product rule (dep on first M1) A1: correct (unsimplified) differentiation	
	$\frac{dx}{d\theta} = 0 \Rightarrow \tan \theta = 1 \Rightarrow \theta = \dots$	Set their derivative = 0 and attempt to solve for θ (Dependent on second M mark above)	dM1
	$\theta = \frac{\pi}{4}, r = 2$	Both	A1
	NB: Use of $x = r \sin \theta$ can score M0M1A0M1A0 max		(5)
(b)	$\int r^2 d\theta = \int (1 + \tan \theta)^2 d\theta$	Use of $\int r^2 d\theta$ and $r = 1 + \tan \theta$ No limits needed	M1
	$(1 + \tan \theta)^2 = 1 + 2 \tan \theta + \tan^2 \theta$ $= 1 + 2 \tan \theta + \sec^2 \theta - 1$	Expands and uses the correct identity	M1
	$\int (2 \tan \theta + \sec^2 \theta) d\theta$	Correct expression Need not be simplified, no limits needed.	A1
	$\left[2 \ln \sec \theta + \tan \theta \right]_{\left(\frac{\pi}{4}\right)}^{\left(\frac{\pi}{3}\right)}$	M1: Attempt to integrate – at least one trig term integrated. Dependent on the second M mark A1: Correct integration. Need not be simplified or include limits.	dM1A1
	$R = \frac{1}{2} \left\{ \left(2 \ln \sec \frac{\pi}{3} + \tan \frac{\pi}{3} \right) - \left(2 \ln \sec \frac{\pi}{4} + \tan \frac{\pi}{4} \right) \right\}$	Substitutes $\frac{\pi}{3}$ and their $\frac{\pi}{4}$ and subtracts (Dependent on 2 previous method marks in (b))	dM1
	$R = \frac{1}{2} \{ \ln 2 + \sqrt{3} - 1 \}$	Cao and cso	A1
			(7)
			Total 12