

Write your name here

Surname	Other names
---------	-------------

**Pearson Edexcel**  
**International**  
**Advanced Level**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

# Core Mathematics C12

## Advanced Subsidiary

Tuesday 13 January 2015 – Morning  
**Time: 2 hours 30 minutes**

Paper Reference  
**WMA01/01**

**You must have:**  
Mathematical Formulae and Statistical Tables (Blue)

Total Marks

--

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided – *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

### Information

- The total mark for this paper is 125.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P45057A

©2015 Pearson Education Ltd.

5/5/5/



**PEARSON**

Leave blank

1. Simplify the following expressions fully.

(a)  $(x^6)^{\frac{1}{3}}$

(1)

(b)  $\sqrt{2}(x^3) \div \sqrt{\frac{32}{x^2}}$

(2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



January 2015  
International A Level WMA01/01 Core Mathematics C12  
Mark Scheme

Question Number	Scheme	Marks
<b>1.</b>	(a) $x^2$	B1 [1]
	(b) $\frac{1}{4}x^4$ or $\frac{1}{2^2}x^4$ or $0.25x^4$	B1, B1 [2]
		<b>3 marks</b>
	<b>Notes</b>	
<p>(a) <b>B1:</b> This answer only</p> <p>(b) <b>B1:</b> For <math>\frac{1}{4}x^k</math> <b>as final answer</b>, <math>k</math> can even be 0. Also accept <math>\frac{1}{2^2}</math> for B1 but <math>2^{-2}</math> is not simplified and is B0</p> <p><b>B1:</b> for <math>x</math> to power 4 (independent mark) so <math>kx^4</math> with <math>k</math> a constant (could even be 1) <b>as final answer</b></p> <p>n.b. Can score B0B1 or B1B0 or B0B0 or B1B1</p> <p><b>Mark the final answer on this question</b></p> <p>Also note : Candidates who misread question as <math>\sqrt{2x^3} \div \sqrt{\frac{32}{x^2}}</math> should get <math>\frac{1}{4}x^{\frac{5}{2}}</math> This is awarded B1B0</p> <p><b>Special case:</b> The answer <math>\left(\frac{1}{\sqrt{2}}x\right)^4</math> is awarded B0 B1 as <math>x</math> may be in a bracket with power 4 outside.</p>		

2.




Figure 1

Figure 1 shows a sketch of part of the graph of  $y = \frac{12}{\sqrt{(x^2 - 2)}}$ ,  $x \geq 2$

The table below gives values of  $y$  rounded to 3 decimal places.

$x$	2	5	8	11
$y$	8.485	2.502	1.524	1.100

- (a) Use the trapezium rule with all the values of  $y$  from the table to find an approximate value, to 2 decimal places, for

$$\int_2^{11} \frac{12}{\sqrt{(x^2 - 2)}} dx$$

**(4)**

- (b) Use your answer to part (a) to estimate a value for

$$\int_2^{11} \left( 1 + \frac{6}{\sqrt{(x^2 - 2)}} \right) dx$$

**(3)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks										
<p>2.</p> <p>(a)</p>	<table border="1" data-bbox="304 241 1259 322"> <tr> <td><math>x</math></td> <td>2</td> <td>5</td> <td>8</td> <td>11</td> </tr> <tr> <td><math>y</math></td> <td>8.485</td> <td>2.502</td> <td>1.524</td> <td>1.100</td> </tr> </table> <p>State <math>h = 3</math>, or use of <math>\frac{1}{2} \times 3</math></p> <p><math>\{ 8.485 + 1.100 + 2( 2.502 + 1.524 ) \}</math> For structure of <math>\{ \dots \}</math></p> <p><math>\frac{1}{2} \times 3 \times \{ 17.637 \}</math> (= 26.4555) = awrt 26.46</p> <p>(b) Adds 9+.....</p> <p>half of <b>their answer from (a)</b> seen ( allow use of half of 26.4555)</p> <p>So required estimate = <math>9 + 13.23 = 22.23</math></p>	$x$	2	5	8	11	$y$	8.485	2.502	1.524	1.100	<p>B1 aef</p> <p>M1A1</p> <p>A1</p> <p>[4]</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>
$x$	2	5	8	11								
$y$	8.485	2.502	1.524	1.100								
<b>7 marks</b>												
<p>(b)</p>	<p>Way 2: <b>Begins again with trapezium rule</b></p> <table border="1" data-bbox="304 857 1259 938"> <tr> <td><math>x</math></td> <td>2</td> <td>5</td> <td>8</td> <td>11</td> </tr> <tr> <td><math>y</math></td> <td>5.2425</td> <td>2.251</td> <td>1.762</td> <td>1.550</td> </tr> </table> <p>Uses <math>\frac{1}{2} \times 3 \times \{ 5.2425 + 1.550 + 2(2.251 + 1.762) \}</math></p> <p>= 22.23</p>	$x$	2	5	8	11	$y$	5.2425	2.251	1.762	1.550	<p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>
$x$	2	5	8	11								
$y$	5.2425	2.251	1.762	1.550								
<p><b>Notes</b></p>												
<p>(a) <b>B1:</b> for using <math>\frac{1}{2} \times 3</math> or 1.5 or equivalent or just states <math>h = 3</math></p> <p><b>M1:</b> requires the correct <math>\{ \dots \}</math> bracket structure. It needs the first bracket to contain first <math>y</math> value plus last <math>y</math> value and the second bracket to be multiplied by 2 and to be the summation of the remaining <math>y</math> values in the table with no additional values. If the only mistake is a copying error or is to omit one value from 2nd bracket this may be regarded as a slip and the M mark can be allowed (An extra repeated term forfeits the M mark however). M0 if values used in brackets are <math>x</math> values instead of <math>y</math> values</p> <p><b>A1: for the completely correct bracket</b> <math>\{ \dots \}</math></p> <p><b>A1:</b> for answer which rounds to 26.46 after attempt at trapezium rule</p> <p>NB: Separate trapezia may be used: B1 for 1.5, M1 for <math>\frac{1}{2} h(a + b)</math> used 2 or 3 times (and A1 if it is all correct ) Then A1 for 26.46.</p> <p>Special case: Bracketing mistake <math>1.5 \times (8.485 + 1.1) + 2(2.502 + 1.524)</math> scores B1 M1 A0 A0 unless the final answer implies that the calculation has been done correctly (then full marks can be given).</p> <p>(b) <b>Way 1:</b></p> <p><b>M1:</b> Adds Area of Rectangle = <math>1 \times 9</math> or <math>\int 1 dx = [x]_2^{11}</math> to their “13.23” or to their “26.46” or to their “52.92”</p> <p><b>M1:</b> Half answer to part (a) seen</p> <p><b>A1:</b> Accept awrt 22.23</p> <p><b>Or Way 2: (If they begin again with a trapezium rule)</b></p> <p><b>M1:</b> for correct table <b>M1:</b> for correct use of trapezium rule</p> <p><b>A1:</b> awrt 22.23</p>												

3.




Figure 2

Figure 2 shows a sketch of part of the curve with equation  $y = f(x)$ .

The curve crosses the coordinate axes at the points  $(2.5, 0)$  and  $(0, 9)$ , has a stationary point at  $(1, 11)$ , and has an asymptote  $y = 3$



On **separate** diagrams, sketch the curve with equation

(a)  $y = 3f(x)$  (3)

(b)  $y = f(-x)$  (3)

On each diagram show clearly the coordinates of the points of intersection of the curve with the two coordinate axes, the coordinates of the stationary point, and the equation of the asymptote.



Question Number	Scheme	Marks
3.	<p>(a)</p>  <div style="border: 1px solid black; padding: 5px; margin: 10px 0;"> <p>Shape- similar to before but with indication of <b>stretch</b> in y direction by at least <b>one correct</b> from the three traits:                      y intercept, (0, 27)                      maximum point (1, 33)                      or asymptote indicated at 9</p> <p>Intercept (0,27), max (1,33) and x intercept (2.5,0) <b>all three of these seen</b></p> </div> <p>(b)</p>  <div style="border: 1px solid black; padding: 5px; margin: 10px 0;"> <p>Shape (reflection in y axis)</p> <p>(-1,11), (0,9) and (-2.5,0) seen</p> <p><math>y = 3</math> (must be equation)</p> </div>	<p>B1</p> <p>B1</p> <p>B1</p> <p>[3]</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>[3]</p>
<b>6 marks</b>		
	<b>Notes</b>	

(a)**B1**: Correct shape with curve crossing  $x$  axis and one label correct from the three listed (i.e. a correct new  $y$  value). Condone “slight” imperfections in the curvature of the sketches.

**B1**: All three specified labels given to indicate the three new point positions. Do not need coordinates if clearly labelled on the axes. Accept 27 and accept 2.5 and even allow (27, 0) and (0, 2.5) on  $y$  and  $x$  axes respectively.

**B1**: Equation of asymptote correct (asymptote on figure takes precedence) Asymptote does not need to be drawn dotted.

(b)**B1**: Correct shape (maximum in 2<sup>nd</sup> quadrant, intercept on negative  $x$  - axis and approaches asymptote for large positive  $x$ ) Condone “slight” imperfections in the curvature of the sketches.

**B1**: All three specified labels given to indicate the three new point positions. Accept 9 and accept -2.5 and even allow (9, 0) and (0, -2.5) on  $y$  and  $x$  axes respectively.

**B1**: Equation of asymptote correct (asymptote on figure takes precedence) Do not award this mark if they merely copy the original graph.

If there is no sketch – the maximum mark in part (a) is B0B1B1 and in part (b) is B0B1B0 so 3/6

Special case: Stretch in  $y$  direction of scale factor  $1/3$ . If there is a graph of the correct shape with (0,3), (1, 11/3), (2.5,0) and asymptote  $y = 1$  then award B0B0B1





Question Number	Scheme	Marks
4.	<p>(a) <math display="block">\left(2 + \frac{x}{4}\right)^{10} = 2^{10} + \binom{10}{1}2^9 \cdot \left(\frac{x}{4}\right) + \binom{10}{2}2^8 \cdot \left(\frac{x}{4}\right)^2 + \binom{10}{3}2^7 \cdot \left(\frac{x}{4}\right)^3 \dots</math> <math display="block">= 1024 + 1280x + 720x^2 + 240x^3 \dots</math></p> <p>(b) State or Use <math>x = 0.1</math>                      Estimate = <math>1024 + 1280 \times 0.1 + 720 \times (0.1)^2 + 240 \times (0.1)^3 \dots</math>                      = 1159.44 or 1159.440 or 1159 or 1159.4 (after correct working)</p>	<p>M1 B1, A1 A1 [4]</p> <p>B1 M1 A1 [3]</p> <p><b>7 marks</b></p>
	<b>Notes</b>	
<p>(a) <b>M1:</b> The <b>method</b> mark is awarded for an attempt at Binomial to get one or more of the terms in <math>x</math> – need <b>correct</b> binomial coefficient multiplied by the correct power of <math>x</math>. Ignore bracket errors or errors (or omissions) in powers of 2 or 4 or bracket errors. Accept any notation for <math>{}^{10}C_1</math>, <math>{}^{10}C_2</math> and <math>{}^{10}C_3</math>, e.g. <math>\binom{10}{1}</math>, <math>\binom{10}{2}</math> and <math>\binom{10}{3}</math> (unsimplified) or 10, 45 and 120 from Pascal's triangle This mark may be given if no working is shown, but any of the terms including <math>x</math> is correct.  <b>B1:</b> must be simplified to 1024 ( writing just <math>2^{10}</math> is <b>B0</b> ). <b>If miscopied later then isw</b>  <b>A1:</b> is cao and is for <b>two correct</b> from <math>1280x</math>, <math>720x^2</math> and <math>240x^3</math>  <b>A1:</b> is c.a.o and is for all of <math>1280x</math>, <math>720x^2</math> and <math>240x^3</math> correct (ignore extra terms) if divided by 2 or 4 then isw                      Allow terms given separately without + signs and with commas. Ignore extra terms. Ignore subsequent work once correct answer is seen in simplified form.                      N.B. If the series is given <b>in Descending Order</b> the first M mark may be awarded and if the whole expansion is given (all 11 terms) then full marks is possible.                      (b) <b>B1:</b> States or Uses <math>x = 0.1</math>  <b>M1:</b> Uses their solution of <math>\frac{x}{4} = 0.025</math> substituted in to their series expansion – If no equation stated could see evidence of use of 0.1 or 0.01 (not 0.025) substituted consistently for example  <b>A1:</b> This is cao and must follow M1.                      NB 1159.45 or 1159.44533 is A0 (used <math>2.025^{10}</math>) But correct working followed by an answer 1159 or 1159.4 can be awarded A1</p>		



Question Number	Scheme	Marks
5. (a)	$S_n = a + (a+d) + (a+2d) + \dots + (a+(n-1)d)$	M1
	$S_n = (a+(n-1)d) + (a+(n-2)d) + \dots + (a+d) + a$	M1
	$2S_n = (2a+(n-1)d) + (2a+(n-1)d) + \dots + (2a+(n-1)d)$	M1
	$S_n = \frac{n}{2}[2a+(n-1)d]$ * See notes below for those who use triangle numbers in their proof	A1*
(b)	Uses either $\frac{n}{2}(2 \times a + (n-1)7)$ or $\frac{n}{2}(a+497)$ or $7 \times \sum_{i=1}^{71} i$	M1
	i.e. $\frac{71}{2}(2 \times 7 + 70 \times 7)$ or $\frac{72}{2}(2 \times 0 + 71 \times 7)$ or $\frac{71}{2}(7 + 497)$ or $7 \times \frac{71}{2}(72)$	A1
	= 17892	A1
		<b>[4]</b>
		<b>[3]</b>
		<b>7 marks</b>
<b>Notes</b>		

(a) **M1:** List terms including **at least first two and a last term which may be  $a + nd$  or  $a + (n - 1)d$  or  $L$**   
**M1:** List terms in reverse including **at least their last term ( or correct last term) and finally their first term**  
**M1:** The LHS should be  $2S$ . The RHS must follow from at least two terms correctly matching in the addition and should include at least two terms which are each **correctly**  $\{2a + (n - 1)d\}$  or  $(a + L)$  **or should be**  $n\{2a + (n - 1)d\}$  or  $n(a + L)$   
**A1:** Need some indication of at least three terms being added (i.e at least three terms and their pairs listed with terms correctly matching or three additions seen) and also need to achieve final answer with no errors and if  $L$  was used need to state that  $L = a + (n - 1)d$   
 NB: Some candidates use a variation of  

$$\sum_{r=1}^n (a + (r - 1)d) = \sum_{r=1}^n a + d \sum_{r=1}^n (r - 1) = na + d \frac{n}{2}(n + 1) - dn$$
 or  $na + d \frac{(n - 1)}{2}(n)$   
 And conclude that  $S_n = \frac{n}{2}[2a + (n - 1)d]$ . This gains the full 4 marks M1M1M1A1, but must be completely correct.

(b) **M1:** Uses correct formula (with their  $a$  and  $n$ ) with  $d = 7$  or with last term correct  
**A1:** Uses consistent and correct  $a$  and  $n$   
**A1:** Correct answer



Question Number	Scheme	Marks
6.	(a) Use or state $2 \log_4(2x+3) = \log_4(2x+3)^2$	M1
	Use or state $\log_4 4 = 1$ or $4^1 = 4$	M1
	Use or state $\log_4 x + \log_4(2x-1) = \log_4 x(2x-1)$ or $\log_4(2x+3)^2 - \log_4 x = \log_4 \frac{(2x+3)^2}{x}$ etc	M1
	$(2x+3)^2 = 4x(2x-1)$ or equivalent including correct rational equations	A1
	Then $4x^2 + 12x + 9 = 8x^2 - 4x$ and so $4x^2 - 16x - 9 = 0$ *	A1* [5]
(b) $(2x+1)(2x-9) = 0$ so $x =$ (or use other method e.g formula or completion of square)	M1	
$x = (-\frac{1}{2}$ or $\frac{9}{2}$	A1 [2]	
		<b>7 marks</b>
<b>Notes</b>		

(a) M1: Uses power law for logs

M1: Connects 1 with 4 correctly

M1: Uses addition ( or subtraction) law correctly

e.g.  $\log_4 x + \log_4(2x-1) = \log_4 x(2x-1)$  or  $\log_4(2x+3)^2 - \log_4 x = \log_4 \frac{(2x+3)^2}{x}$  or

$\log_4(2x+3)^2 - \log_4 x - \log_4(2x-1) = \log_4 \frac{(2x+3)^2}{x(2x-1)}$  or even  $\log_4 x + \log_4 4 = \log_4 4x$  or

$\log_4(2x-1) + \log_4 4 = \log_4 4(2x-1)$  or  $\log_4(2x-1) + \log_4 4 + \log_4 x = \log_4 4x(2x-1)$  etc...

A1: Correct equation (unsimplified) after correct work. e.g.  $\frac{(2x+3)^2}{x(2x-1)} = 4$

A1: Obtains printed answer correctly (This is a given answer so needs previous A mark to have been awarded and needs correct expansion)

Special case :

$$\log_4 (2x+3)^2 = 1 + \log_4 x(2x-1) \text{ so } \frac{\log_4 (2x+3)^2}{\log_4 x(2x-1)} = 1 \text{ so } \frac{4x^2 + 12x + 9}{2x^2 - x} = 4$$

This can have M1, M1, M1, A0, A0 so 3/5 losing accuracy because of the error in the second step.

(b) Some candidates who did not achieve marks in part (a) begin the log work again and make more progress here. Mark the better work. So credit for (a) may be given in (b). Credit for (b) should not be given in (a)

M1: Uses solution of their quadratic or of printed quadratic(see notes). This must be in part (b)

A1:  $x = 4.5$  and discards  $x = -0.5$  (any equivalent form) Giving  $x = -\frac{1}{2}, \frac{9}{2}$  is A0 This must be in part (b)

Leave blank

7. The circle *C* has equation

$$x^2 + y^2 + 10x - 6y + 18 = 0$$

Find

(a) the coordinates of the centre of *C*,

(2)

(b) the radius of *C*.

(2)

The circle *C* meets the line with equation  $x = -3$  at two points.

(c) Find the exact values for the *y* coordinates of these two points, giving your answers as fully simplified surds.

(4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question number	Scheme	Marks
<p>7 (a)</p> <p>(b)</p> <p>(c)</p>	<p>Obtain <math>(x \pm 5)^2</math> <b>and</b> <math>(y \pm 3)^2</math></p> <p>Centre is <math>(-5, 3)</math>.</p> <p>See <math>(x \pm 5)^2 + (y \pm 3)^2 = 16 (= r^2)</math> or <math>(r^2 = )"25"+"9"-18</math>  <math>r = 4</math></p> <p>Use <math>x = -3</math> in either form of equation of circle to obtain simplified quadratic in <math>y</math>                      e.g <math>x = -3 \Rightarrow (-3+5)^2 + (y-3)^2 = 16 \Rightarrow (y-3)^2 = 12</math>                      or <math>(-3)^2 + y^2 + 10 \times (-3) - 6y + 18 = 0 \Rightarrow y^2 - 6y - 3 = 0</math>                      solve resulting quadratic to give <math>y =</math>  <math display="block">y = 3 \pm 2\sqrt{3}</math></p>	<p>M1</p> <p>A1</p> <p>[2]</p> <p>M1</p> <p>A1</p> <p>[2]</p> <p>M1</p> <p>M1</p> <p>A1, A1</p> <p>[4]</p> <p><b>8 marks</b></p>
<p><u>Alternatives</u></p> <p>(a)</p> <p>OR</p> <p>(b)</p> <p>(c)</p>	<p><i>Method 2:</i> From <math>x^2 + y^2 + 2gx + 2fy + c = 0</math> centre is <math>(\pm g, \pm f)</math>                      Centre is <math>(-g, -f)</math> , <b>and so centre is</b> <math>(-5, 3)</math>.</p> <p><i>Method 3:</i> Use any value of <math>y</math> to give two points (<math>L</math> and <math>M</math>) on circle. <math>x</math> co-ordinate of mid point of <math>LM</math> is <math>"-5"</math> <b>and</b> Use any value of <math>x</math> to give two points (<math>P</math> and <math>Q</math>) on circle.  <math>y</math> co-ordinate of mid point of <math>PQ</math> is <math>"3"</math> (Centre – chord theorem) . <math>(-5, 3)</math> is                      M1A1</p> <p><i>Method 2:</i> Using <math>\sqrt{g^2 + f^2 - c}</math> or <math>(r^2 = )"25"+"9"-18</math> <math>r = 4</math></p> <p><i>Method 2:</i> Divide triangle <math>PTQ</math> and use Pythagoras with  <math>r^2 - (-3 - "-5")^2 = h^2</math>, then evaluate <math>"3 \pm h"</math> - then get <math>3 \pm 2\sqrt{3}</math></p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>M1 A1</p> <p>(2)</p> <p>M1</p> <p>M1 A1</p> <p>A1</p> <p>(4)</p>
<b>Notes</b>		



**Mark (a) and (b) together**

(a) **M1** as in scheme and can be implied by  $(\pm 5, \pm 3)$  **A1**: for correct centre and **(-5, 3) (without working) implies M1A1**

(b) **M1** for a complete and correct method leading to  $r^2 = "25" + "9" - 18$  or  $r = \sqrt{"25" + "9" - 18}$   
or for using equation of circle in  $(x \pm 5)^2 + (y \pm 3)^2 = k^2$  form to identify  $r = k$

N.B.  $r^2 = k$  or  $r = k^2$  is M0 Also - "25" - "9" - 18 is M0 and  $r^2 = "25" + "9"$  (without the 18) is M0

**A1**  $r = 4$  (only and not with  $r = -4$ ) Again **correct answer with no working implies M1A1**

**Special case:** if centre is given as **(5, -3) or (5, 3) or (-5, -3)** allow **M1A1** for  $r = 4$  worked correctly as  $(r^2 = )"25" + "9" - 18$  i.e if they obtain  $r = 4$  after sign error give final A1 (So M1A0M1A1)

(c) **M1** For substituting  $x = -3$  into an equation for the circle and attempt to simplify to 3 term quadratic or to  $(y - a)^2 = b$

**M1** For attempting to solve their quadratic (following usual rules - see notes)

**A1, A1** Answers must be given as surds - A1 for each correct answer. To earn both A marks, answers must be simplified.

8. A sequence is defined by

$$\begin{aligned} u_1 &= k \\ u_{n+1} &= 3u_n - 12, \quad n \geq 1 \end{aligned}$$

where  $k$  is a constant.

(a) Write down fully simplified expressions for  $u_2$ ,  $u_3$  and  $u_4$  in terms of  $k$ . (4)

Given that  $u_4 = 15$

(b) find the value of  $k$ , (2)

(c) find  $\sum_{i=1}^4 u_i$ , giving an exact numerical answer. (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
8.	$u_2 = 3k - 12, u_3 = 3(u_2) - 12$ $u_2 = 3k - 12, u_3 = 9k - 48$ $u_4 = 3(9k - 48) - 12 = 27k - 156 \quad (\text{ft their } u_3) \quad .$	M1
		A1
		M1 A1ft [4]
		M1
(b)	$27k - 156 = 15 \text{ so } k =$ $k = 6\frac{1}{3} \text{ or } \frac{19}{3} \text{ or } 6.33 \text{ (3sf)}$	A1 [2]
(c)	$\sum_{i=1}^4 u_i = 6\frac{1}{3} + 7 + 9 + 15 \text{ or } \sum_{i=1}^4 u_i = k + 3k - 12 + 9k - 48 + 27k - 156$ $= 40k - 216, = 37\frac{1}{3} \text{ or } \frac{112}{3}$	M1 A1ft, A1cao [3]
		<b>9 marks</b>

**Notes**

(a) **M1:** Attempt to use formula twice to find  $u_2$  and  $u_3$   
**A1:** two correct simplified answers  
**M1:** Attempt again to find  $u_4$   
**A1ft:** 4<sup>th</sup> term correct and simplified - follow through their  $u_3$

(b) **M1:** Put their 4<sup>th</sup> term ( not 5<sup>th</sup>) equal to 15 and attempt to find  $k =$   
**A1:** accept any correct fraction or decimal answer (allow 6.33 or better here)

(c) **M1:** Uses 1<sup>st</sup> term and their following 3 terms **with plus signs** (either numerical or in terms of  $k$ ). Must be using terms from iteration and not formula for an AP or GP. May make a copying slip.  
**A1ft:** for  $40k - 216$  or follow through on their  $k$  so check  $40k - 216$  for their  $k$   
**A1:** obtains  $37\frac{1}{3}$  (must be exact) if exact answer given, then isw

Those who use 6.3 will obtain 36 They should have M1A1ftA0 – should have used exact  $k$  to give exact answer here.

Those who use 6.33 will obtain 37.2 This should have M1A1ftA0 – should have used exact  $k$  to give exact answer here.

Those who use 6.333 will obtain 37.32 This should have M1A1ftA0 – should have used exact  $k$  to give exact answer here.

6.3333 will obtain 37.332 This should have M1A1ftA0 – should have used exact  $k$  to give exact answer here.

6.33333 will obtain 37.3332 etc All these answers should have M1A1ftA0 – should have used exact  $k$  to give exact answer here. Etc

**Special case:** Those who use  $k = 6$  will obtain  $6 + 6 + 6 + 6 = 24$  This is M1 A0 A0 in part (c) – as over simplified

9.




Figure 3

In Figure 3, the points  $A$  and  $B$  are the centres of the circles  $C_1$  and  $C_2$  respectively. The circle  $C_1$  has radius 10 cm and the circle  $C_2$  has radius 5 cm. The circles intersect at the points  $X$  and  $Y$ , as shown in the figure.

Given that the distance between the centres of the circles is 12 cm,

- (a) calculate the size of the acute angle  $XAB$ , giving your answer in radians to 3 significant figures, (2)
- (b) find the area of the major sector of circle  $C_1$ , shown shaded in Figure 3, (3)
- (c) find the area of the kite  $AYBX$ . (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
9. (a)	$5^2 = 10^2 + 12^2 - 2 \times 10 \times 12 \cos \angle XAB$ , or $\cos \angle XAB = \frac{10^2 + 12^2 - 5^2}{2 \times 10 \times 12}$ or $\frac{219}{240}$ or 0.9125 or $\frac{73}{80}$ $\angle XAB = 0.421$ or $0.134\pi$	M1 A1 [2]
(b)	Area of sector is $\frac{1}{2}r^2\theta = \frac{1}{2} \times 10^2 \times \theta$ Area of major sector is $\frac{1}{2} \times r^2 (2\pi - 2 \times "0.421")$ or $\pi \times r^2 - \frac{1}{2} \times r^2 \times 2 \times "0.421")$ = 272	M1 M1 A1 [3]
(c)	area of triangle $AXB = \frac{1}{2}10 \times 12 \times \sin XAB$ <b>Way 2:</b> Find angle $XBA$ and hence area $XY$ area of kite = $2 \times \text{triangle}AXB$ Area of kite = area of $XY + \text{Area } XAY$ = awrt 49      = $37.298 + 11.76 = 49$ <b>Way 3:</b> Finds length $XY$ by cosine rule or elementary trigonometry (8.173) Uses area of kite = $\frac{1}{2} "8.173" \times 12$ = awrt 49	M1 dM1 A1 [3] M1 dM1 A1 [3] <b>8 marks</b>
	<b>Notes</b>	

(a) **M1:** Uses cosine rule – must be a correct statement, allow statement  $5^2 = 10^2 + 12^2 - 2 \times 10 \times 12 \cos \angle XAB$   
**A1:** accept awrt 0.421 (answers in degrees gain M1 A0). Also 0.42 is A0

(b) **M1:** Uses area formula with  $r = 10$  and any angle in radians. If they use degrees they must use the formula  $\frac{\theta}{360} \times \pi 10^2$   
**M1:** Finds angle in major sector ft their angle from (a) and uses sector formula **or** subtracts minor area from circle (allow work in degrees) Must use  $(2\pi - 2 \times "0.421")$  but  $r$  may be 5 instead of 10 for this mark  
**A1:** Accept awrt 272 (may reach this using degrees)

(c) Way 1: **M1:** Finds area of triangle  $AXB$ , using 10, 12 and their angle  $XAB$   
**dM1:** Doubles area of triangle  $AXB$   
 Way 2: **M1:** Finds angle  $XBA$  (0.958..) by valid method (cosine rule) (**NOT 90 –  $XAB$** ) and hence area  $XY = \frac{1}{2} 5 \times 5 \times \sin 1.9163$   
**dM1:** Adds areas of triangles  $XY$  and  $XAY$  (37.298 and 11.76)  
 Way 3: **M1:** Finds length  $XY$  by cosine rule or elementary trigonometry (8.173)  
**dM1:** Uses area of kite =  $\frac{1}{2} "8.173" \times 12$

**For each method A1:** awrt 49- do not need units



Question Number	Scheme	Marks
<p><b>10.</b></p> <p>(a)</p> <p>(b)</p>	<p style="text-align: center;"><math>f(x) =</math></p> <p><math>6x^3 + ax^2 + bx - 5</math></p> <p>Attempts <math>f(\pm 1)</math> <b>or</b> Attempts <math>f(\pm \frac{1}{2})</math> Or Use long division as far as remainder*</p> <p>Obtains <math>6(-1)^3 + a(-1)^2 + b(-1) - 5 = 0</math> or <math>-6 + a - b - 5 = 0</math> or <math>a - b = 11</math> or equivalent</p> <p>Obtains <math>6(\frac{1}{2})^3 + a(\frac{1}{2})^2 + b(\frac{1}{2}) - 5 = -15</math> or <math>\frac{6}{8} + \frac{a}{4} + \frac{b}{2} - 5 = -15</math> or <math>a + 2b = -43</math> or equivalent</p> <p>Solve simultaneous equations to obtain <math>a = -7</math> and <math>b = -18</math></p> <p><math>6x^3 + ax^2 + bx - 5 = (x+1)(6x^2 + \dots x + \dots)</math></p> <p><math>6x^3 - 7x^2 - 18x - 5 = (x+1)(6x^2 - 13x - 5)</math></p> <p><math>(6x^2 - 13x - 5) = (ax+b)(cx+d)</math> where <math>ac = "6"</math> and <math>bd = "\pm 5"</math></p> <p><math>= (x+1)(2x-5)(3x+1)</math></p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>M1 A1</p> <p style="text-align: right;"><b>[5]</b></p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p style="text-align: right;"><b>[4]</b></p> <p style="text-align: right;"><b>9 marks</b></p>
	<p><b>Notes</b></p>	

(a) **M1**: Using remainder theorem: As on scheme. One of these is sufficient do not need to equate to 0 and to -15

\*Using Long division: need at least  $6x^2 + (a-6)x + \dots$  as quotient, and get as far as remainder **or for the other**

**division reaches**  $3x^2 + (\frac{a+3}{2})x + \dots$  as quotient, and get as far as remainder .

**A1**: Any equivalent form \*e.g.  $-11 - b + a = 0$  (using remainder after division) The mark is earned for  $a - b = 11$

even if “=0” not explicitly seen

**A1**: Any equivalent form \*e.g.  $-5 + \frac{b}{2} + \frac{a+3}{4} = -15$  (using remainder after division) Must be accurate but may be

unsimplified. NB Using 15 instead of -15 is A0

**M1**: Solves their **linear** equations to obtain  $a$  or  $b$

**A1**: Both  $a$  and  $b$  correct. Correct answers without working can earn M1A1.

(b) **M1**: Recognises  $(x+1)$  is factor and obtains quadratic expression with correct first term by any method.

Use of  $(x-1)$  is M0. NB Starting with  $(x+1)(2x-1)(ax+b)$  is also M0

**A1**: Correct quadratic  $(6x^2 - 13x - 5)$

**M1**: Attempt to factorise quadratic where  $ac = "6"$  and  $bd = "\pm 5"$

**A1**: any correct combination e.g.  $= 2(x+1)(x - \frac{5}{2})(3x+1)$  or  $= 6(x+1)(x - \frac{5}{2})(x + \frac{1}{3})$  etc... (on one line)

**Following a correct value for  $a$  and for  $b$ :**

**They may just write the factorised answer down.**

For a correct answer this is M1A1M1A1

For  $= (x+1)(x - 2.5)(x + \frac{1}{3})$  award M1A0M1A0

For **correct answer** following **incorrect quadratic** give M1 A0 M1 A0 – fortuitous

If the correct answer follows incorrect  $a$  and  $b$ , it is fortuitous and again M1A0M1A0 should be given.



11.




Figure 4

Figure 4 shows a sketch of the curve  $C$  with equation  $y = \sin(x - 60^\circ)$ ,  $-360^\circ \leq x \leq 360^\circ$

(a) Write down the exact coordinates of the points at which  $C$  meets the two coordinate axes. (3)

(b) Solve, for  $-360^\circ \leq x \leq 360^\circ$ ,

$$4 \sin(x - 60^\circ) = \sqrt{6} - \sqrt{2}$$

showing each stage of your working. (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>11</b> (a)</p> <p>(b)</p>	<p><math>\left(0, -\frac{\sqrt{3}}{2}\right)</math></p> <p>and <math>(60^\circ, 0)</math> and <math>(240^\circ, 0)</math> and <math>(-120^\circ, 0)</math> and <math>(-300^\circ, 0)</math></p> <p><math>\sin(x - 60^\circ) = \frac{\sqrt{6} - \sqrt{2}}{4}</math> (= 0.2588 )</p> <p><math>x - 60^\circ = 15^\circ</math> (or <math>165^\circ</math> or <math>-195^\circ</math> or <math>-345^\circ</math>) or 0.262 or <math>\frac{\pi}{12}</math> radians</p> <p>So <math>x = 75^\circ</math> or <math>225^\circ</math> or <math>-135^\circ</math> or <math>-285^\circ</math> (allow awrt)</p>	<p>B1</p> <p>B1 B1 [3]</p> <p>M1</p> <p>A1</p> <p>M1 A1 A1 [5]</p> <p><b>8 marks</b></p>

**Notes**

(a) **B1** : Correct exact y intercept (not decimal) – allow on the diagram or in the text. Allow  $y = -\frac{\sqrt{3}}{2}$

**B1** for 2 correct x intercepts then **third B1** for all 4 correct x intercepts ( may or may not be given as coordinates – may be given on graph) Must be in degrees. (Extra answers in the range lose the **third B1**)

(b) **M1**: Divides by 4 first giving correct statement  $\sin(x - 60^\circ) = \frac{\sqrt{6} - \sqrt{2}}{4}$  but  $(x - 60^\circ) = \frac{\sqrt{6} - \sqrt{2}}{4}$  is **M0** and  $\sin x - \sin 60^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$  is also **M0** and  $\sin(x - 60^\circ) = \frac{\sqrt{4}}{4}$  is **M0** if not preceded by correct statement

**A1**: Obtains  $15^\circ$  (or  $165^\circ$  or  $-195^\circ$  or  $-345^\circ$ )

**M1**: Adds  $60^\circ$  to their previous answer which should have been in degrees and obtained by using inverse sine

**A1**: Two correct answers second **A1**: All four correct answers Extra answers outside range are ignored. Lose final A mark for extra wrong answers in the range.

**If they approximate too early allow awrt answers given for full marks.** (e.g. 75.01 etc)

**Answers in mixture, degrees and radians:** Allow first M A1 only so M1A1M0A0A0 for 60.262 for example



Question Number	Scheme	Marks
<p>12.(a)</p> <p>(b)</p> <p>(c)</p>	<p>Uses <math>275000 \times (1.1)^5</math> or finds £442890.25 or uses <math>275000 \times (1.1)^4</math> or finds £402627.50</p> <p>Finds both of the above and subtracts to give £40 262.75 and concludes approx. £40300*</p> <p>Or</p> <p>Uses <math>275000 \times (1.1)^5 - 275000 \times (1.1)^4</math>, = awrt 40260 = 40300 (3sf) *</p> <p>Puts <math>275000 \times (1.1)^{n-1} &gt; 1000000</math> or <math>275000 \times (1.1)^{n-1} = 1000000</math></p> <p><math>(1.1)^{n-1} &gt; \frac{1000000}{275000}</math> (or <math>\frac{40}{11}</math> or 3.63 or 3.64). Or</p> <p><math>(1.1)^{n-1} = \frac{1000000}{275000}</math> (or <math>\frac{40}{11}</math> or 3.63 or 3.64)</p> <p><math>n-1 &gt; \frac{\log(\frac{40}{11})}{\log 1.1}</math> or <math>n-1 = \frac{\log(\frac{40}{11})}{\log 1.1}</math></p> <p>(<math>n &gt; 14.5</math> or <math>n &gt; 14.6</math> or <math>n = 15</math>) so the year is 2030</p> <p>Uses <math>S = \frac{275000(1.1^n - 1)}{1.1 - 1}</math> or uses <math>S = \frac{275000(1 - 1.1^n)}{1 - 1.1}</math></p> <p>Uses <math>n = 11</math> in formula</p> <p>Awrt £5 096 100</p> <p>Or: adds 11 terms £275000 + 302500 + 332750 + 366025 + 402627.5 + 442890.25 + 487179.275 + 535897.2025 + 589486.9228 + 648435.615 + 713279.1765 = awrt 5096100 (see notes below)</p>	<p>M1</p> <p>M1 A1*</p> <p>[3]</p> <p>M1 M1, A1*</p> <p>[3]</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[4]</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[3]</p> <p><b>10</b></p> <p><b>marks</b></p>
	<p><b>Notes</b></p>	

(a) **M1**: for correct expression for profit in 2021 **or** in 2020, by any method (including subtracting the sums  $S_{n+1} - S_n$ ) to give a term

**M1**: for finding both correct expressions and subtracting

**A1**: answers wrt£442900 and wrt£402600 subtracted **or** wrt£40260 obtained **then rounded to** £40300

(answer given)

(b) **M1**: Correct inequality – or allow equality . N.B.  $250000 \times (1.1)^n$  or  $302500 \times (1.1)^{n-2}$  on LHS are also correct.

**M1**: Division – isw if initial fraction is correct. Not dependent on previous mark. It could follow wrong combination of  $a$  and  $n$  for example, which would give M0 M1

**M1**: Correct use of logs to give  $n$  or  $n-1 > \frac{\log(k)}{\log 1.1}$  or  $\log_{1.1} k$  after  $(1.1)^{n-1} > k$  Allow equality for this mark

( 3.63 is truncated value of  $\frac{40}{11}$  and 3.64 is rounded value – allow either of these if used in place of fraction)

**A1**: 2030 is required . If inequalities are used and errors are seen, then this mark is A0 (even for 2030) (Trial and improvement or listing can have full marks for the correct answer, need to see both 14<sup>th</sup> and 15<sup>th</sup> term – otherwise zero)

Special case: If  $n$  is used instead of  $n-1$  and they reach 2029 then mark profile is likely to be M0 M1 M1 A0

**unless they recover to the correct answer when full marks may be earned**

**If an equals sign is used throughout and then correct answer is obtained allow 4/4**

**Special case: Uses Sum formula** – Can earn M0 M0 M1 A1 for “correct work”

Uses  $S = \frac{275000(1.1^n - 1)}{1.1 - 1} > 1000000$  (M0)  $1.1^n > 1 + \frac{1000000}{275000}$  (M0)  $n > \frac{\log(15/11)}{\log 1.1}$  (M1)  $n > 3.254\dots$  so 2019 (A1)

Using this method with errors can earn M0M0M1A0 for proceeding from  $1.1^n > k$  with  $k > 0$  to  $n > \frac{\log(k)}{\log 1.1}$

(c) **M1**: Correct  $a$  and  $r$  but  $n$  may be wrong

**A1**: Correct use of formula with  $n = 11$

**A1**: awrt £5 096 100 (again – this answer implies all 3 marks)

Or M1: adds 11 terms ( mostly correct)

A1: lists 11 correct terms £275000 + £302500 + £332750 + £366025 + £402627.5 + £442890.25 + £487179.275 + £535897.2025 + £589486.9228 + £648435.615 + £713279.1765

A1: correct answer = awrt £5096100 (this implies two previous marks)

13. The curve  $C$  has equation

$$y = 3x^2 - 4x + 2$$

The line  $l_1$  is the normal to the curve  $C$  at the point  $P(1, 1)$

(a) Show that  $l_1$  has equation

$$x + 2y - 3 = 0$$

(5)

The line  $l_1$  meets curve  $C$  again at the point  $Q$ .

(b) By solving simultaneous equations, determine the coordinates of the point  $Q$ .

(4)

Another line  $l_2$  has equation  $kx + 2y - 3 = 0$ , where  $k$  is a constant.

(c) Show that the line  $l_2$  meets the curve  $C$  once only when

$$k^2 - 16k + 40 = 0$$

(4)

(d) Find the two exact values of  $k$  for which  $l_2$  is a tangent to  $C$ .

(2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>13.</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p><math>y = 3x^2 - 4x + 2</math></p> <p><math>\frac{dy}{dx} = 6x - 4 + \{0\}</math></p> <p>At (1, 1) gradient of curve is 2 and so gradient of normal is <math>-\frac{1}{2}</math></p> <p><math>\therefore (y - 1) = -\frac{1}{2}(x - 1)</math> and so <math>x + 2y - 3 = 0^*</math></p> <p>Eliminate <math>x</math> or <math>y</math> to give <math>2(3x^2 - 4x + 2) + x - 3 = 0</math> or <math>y = 3(3 - 2y)^2 - 4(3 - 2y) + 2</math></p> <p>Solve three term quadratic e.g. <math>6x^2 - 7x + 1 = 0</math> or <math>12y^2 - 29y + 17 = 0</math> to give <math>x =</math> or <math>y</math></p> <p><math>=</math></p> <p><math>x = \frac{1}{6}</math> or <math>y = 1\frac{5}{12}</math></p> <p><b>Both</b> <math>x = \frac{1}{6}</math> and <math>y = 1\frac{5}{12}</math> i.e. <math>(\frac{1}{6}, 1\frac{5}{12})</math> or <math>(0.17, 1.42)</math> { Ignore (1, 1) listed as well }</p> <p>When this line meets the curve <math>2(3x^2 - 4x + 2) + kx - 3 = 0</math></p> <p>So <math>6x^2 + (k - 8)x + 1 = 0</math></p> <p><b>Uses condition for equal roots</b> "<math>b^2 = 4ac</math>" on their three term quadratic to get expression in <math>k</math></p> <p>So obtain <math>(k - 8)^2 = 24</math> i.e. <math>k^2 - 16k + 40 = 0</math> *</p> <p>If they use <b>gradient of tangent</b> to do part (c) see the end of the notes below*.</p> <p>Solve the given quadratic or their quadratic by formula or completion of the square to give</p> <p><math>k = 8 \pm \sqrt{24}</math> or <math>8 \pm 2\sqrt{6}</math> or <math>\frac{16 \pm \sqrt{96}}{2}</math> .....</p>	<p>M1A1</p> <p>M1</p> <p>M1 A1*</p> <p><b>[5]</b></p> <p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p><b>[4]</b></p> <p>M1</p> <p>dM1</p> <p>ddM1</p> <p>A1 *</p> <p><b>[4]</b></p> <p>M1A1</p> <p><b>[2]</b></p> <p><b>15 marks</b></p>
	<p><b>Notes</b></p>	

- (a) **M1:** Evidence of differentiation, so  $x^n \rightarrow x^{n-1}$  at least once  
**A1:** Both terms correct  
**M1:** Substitutes  $x = 1$  into their derivative and uses perpendicular property  
**M1:** Correct method for Linear equation, using (1,1) and their changed gradient  
**A1:** Should conclude with printed answer (this answer is given in the question)
- (b) **M1:** May make sign slips in their algebra; {e.g. substitute  $3 + 2y$ } – does not need to be simplified so isw.  
 But putting  $3(3 - 2y)^2 - 4(3 - 2y) + 2 = 0$  instead of  $= y$  is M0  
**M1:** Solve three term quadratic to give one of the two variables  
**A1:** One Correct coordinate – accept any equivalent  
**A1:** Both correct – any equivalent form. Allow decimals if correct awrt (0.17, 1.42) ( ignore (1,1) given as well)
- (c) **M1:** Eliminate  $y$  (condone small copying errors)  
**dm1:** Collect into 3 term quadratic in  $x$  or identifies “ $a$ ”, “ $b$ ” and “ $c$ ” clearly (may be implied by later work).  
**ddM1:** Uses condition “ $b^2 = 4ac$ ” on quadratic in  $x$  (dependent on both previous M marks)  
 NB **M0** for  $b^2 > 4ac$  or  $b^2 \geq 4ac$  or  $b^2 < 4ac$  or  $b^2 \leq 4ac$   
**A1:** Need  $(k - 8)^2 = 24$  or equivalent before stating printed answer  
 \*Alternative method for part (c)  
**M1:** Use gradient of line = gradient of curve so “ $6x - 4 = -\frac{k}{2}$ ”  
**M1:** Find  $x = \frac{2}{3} - \frac{k}{12}$  and use line equation to get  $y = \frac{3}{2} - \frac{1}{3}k + \frac{k^2}{24}$  (these equations do not need to be simplified)  
**M1:** Find  $x = \frac{2}{3} - \frac{k}{12}$  and use curve equation to get  $y = \frac{2}{3} + \frac{k^2}{48}$  ( these equations do not need to be simplified)  
**A1:** Puts two correct expressions for  $y$  equal and obtains printed answer without error.
- (d) **M1:** Solve by formula or completion of the square to give  $k =$  (Attempt at factorization is M0)  
**A1:** Correct answer – should be one of the forms given in the main scheme or equivalent exact form  
 Answers only with no working 2 marks (exact and correct) or 0 marks ( approximate or wrong)





Question Number	Scheme	Marks
14. (i)	Way 1: Use $\frac{\sin x}{\cos x} = \tan x$ to give $\tan x =$ Way2: complete method to find $\sin x =$ or $\cos x =$	M1
	$\tan x = -\frac{7}{3}$ or $\sin x = \pm \frac{7}{\sqrt{58}}$ or $\cos x = \pm \frac{3}{\sqrt{58}}$	A1
	So $x = 113.2, 293.2$	M1 A1 [4]
(ii)	$10\cos^2 \theta + \cos \theta = 11(1 - \cos^2 \theta) - 9$	M1
	Solves their three term quadratic “ $21\cos^2 \theta + \cos \theta - 2 = 0$ ” to give $\cos \theta = \dots$	M1
	So ( $\cos \theta =$ ) $-\frac{1}{3}$ or $\frac{2}{7}$	A1
	$\theta = 1.91, 4.37, 1.28$ or $5.00$ (allow 5 instead of 5.00)	M1 A1 A1 [6]
		<b>10 marks</b>
<b>Notes</b>		

(i) **M1: (Way 1)** Attempts to use  $\frac{\sin x}{\cos x} = \tan x$  (there may be a sign error or may omit  $x$  and write  $\tan =$ )

**(Way 2)**  $3\sin x = -7\cos x$  so  $9\sin^2 x = 49\cos^2 x$  and uses  $\sin^2 x + \cos^2 x = 1$  to find  $\sin x =$  or  $\cos x =$

**A1:** must be  $\tan x = -\frac{7}{3}$  (way 1) or allow  $\sin x = \pm \frac{7}{\sqrt{58}}$  or  $\cos x = \pm \frac{3}{\sqrt{58}}$  (way 2). Ignore  $\cos x = 0$  as extra answer.

**M1:** One correct angle in degrees in range – so need either 113.2 or 293.2 in most cases

But If they had  $\tan x = -\frac{3}{7}$ , then obtaining 156.8 or 336.8 is equivalent work and gains M1

If however they had  $\tan x = +\frac{7}{3}$ , then obtaining an answer in the range is not equivalent work – so is M0

**A1:** These two answers - accept awrt 113.2 and 293.2 Extra answers in range – lose this mark

Working in radians gives a maximum of M1A1M0A0

(ii) **M1:** Replaces  $\sin^2 \theta$  by  $(1 - \cos^2 \theta)$

**M1:** Collects terms and solves their three term quadratic by usual methods (see notes)

**A1:** Both correct answers needed, but isw if one then rejected. Allow awrt -0.333 and 0.286

**M1** Uses inverse cosine to obtain at least two correct answers for their values of cosine (check with calculator if they have followed wrong values)

**A1: Any** two completely correct answers (allow awrt)

**A1:** All four correct (awrt) Allow  $0.608\pi, 1.39\pi, 0.408\pi,$  or  $1.59\pi$

Extra answers outside range – ignore **Extra answers in the range** – lose final mark. Inaccurate answers to 3sf lose final A mark

Answers in degrees lose final two marks

So two of awrt 73, 287, 109 (or 109.5), 251 (or 250.5) would earn M1A0A0

15.




Figure 5

Figure 5 shows a sketch of part of the curve  $C$  with equation

$$y = x^3 + 10x^{\frac{3}{2}} + kx, \quad x \geq 0$$

where  $k$  is a constant.

(a) Find  $\frac{dy}{dx}$

**(2)**

The point  $P$  on the curve  $C$  is a minimum turning point.  
Given that the  $x$  coordinate of  $P$  is 4

(b) show that  $k = -78$

**(2)**

The line through  $P$  parallel to the  $x$ -axis cuts the  $y$ -axis at the point  $N$ .

The finite region  $R$ , shown shaded in Figure 5, is bounded by  $C$ , the  $y$ -axis and  $PN$ .

(c) Use integration to find the area of  $R$ .

**(7)**


---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Question Number	Scheme	Marks
<p><b>15.</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<p><math>y = x^3 + 10x^{\frac{3}{2}} + kx</math></p> <p><math>\frac{dy}{dx} = 3x^2 + 10 \times \frac{3}{2}x^{\frac{1}{2}} + k</math></p> <p>Substitutes <math>x = 4</math> and <math>\frac{dy}{dx} = 0</math> to give <math>3(4)^2 + 15(4)^{\frac{1}{2}} + k = 0 \Rightarrow k = -78</math> *</p> <p>When <math>x = 4</math>, <math>y = -168</math> ( see this stated – or see rectangle has height 168)</p> <p><math>\int x^3 + 10x^{\frac{3}{2}} - 78x (+168)dx = \frac{1}{4}x^4 + \frac{10}{\frac{5}{2}}x^{\frac{5}{2}} - \frac{78}{2}x^2 (+168x + c)</math></p> <p>Use limits 0 and 4 to give <math>\pm 432</math> or if <math>168x</math> included to give <math>\pm 240</math></p> <p>Rectangle area is <math>4 \times "168"</math> (= 672) or see <math>168x</math> in integrated answer with limits</p> <p>So <math>R</math> has area "672 – 432" or see <math>+168</math> in original integrand = 240</p>	<p>M1 A1 [2]</p> <p>M1 A1* [2]</p> <p>B1 M1 A1 dB1 M1 M1 A1 [7]</p> <p><b>11 marks</b></p>
	<p><b>Notes</b></p>	
<p>(a)</p> <p>(b)</p> <p>(c)</p>	<p><b>M1:</b> Fractional power dealt with correctly so becomes <math>\frac{3}{2}x^{\frac{1}{2}}</math> ( may be implied by simplification to 15)</p> <p><b>A1:</b> All terms correct, may not be simplified</p> <p><b>M1: Substitutes</b> <math>x = 4</math> and <math>\frac{dy}{dx} = 0</math> Must see <math>3(4)^2 + 15(4)^{\frac{1}{2}} + k = 0</math> or <math>48 + 30 + k = 0</math></p> <p><b>*A1:</b> This is a printed answer so all must be correct in the working and conclusion <math>k = -78</math> is needed.</p> <p><b>B1:</b> Substitute into <math>y =</math> to find <math>y</math> (This may appear anywhere in the answer)</p> <p><b>M1:</b> Attempt to integrate so at least one power increases</p> <p><b>A1:</b> Accept unsimplified correct answer and allow with or without their <math>+168x</math>, or even with their <math>-168x</math></p> <p><b>dB1:</b> Use limit 4 to give 432 but may be implied by later answer 240- needs to follow M1A1 for integration</p> <p><b>M1:</b> Calculates rectangle area (may be by integration). Must be rectangle and not triangle area</p> <p><b>M1:</b> Subtracts (either way round) numerical areas – should be (+) – (+) or (-) - (-) (subtraction may be in their original integral but penalize wrong sign here eg <math>-168x</math> instead of <math>+168x</math>) (Again use of triangle is M0)</p> <p><b>A1:</b> 240 only (Can recover from -240 to 240)</p> <p>Common error: If <math>168x</math> (instead of 168) is integrated this may only gain a maximum of B1 M1 A1 dB1 (for seeing 432 calculated if integrals are separated) M0 M0 A0 4/7</p>	